Biology of the Tropical Catfish (Family: Clariidae) with Special Emphasis on its Suitability for Culture (Including a Bibliography of the Clariidae and Related Topics)

D. Clay

Biological Station,
St. Andrews, N.B., EOG 2XO

December 1977

Fisheries & Marine Service Manuscript Report No.1458

Fisheries and Marine Service

Manuscript Reports

These reports contain scientific and technical information that represents an important contribution to existing knowledge but which for some reason may not be appropriate for primary scientific (i.e. *Journal*) publication. They differ from Technical Reports in terms of subject scope and potential audience: Manuscript Reports deal primarily with national or regional problems and distribution is generally restricted to institutions or individuals located in particular regions of Canada. No restriction is placed on subject matter and the series reflects the broad interests and policies of the Fisheries and Marine Service, namely, fisheries management, technology and development, ocean sciences and aquatic environments relevant to Canada.

Manuscript Reports may be cited as full publications. The correct citation appears above the abstract of each report. Each report will be abstracted by *Aquatic Sciences and Fisheries Abstracts* and will be indexed annually in the Service's index to scientific and technical publications.

Numbers 1-900 in this series were issued as Manuscript Reports (Biological Series) of the Biological Board of Canada, and subsequent to 1937 when the name of the Board was changed by Act of Parliament, as Manuscript Reports (Biological Series) of the Fisheries Research Board of Canada. Numbers 901-1425 were issued as Manuscript Reports of the Fisheries Research Board of Canada. The series name was changed with report number 1426.

Details on the availability of Manuscript Reports in hard copy may be obtained from the issuing establishment indicated on the front cover.

Service des pêches et des sciences de la mer

Manuscrits

Ces rapports contiennent des renseignements scientifiques et techniques qui constituent une contribution importante aux connaissances actuelles mais qui, pour une raison ou pour une autre, ne semblent pas appropriés pour la publication dans un journal scientifique. Ils se distinguent des Rapports techniques par la portée du sujet et le lecteur visé; en effet, ils s'attachent principalement à des problèmes d'ordre national ou régional et la distribution en est généralement limitée aux organismes et aux personnes de régions particulières du Canada. Il n'y a aucune restriction quant au sujet; de fait, la série reflète la vaste gamme des intérêts et des politiques du Service des pêches et de la mer, notamment gestion des pêches; techniques et développement, sciences océaniques et environnements aquatiques, au Canada.

Les Manuscrits peuvent être considérés comme des publications complètes. Le titre exact paraît au haut du résumé de chaque rapport, qui sera publié dans la revue *Aquatic Sciences and Fisheries Abstracts* et qui figuera dans l'index annuel des publications scientifiques et techniques du Service.

Les numéros de 1 à 900 de cette série ont été publiés à titre de manuscrits (Série biologique) de l'Office de biologie du Canada, et après le changement de la désignation de cet organisme par décret du Parlement, en 1937, ont été classés en tant que manuscrits (Série biologique) de l'Office des recherches sur les pêcheries du Canada. Les numéros allant de 901 à 1425 ont été publiés à titre de manuscrits de l'Office des recherches sur les pêcheries du Canada. Le nom de la série a été changé à partir du rapport numéro 1426.

La page couverture porte le nom de l'établissement auteur où l'on peut se procurer les rapports sous couverture cartonnée.

•Cover design by Christine Rusk

Fisheries and Marine Service
Manuscript Report 1458

December 1977

BIOLOGY OF THE TROPICAL CATFISH (FAMILY: CLARIIDAE) WITH SPECIAL EMPHASIS ON ITS SUITABILITY FOR CULTURE (INCLUDING A BIBLIOGRAPHY OF THE CLARIIDAE AND RELATED TOPICS)

by

D. Clay
Marine Fish Division
Department of Fisheries and Environment
Bedford Institute of Oceanography
Dartmouth, Nova Scotia B2Y 4A2

This is the fourth Manuscript Report in this series from the Biological Station, St. Andrews, N.B.

© Minister of Supply and Services Canada 1977
Cat. no. Fs 97-4/1458 ISSN 0701-7618

TABLE OF CONTENTS

	Page
Abstract/Résumé	v
Introduction	1
Commercial Importance of the Clariidae	3
Anatomy	7
Respiration	9
Habitat and Distribution	10
Reproduction	11
Feeding and Digestion	15
Growth and Production	19
Acknowledgments	21
Bibliography	23

PREFACE

This review and the work entailed in compiling this report were carried out while the author was employed by the Department of Agriculture, University of Rhodesia. At this time it was decided to prepare a detailed "Species Synopsis" on the family Clariidae. The present manuscript is a preliminary coverage of the topic and the aim of producing this interim publication is to obtain suggestions and criticism of all interested parties involved in work on the Clariidae.

The author would appreciate any suggestions, additions, or alterations that can be forwarded by workers in this field before July 1978.

ABSTRACT

Clay, D. 1977. Biology of the tropical catfish (Family: Clariidae) with special emphasis on its suitability for culture (including a bibliography of the Clariidae and related topics).. Fish. Mar. Serv. MS Rep. 1458. 68 p.

All available literature relevant to the culture of the Clariidae have been reviewed and evaluated. The present production of the catfish in inland African fisheries is estimated at over 20% by weight of the commercial catch. *Clarias* spp. is a particularly suitable fish for culture because of its dual respiratory system, its extreme hardiness, its broad feeding habits, and its very wide range of habitat. Emphasis is placed on reviewing the reproduction and digestion of these fish and their importance for culture. A need of local variety trials is shown to be necessary in order to investigate the potential for culture of the indigenous fishes of Africa and Asia.

Key words: Clariidae, culture, bibliography, growth, commercial fishing, reproduction, feeding, habitat, respiration, anatomy

RÉSUMÉ

Clay, D. 1977. Biology of the tropical catfish (Family: Clariidae) with special emphasis on its suitability for culture (including a bibliography of the Clariidae and related topics). Fish. Mar. Serv. MS Rep. 1458, 68 p.

Toute la documentation accessible sur l'élevage des Clariidae a été passée en revue et évaluée. On estime que la pêche actuelle de poissons-chats dans l'intérieur du continent africain représente plus de 20%, en poids, de la pêche commerciale. L'espèce *Clarias* convient particulièrement à l'élevage à cause de son système respiratoire double, de sa vigueur incomparable, de la diversité de ses habitudes alimentaires et de sa très grande faculté d'adaptation. L'accent est mis sur l'étude de leur système respiratoire et digestif, et sur leur importance piscicole. Il apparaît nécessaire de procéder à une sélection des variétés locales pour être en mesure d'évaluer les possibilités d'élevage des poissons indigènes d'Afrique et d'Asie.

INTRODUCTION

The tropical catfish is an air breathing, non-scaled, omnivorous, potamdromous member of the family Clariidae. The Indian catfish (*Clarias batrachus* Linnaeus), is found throughout the Indian sub-continent, the Far East, and has recently been introduced into both Hawaii and continental U.S.A. The Asian catfish (*Clarias macrocephalus* Gunther) is located in the Far East. The major commercial members of the African catfish, the subject of this review, are *Clarias gariepinus* Burchell, *Clarias lazera* Cuvier and Valenciennes, and *Clarias mossambicus* Peters. They are thought by some, including the author, to be geographical variations of the same species. Jubb (1967) has synonymized *C. gariepinus* and *C. mossambicus*.[1] These three species are distributed throughout Africa from the Orange River in the south to the Nile delta in the north (Fig. 1).



Fig. 1. World distribution of the family Clariidae (after Hora 1937; Menon 1951; Ogilvie and Goodrick 1968).

Although many authors have worked in the field of Clariidae taxonomy (Barnard 1943; Boulenger 1886, 1896, 1899, 1902, 1903, 1911; Deraniyagala 1958; Greenwood 1958, 1961; Gunther 1867; Jackson 1955, 1959; Jubb, 1964; Leriche 1901; Mann 1971; Poll 1942, 1943; Vaillant 1897) a detailed synthesis is now needed in light of present knowledge. Table 1 gives a brief résumé on the commercially important Clariidae with their respective scientific and common local names.

[1]Throughout this paper wherever *Clarias mossambicus* is discussed it will be referred to as *Clarias gariepinus*.

C. gariepinus can make up over 50% by weight of the commercial catch in some Central African fisheries (Harding 1960). More frequently, as in Lake Kariba, they make up less than 20% by weight of the catch (Anon. 1972, 1973, 1974; Harding 1964); even with this lower figure they must be considered one of the most important fish of African waters. Mwanza (1972) found *Clarias* sp. to be the only one of the three important commercial species to be unaffected by the 1968 drying out of Lake Chilwa in Malawi.

The tropical catfish has for many years suffered an unjust stigma in Africa as a trash fish, and recently as a nuisance fish in the U.S.A. (Idyll 1969). Jubb (1953) stated that all fish in farm dams were useful except the 'barbel' and furthermore that it was the only common fish that did not require a fish ladder to help populate new dams. More recent inquiry regarding this fish has sparked interest in its potential for aquaculture. Many authors (Brown 1955; Coureur and Spaas 1956; Van der Waal 1970) as well as the United Nations Development Program (UNDP) (Micha, pers. comm.) have begun investigations into various aspects of the biology of the tropical catfish, specifically for culture. Farming the Asian catfish in Thailand has developed over the last 15 yr into a large and growing industry (Sidthimunka 1972). The tropical catfish is very important in Africa and the Far East in both commercial catches and consumer preference. One consideration, important to the African situation, is that many tribes will not willingly eat the catfish because of taboos related to its scaleless body. Such feelings are also found amongst the Jewish population of Israel.

One of the earliest and most interesting accounts of *Clarias* sp. was written by the Rev. Boake in 1866. He reported that in the Mootoo Rajawelle swamp of [Sri Lanka] the vegetation was a "continuous sod ... firm enough to support the weight of natives ... cutting (the) long grass ... while on some of the firmer parts bullocks were to be seen grazing." Under this thick mat was 60 cm of water kept very muddy with "the consistency of thick pea soup" by the "large numbers of Hoongas *(Heteropneustes fossilis)* and Magooras *(Clarias teysmanni* Bleeker)." The local people of Ceylon used the knowledge that the tropical catfish were air breathers in order to catch them (Boake 1866). The fishermen cut several holes about 1 m diameter in the "sod" (described above). This was done at night when the fish were more active, and the frequent breathing at the surface indicated their presence. The next day the sod and peat in a large diameter ring around the hole were trampled into the swamp bottom, thus forming an enclosed ring of very muddy water with the fish trapped inside. The hole was then covered with long grass criss-crossed in such a way as to form a mesh-like net. When the fish began rising to breathe, it gave an advance warning by the air bubbles released before surfacing. The fishermen thus forwarned had little difficulty in catching the fish as it forced its head up

Table 1. A list of the major species of the Clariidae of commercial importance and their common names.

Scientific name	Synonym	Common name/location	Language
Clarias macrocephalus (Gunther)		Pla Duk Oui/Thailand	Thai
		Pla Duk/Thailand	Thai
Clarias batrachus (Linnaeus)	*C. magur*	Freshwater catfish/India	English
		Teysmann's spotted catfish/India	English
		Magur, Kagga, Ana meenu, Marpoo/India	?
		Magura/Sri Lanka	?
		Pla Duk Dan/Thailand	Thai
		Pla Duk/Thailand	Thai
		Ca tre, Trey andeng/Cambodia	?
		Ca tre trang/Vietnam	Vietnamese
		Alabbiyog, Kawatsi/Philippines	?
		Keli, Ikan keli/Malaysia	Malayan
		Leleh, Ikan kalang, Pintet, Tjepi/Indonesia	Indonesian
		Nga rheo/Kymer Republic	?
		Walking catfish/U.S.A.	English
Clarias teysmanni (Blecker)		Magoora/Sri Lanka	?
Clarias lazera (Cuvier)		Nigerian mudfish/Nigeria	English
		Kullune, Kulumni, Arungu, Tarwada/Nigeria	Hausa
		Mali/Uganda	?
		Nyai/Uganda	?
		Garmout/Sudan	?
		Harmut/Egypt	Arabic
		Armut/Israel	Arabic
Clarias senegalensis (Cuvier & Valenciennes)		Catfish/Ghana	English
Clarias ngamensis (Castelnau)		Ndombi/Zambia, Rhodesia	?
		Blunt-toothed barbel/Zambia	English
		Blunt-toothed catfish[a]/Zambia	English
Clarias theodorae (Weber)		Brown barbel, Snake barbel/Zambia	English
		Mulonfi/Zambia	?
		Mulonge/Zambia	?
		Minga/Zambia, Rhodesia	?
Clarias gariepinus (Burchell)	*C. mossambicus* (Peters)	Barber/Republic of South Africa	Afrikaans
		Barbel/R.S.A., Rhodesia	English
		African catfish/Rhodesia	English
		Mpongo, Mulonge, Muta/Zambia, Tanzania	?
		Sharp-toothed barbel/Rhodesia, Zambia	English
		Sharp-toothed catfish[a]/Rhodesia, Zambia	English
		Haramba/Rhodesia	Shona
		Ndombi/Zambia, Rhodesia	?
		Mubando/Zambia, Rhodesia	?
		Black barbel, Zambezi barbel/Zambia	English
		Mlamba/Malawi	?
		Mali (Twang)/Uganda	?
		Nyaki (Kasonzi)/Uganda	?
		Mandev/Mozambique	Shangan
		Slamue chilu/Mozambique, Rhodesia	?
		Gariep/R.S.A., Botswana	Hottentot

[a]Name now accepted as 'official' common name (Smith 1975).

through the woven grass net. This method resulted in about 11 fish per hour being caught.

The hardy nature of these fish, exploited by the people of [Sri Lanka] for so many years, is still a source of frequent reports. Tait (1967) found that in a mass fish kill in the Kafue River system of Zambia no catfish died, although *Tilapia* spp., *Hepsetus* sp., *Synodontis* sp., and *Schilbe* sp. all died of unknown causes.

COMMERCIAL IMPORTANCE OF THE CLARIIDAE

The Clariidae of Africa is one of the most important commercial families of freshwater fishes in traditional subsistence and modern commercial fisheries[2] (Tables 2 and 3). Many methods are used in fishing but the majority of catfish are landed by the many gillnet fisheries of African inland waters. Although Bowmaker (1973) found *C. gariepinus* not effectively caught in gillnets in a riverine estuary of Lake Kariba, Ratcliff (1972) found gillnets to be the major means of catching the catfish in Lake Malawi. There would appear to be great variation in the catchability of *Clarias* spp. in gillnets in various locations. Other methods seem to have less variability; seine nets appear to be universally ineffective in catching catfish; this may be due to the diving behaviour of the Clariidae. When cornered or disturbed, these fish dive into the mud allowing the toe rope of the seine to slide upward over their bodies. Williams (1960) gives records for the same river for a gillnet fishery and a seine-net fishery, the latter catches ½ to ¼ by weight of the former (Table 2).

Longlines are widely used in Tanzania (Bailey 1966; Soulsby 1960) and other African countries; in some lakes they are nearly 100% selective for *Clarias* spp. (Clay 1972, Mortimer 1959) because few other indigenous fish take still bait. Such knowledge would be a useful management tool to allow more complete cropping of this very productive fish. In small dams (as in Table 3) where the Clariidae can make up over 30% of the production, this very selective form of fishing requiring a low capital investment would allow an economic harvest of at least one-third of the total production. This would be especially important where gillnets could not offer a large enough economic return. During the annual spawning migrations in the Barotse flood plains of Zambia, traditional spear fishing is very widely used to harvest *Clarias* spp. (Bell-Cross 1974). Many of the very colourful traditional fishing methods such as basket fishing, trap fishing, and weir fishing are all effective in catching *Clarias* spp. although not totally selective for them.

[2]The purse-seine fishing of Lake Tanzania and Lake Kariba are notable exceptions to this.

As well as being an important fish in tonnage landed, it is one of the most sought after fish in Africa. Balon (1972) lists *C. gariepinus* in the top four highly preferred fish of Lake Kariba, while Mann (1964) found *Clarias* sp. liked by the African consumers for taste and high fat content. Often the likes or more particularly the dislikes are regional in nature and related to taboos dealing with scaleless fish.

The nutritive value of *C. gariepinus* has been investigated by Watanabe (1971). Water was found to be approximately 81% of the live weight, rising in winter with a corresponding drop in oil level which is generally about 2%. Protein levels are about 16% with ash levels running at 1%. Watanabe placed the loss after commercial gutting and removing the head at 44%, and after filleting at 66%.

Figure 2 gives some indication of the seasonal nature of catfish catches in gillnets.



Fig. 2. Seasonal variation in the catch per unit effort of *Clarias gariepinus* of the commercial gillnet fishery of Lake McIlwaine (data averaged over a 13-yr period). The numbers at the end of each of the four curves are gillnet stretched-mesh measurements.

Although the "seasons" do not necessarily coincide in all lakes, a cyclical pattern is evident in most. In the Mwenda estuary of Lake Kariba the catch per unit effort (CPU) of *Clarias* sp. ranged throughout the year from 0.3-1.6 kg 100 m^{-1} per night set (Bowmaker 1973). Great variation also occurs in catfish catches from different lakes. In contrast to the low yield of the Mwenda estuary of Lake Kariba, Lake Mweru, Zambia, has yielded approximately 6.3 kg 100 m^{-1} per night set (Bowmaker 1963), while the CPU in Lake McIlwaine, Rhodesia, has ranged from 2.5 to

Table 2. Importance of the family Clariidae in Africa as percent of the catches of commercially valuable fishes (species of Clariidae as per species code and fishing method as per method code).

Location	Fishing method	Percent catch by weight	Percent catch by numbers	Author
Lake Tanzania, Tanzania	L.F.	75 (C.spp.)		Soulsby (1960)
Malya Dam, Tanzania	C.G.N.F.	3 (C.g.)		Bailey (1966)
Lake Kitangiri, Tanzania	L.F. & S.F.	major[d] (C.g.)		Lockley (1960)
Lake Ruka, Tanzania	C.G.N.F. (B)	10-50 (C.spp.)		Mann (1964)
Barotse[a], Zambia	C.G.N.F.	4.5 (C.g.)		Bell-Cross (1974)
	C.G.N.F.	4.5 (C.n.)		
	W.F.	6.8 (C.n.)		
	T.F.	major[d] (C.g.)		
	E.G.N.F.	7 (C.g.)	2 (C.g.)	Duerre (1969)
	E.G.N.F.	8.5 (C.n.)	4 (C.n.)	
	C.G.N.F.	4.5 (C.g.)	4.3 (C.g.)	
	C.G.N.F.	2.2 (C.n.)	2.2 (C.n.)	
Kafue[a], Zambia	C.G.N.F.	major[d] (C.g.)		Carey (1971)
	C.G.N.F.	major[d] (C.n.)		
	C.G.N.F.	43 (C.spp.)		MacLaren (1956)
	S.N.F.	6 (C.spp.)		
Kafue River, Zambia	C.G.N.F. (1954-1958)	29-52 (C.spp.)		Williams (1960)
	S.N.F. (1954-1958)	7-24 (C.spp.)		
	S.N.F.	31 (C.spp.)	14 (C.spp.)	Everett (1971)
Lake Bangweulu, Zambia	C.G.N.F.	60 (C.g.)		Harding (1960)
	C.G.N.F.	7 (C.m.)		
Lake Mweru-Wantipa, Zambia	L.F.		90 (C.g.)	Allen (1964)
	C.G.N.F.		7 (C.g.)	
	C.G.N.F.		<1 (C.g.)	Allen (1965)
Lundazi Dam, Zambia	E.G.N.F. (1954)	42 (C.g.)	25 (C.g.)	Mortimer (1959)
	E.G.N.F. (1955)	50 (C.g.)	28 (C.g.)	
	E.G.N.F. (1956)	50 (C.g.)	27 (C.g.)	
	E.G.N.F. (1957)	44 (C.g.)	23 (C.g.)	
	E.G.N.F. (1958)	20 (C.g.)	8 (C.g.)	
Lungwewungu River, Zambia	C.G.N.F.	14 (C.g.)		Bell-Cross (1974)
North Kashiji River, Zambia	C.G.N.F.	30 (C.g.)		
South Kashiji River, Zambia	C.G.N.F.	4.6 (C.g.)		
Small stream, Zambia	A.		14 (C.spp.)	Mortimer (1965)
Stream, Zambia	W.F.	7 (C.spp.)		Bell-Cross (1971)
Zambezi River, Zambia	C.G.N.F.(Below dam wall)	6 (C.g.)		Bowmaker (1960)
Zambezi River, Rhodesia	C.G.N.F.(before closure of Kariba Dam wall)	3-5 (C.g.)		Harding (1964)
Lake Kariba, Rhodesia	C.G.N.F.	20 (C.g.)		Hard' ng (1964)
	C.G.N.F.	18 (C.g.)		Coke (1968)
	C.G.N.F.	16 (C.g.)		Van der Lingen (1971)
	E.G.N.F.		5 (C.g.)	Allen (1965)
	C.G.N.F. (1971)	4.5 (C.g.)		Anon. (1972)

Table 2. (cont'd)

Location	Fishing method	Percent catch by weight		by numbers		Author
	C.G.N.F. (1972)	3.3	(C.g.)			Anon. (1973)
	C.G.N.F. (1973)	3.8	(C.g.)			Anon. (1974)
	P.	2	(C.g.)			Balon (1973)
	E.G.N.F.	6.3	(C.g.)	5	(C.g.)	Bowmaker (1973)
(Mwenda Estuary, Lake Kariba)	P.	28	(C.g.)			Unpublished[b]
Mwenda River, Rhodesia	C.G.N.F.	20-76	(C.g.)			Van der Lingen (1962)
Lake McIlwaine, Rhodesia	E.G.N.F.			10	(C.g.)	Minshull (1975)
	E.G.N.F.	5.3	(C.g.)	1.6	(C.g.)	
Lake Kyle, Rhodesia	S.N.F.	1.9	(C.g.)	0.08	(C.g.)	
	A.	2.5	(C.g.)	0.9	(C.g.)	Ratcliff (1972)
	C.G.N.F. & P.S.F. (1965)	25	(C.spp.)			
Lake Chilwa[c], Malawi	C.G.N.F. & P.S.F. (1966)	40	(C.spp.)			
	C.G.N.F. (before closure of Cabora Basa Dam)	17	(C.g.)	3.5	(C.g.)	Morais (1974)

Footnotes

[a]Flood plain; [b]Unpublished data from Tiger Bay Fisheries (Pvt) Ltd., Norton, Rhodesia (13-yr range) average over 50%; [c]Lake on a drying phase; [d]No figure given.

Species Code

C.g. - *Clarias gariepinus*
C.n. - *Clarias ngamensis*
C.m. - *Clarias mellandi*
C.spp. - *Clarias* spp.

Fishing Method Codes

C.G.N.F. - Commercial gillnet fishery
E.G.N.F. - Experimental gillnet fishery
(B) - Bottom set
L.F. - Longline fishery
T.F. - Trap fishery
S.F. - Spear fishery
W.F. - Weir fishery
S.N.F. - Seine net fishery
P.S.F. - Purse seine fishery
P. - Poisoning
A. - Angling

Table 3. Importance of *Clarias gariepinus* in small dams of Rhodesia. (Data taken from publication and internal reports of Department of National Parks and Wildlife Management, Rhodesia.)

Location	Fishing method/date	% catch by Weight	% catch by Numbers	Author
Cleveland Dam, Salisbury	Experimental gillnet fishing /May 1962	24.1	7.8	Toots (1968)
	/November 1962	6.2	1.7	
	/September 1967	27.7	14.5	
	/January 1968	24.0	7.9	
	Angling/1962-1966	8.4	4.0	
Silalabuhwa Dam, Insiza, TTL	Experimental gillnet fishing /June 1970	3.3	2.0	Toots (1970)
	/March 1971	3.5	0.9	Toots (1971)
Nyambuia Dam, Marandellas	Experimental gillnet & seine net fishing/April 1969	90.0	66.0	Toots (1969a)
Nyakambira Dam, Marandellas	Experimental gillnet & seine net fishing/April 1969	0.0	0.0	Toots (1969a)
Gwenoro Dam, Gwelo	Experimental gillnet fishing /May 1974	58.0	10.0	Toots (1974a)
Ngamo Dam, Gwelo	Experimental gillnet fishing /April 1974	29.1	15.3	Toots (1974c)
Impali Dam, Selukwe	Experimental gillnet fishing /May 1974	61.0	40.0	Toots (1974a)
Whitewaters Dam, Lalapanzi	Experimental gillnet fishing /April 1974	27.4	10.2	Toots (1974c)
Ngondoma Dam, Ngondoma	Experimental gillnet fishing /April 1969	78.0	34.0	Toots (1969b)
	/April 1971	7.9	1.4	
	/April 1974	19.1	6.3	
Pindi Park Dam, Banket	Experimental gillnet fishing /September 1974	35.0	9.0	Toots (1974b)
Chimwemwe Dam, Banket	Experimental gillnet fishing /1972	23.3	6.1	Toots (1972)
Ngesi Dam, Ngesi National Park	Experimental gillnet fishing /March 1962	9.0	3.0	Van der Lingen (1962b)
	/June 1962	8.0	4.5	
	/October 1962	14.5	5.8	
Sebakwe Dam, Sebakwe National Park	Experimental gillnet fishing /June 1962	6.0	4.8	Van der Lingen (1962c)

7.8 kg 100 m^{-1} 24 h^{-1} over a 13-yr period (unpublished data from Tiger Bay Fisheries (Pvt) Ltd., Norton, Rhodesia).

The selectivity of some fishing methods should be utilized as a management technique to permit "catfish fisheries" on the major lakes of southern and central Africa. This would allow a heavy fishing pressure to be placed on this fish while not placing as much pressure on the less plentiful species. Many new dams will be built, hopefully with fish ladders incorporated into them, and it would be valuable to place commercial fisheries on these fish ladders during the floods to remove a percentage of all migrating fish. In the special case of *Clarias* spp. which are the only fish to move up the ladder at night (Bell-Cross 1960), special automatic traps harvesting all fish could be used during the hours of darkness.

From routine observations of commercial fisheries in Zambia, Bell-Cross (1974) found the numbers of large catfish to be increasing. This has been attributed to the reduction in numbers of the crocodiles once so plentiful on large African rivers. Although these larger catfish are of interest to individual sports fishermen, they are unlikely to increase the production of these fish which already produce an estimated one-fifth (20.1%) of the fish protein of East and Central Africa.

ANATOMY

Early writers (Angelopulo 1947) felt that the maximum size of the African catfish was dependent upon the locality in which it was caught. Although the size of the habitat can affect the maximum length, a more important factor appears to be the temperature regime of the environment. Personal observations by the author and data from several sources recorded by Bruton (1976) indicate that in Rhodesia and South Africa there is a definite trend to larger fish in more southern regions. The largest fish are found in the Orange and Vaal Rivers and the smallest "maximum" sized fish are found in Lake Kariba and the Zambezi River.

The head of *Clarias* spp. is a massive, broad, dorso-ventrally flattened structure, heavily armoured by very thick skull bones. It is covered by a very thin layer of skin through which can be seen the many small glanular tubercles (Fig. 3). The trunk stretches from the back edge of the skull to the uro-genital pores and is cylindrical. The caudal region makes up the rest of the body from the uro-genital pores to the tail. This section is elongate and laterally flattened. The variation in body shape is used by this adaptable fish. The dorso-ventral flattening of the anterior region is useful in terrestrial locomotion and benthic



Fig. 3. Skull and brain of Clariidae. A. Skull bones, dorsal view. B. Brain cavity with major nerves, dorsal view. C. Pituitary gland and brain, lateral view. abc. - air bladder capsule; at. - accoustic tubercle; cbl. - cerebellum; crb. - cerebrum; e. - eye; f - frontal bone; ff. - frontal fontanelle; inf. - infundibulum; mb. - maxillary barbel; n. - nasal bone; ob. - olfactory bulb; os. - olfactory sac; ot. - olfactory tract; p. - pituitary gland; pf. - parietal fontanelle; so. - supra occipital bone; II - optic nerve. (Adapted from Angelopulo 1947; Lehri 1966).

feeding, while the laterally flattened tail is well adapted for swimming. Another asset of the dorso-ventral flattening of the head region is the ability to evade seine nets as mentioned previously.

The colour of *Clarias* spp. varies from a mottled beige to dark red-brown or olive green depending upon the environment. When moved from one environment to another different one, the fish's colour can change completely in only 10-15 min. Mills (1966) feels that the fish is mottled in bright light due to pigment concentrations in special cells, called chromatophores. Counter shading common to most fish is present with a dark upper surface and a white ventral surface. *C. gariepinus* of East Africa acquires a red tinge bordering its dorsal fin during the breeding season (Bailey 1966). Similar colouring was noted in spawning *C. lazera* in Israel (Clay, in press).

The entire Clariidae family is scaleless with a mucus coating over a tough skin. All of these catfishes have sensory barbels located around their large terminal mouths. There are five pairs of barbels, the largest are a continuation of the corners of the mouth and are called maxillary barbels. The inner and outer mandibular barbels are located on the lower jaw, while the nasal barbels are located on the upper jaw. All barbels are proportionately longer in the very young fish. The young fish have been observed in Lake Kariba to hang suspended by the surface tension of their long barbels; in this upright position they filter water and zooplankton through their gill rakers in order to feed (Junor, pers. comm.). The barbels have been reported to be richly supplied with taste buds (Das 1927) and are innervated by the Vth nerve (Angelopulo 1947). More recently Rajbanshi (1966) has found through histological studies that all of the barbels on *C. batrachus* are similar in makeup. Taste buds are present but no tactile organs are to be found in the epidermal region. An hypothesis for this is drawn up on the basis that *Clarias* spp. are bottom feeders while other catfishes such as *Wallago attu* which have large numbers of tactile organs are mid-water feeders.

There are two small openings located on the snout. These nasal pores are external sensing devices - the posterior opening is a slit located behind the nasal barbel while the anterior one is slightly raised. The lateral line, another external sensory receptor, extends the entire length of the trunk and caudal regions and is connected to the exterior by a series of raised pores (Mills 1966). Although not visible on the skull of a living fish, the courses of the branches of the lateral line are indicated by grooves in the bones of the skull (Angelopulo 1947).

The eyes are located on each side of the massive skull near the anterior end. They are very susceptible to damage by gillnets even though they are protected by a thick membrane (Mills 1966). Posterior to the eyes and centrally located in the skull is the frontal fontanelle and further back is the smaller parietal fontanelle. The latter is often pierced by fishermen to kill these fish.

The teeth are located in bands in the mouth and throat. The premaxillary band is in a crescent shape and was frequently used for taxonomic identification. Jubb (1967) has recently synonymized *C. gariepinus* and *C. mossambicus* as the only reason for their previous separation into two species was on the basis of their teeth patterns. Plates 1 and 2 show the upper jaws of two *C. gariepinus* from Hartbeespoort Dam in the Republic of South Africa that would have previously been separated into two species. The vomerine pads located in the throat cavity appear to serve a swallowing function. The author has observed that when these pads are stimulated with a blunt object the throat convulses in a swallowing reflex.

In *Clarias* spp. the unpaired dorsal and ventral fins are well developed and do not join the caudal fin which forms a separate rounded homocercal tail. All the fins are supported by soft dermal fin rays (lepidotrichia). As well as these soft rays each of the pectoral fins is supported by a hard calcareous spine. This serrated spine is used for both terrestrial locomotion and defence (Nawar 1954). They can inflict serious wounds which are often infected by the ever present mucus secretions of the catfishes; this has given rise to the belief that these spines are poisonous. Das (1927) reported *C. batrachus* using these spines for fighting. When overcrowded in small containers, the fish will fight until the weakest die. The author has observed that they often appear to lie side by side and by making sudden whip-like motions they endeavour to drive their spines into each other. These spines make a formidable weapon by their locking mechanism - a slight forward clockwise twist locks the spine (Nawar 1954).

The spinal column of *C. gariepinus* is made up of two sets of vertebrae totalling between 60 and 73 (Angelopulo 1947) while *C. lazera* has between 65 and 67 vertebrae (Nawar 1954). This is another indication of the close relationship of these separate species. The first five of these vertebrae are called the "complex vertebrae" (Mills 1966) and have been modified to form the Weberian ossicles which are fused to the skull and enclose a lobed air sac. This may be a special adaptation to provide better muscular attachment and flotation for the relatively large head. The first five (or less) unattached vertebrae have single thick spines. It is from these vertebrae (usually the first three) that the bones for ageing the catfish are chosen (Appelgate 1950; Balon 1974; El Bolock 1972). After these come a short series of vertebrae with double spines; the majority of vertebrae follow with single slender spines. The caudal and trunk vertebrae are similar except for the haemal canal containing the caudal vein and artery. The terminal vertebrae are connected to

the haemal spines as a series of hypurals forming the base of the homocercal tail (Angelopulo 1947). There are 13 pairs of pleural ribs. These "floating ribs" are attached to the anterior surface of the ventro-lateral process of the vertebrae. From a consumer's point of view, Clariidae with their relatively small number of bones are desirable fish when compared to the Cyprinidae family which have large numbers of very fine bones throughout the body.

The sex of *Clarias* spp. can be observed externally on most fish over 17 cm in length and internally on most fish over 13 cm. Males and females can be separated on the basis of the external genitalia. The males have a sharply pointed genital papilla into which the ureter and genital ducts empty. In the female the genital opening is little more than a mole-like growth. In both sexes the uro-genital pores are located behind the anus.

The ovaries in *Clarias* spp. are paired, elongated, sac-like structures lying in the body cavity. They are usually equal in size but occasionally one is larger than the other and rarely one is absent. The caudal end of the ovary is attached to the genital papilla through a common oviduct. As maturity progresses the ovary changes colour from creamy white to greenish or reddish-brown. The testes are also paired, elongated structures but they are more flattened and ribbon-like. When immature, the testes are smooth edged but at maturity the edges become serrated. The size of the testes is generally equal but, as with the ovaries, they can vary. The colour at maturity is usually creamy white.

The pituitary gland (Fig. 3) is a white ovoid structure located immediately posterior to the prosencephalon on the floor of the brain cavity (Angelopulo 1947). It is easily removed and dehydrated in acetone and can be used for induced breeding (Van der Waal 1974).

There are two peculiarities in the African catfish: both the liver and kidney have lateral lobes embedded in the body musculature (Nawar 1955). The liver lobes are embedded behind the pectoral girdle, while the kidney lobes are located behind those of the liver (Plates 3 and 4).

The major nerves of *Clarias* spp. are large and easily traced. This factor along with the wide distribution and extreme hardiness make *Clarias* spp. an ideal fish for classroom study (Bowmaker, pers. comm.; Mills 1966). Supplying African secondary schools with indigenous fish for dissection would be a large industry in itself. This would certainly bear investigation rather than importing exotic fish into Africa and Asia where foreign exchange is always at a premium.

RESPIRATION

These catfish are one of the most widely distributed fish in Africa, often being most plentiful in environments inhospitable to other fishes. One of the greatest assets for *Clarias* spp. in regard to its adaptability is its modified respiratory system. The tropical catfish's ability to utilize atmospheric oxygen allows it to live in shallow tropical pools in which few other freshwater fish can survive due to the low dissolved oxygen levels resulting from decomposing organic matter and the high temperature regime. Most fish that are adapted to aerial respiration live in such environments (Saxena 1963). *Clarias* spp. falls into Hora's (1935) second category for air breathing fishes, i.e. those fishes from pools, marshes, and side streams.

The accessory breathing organs of *C. batrachus* are made up of the following four parts: (1) supra-branchial chamber, (2) the fans or gill plates, (3) the dendritic organ, and (4) the respiratory membrane (Munshi 1961). Recently it has been proven through electron microscopy that the respiratory organs of *C. batrachus* are modified gill structures (Hughes and Munshi 1973). The respiratory organs of the Clariidae are located in two chambers in the posterior part of the skull under the thick dermal parietal bones. Each chamber or arboescent organ consists of two dendritic tree-like growths that arise from the upper ends of the 2nd and 4th gill arches.

The respiratory chambers are walled by "gill-fans", structures that are histologically similar to gill filaments (Greenwood 1961) and which act as valves which control the retention or expulsion of air (Bowmaker, pers. comm.).

The two respiratory trees develop from the epibranchials of the gill arches. The posterior tree is always larger and more fully developed than the anterior. These arborescent organs develop late in the postlarval stage (Greenwood 1956). The first knob or tree develops about 3 cm while the anterior tree (from the second gill arch) develops somewhat later, possibly about the 5-cm stage (Greenwood 1961). This is very suitable as *Clarias* spp. take about 4-6 wk to grow to 5 cm. As the nursery habitat is still well oxygenated at this time, the fry are able to temporarily breathe through their gills.

A *Clarias* spp. releases several large bubbles through its opercular openings when rising to the surface to breathe. It then opens its mouth above the surface, closes it, and dives steeply. The angle of the dive forces the air to the back of the buccal cavity where it rises and fills the two respiratory chambers.

Donnelly (1973) suggested that the barbels are used as trigger mechanisms for stimulating the release of air; this is done by sensing the surface tension of the water as the fish rises.

Early workers thought that *Clarias* spp. would drown if kept from the surface (Boake 1866; Day 1868; Dobson 1874). It has now been proven that *C. batrachus* does not drown if prevented from reaching the surface as long as it is kept in well oxygenated water with the carbon dioxide continuously removed (Abdel Magid 1970; Hora 1935; Jordon 1976). Oxygen uptake by haemoglobin of the catfish is very good and it is remarkably insensitive to high carbon dioxide levels (Fish 1956). This latter point is very important in aerial respiration as the carbon dioxide levels would tend to build up in the intervals between aerial breathing. At oxygen levels below 1 ppm in the water all aquatic respiration stops completely as the oxygen would tend to diffuse from the blood through the gills to the outside environment (Saxena 1963).

The rate of opercular movements increases until just before surfacing, and is lowest immediately after (Abdel Magid 1970). Low oxygen tensions increase the aquatic respiration rate; at very low oxygen levels (>1 ppm) opercular movements cease. *C. batrachus* at temperatures around 30°C come to the surface as frequently as half-minute intervals; this is due to the very low dissolved oxygen (D.O.) levels at high temperatures (Das 1927).

This ability to change from aquatic respiration to aerial respiration makes the family Clariidae ideal for culture. In tropical warm water pond culture one of the greatest assets, high plankton production, is also one of the greatest drawbacks. The high phytoplankton production leads to excessive diurnal fluctuations in D.O. levels, often resulting in anoxic conditions by dawn. *Clarias* spp. are able to withstand this condition with no adverse effects. Even more difficult to use for fish culture are oxidation ponds; the high biological oxygen demand (B.O.D.) in these ponds often means that even though there is a very high level of productivity, it is almost impossible for ordinary fish to survive. Further advantage of aerial respiration for the culturist is in the marketing of live fish (Hora 1934). Because *Clarias* spp. can survive for as long as 18 h in a basket with wet grass, the marketing problems, especially in the tropics, are reduced considerably.

Several theories have been put forward for the evolution of the air-breathing fishes. Most of these are only further adaptations of an early theory put forward by Carter and Beadle (1931) and later modified by Hora (1935). This theory states that lack of oxygen in shallow tropical waters must have been responsible for evolution of the air breathing habit in these fishes. Carter (1931) also pointed out that the high carbon dioxide levels were a very important factor in this evolution. According to Hora (1935) the dendrite tree of the Clariidae was originally developed to increase the surface of the respiratory epithelium.

Although several adaptations have been added to the respiratory system of the Clariidae, several items have also degenerated. The air bladder of *Clarias* spp. is present in small fish and becomes relatively smaller as the fish grows. It is eventually enclosed in a boney capsule projecting laterally from the back of the skull, just posterior to the junction of the fused and free vertebrae (Das 1927). The genus *Xenoclarias* is one group of the Clariidae that has readapted to lacustrine conditions. The aerial respiratory organs have degenerated in these fish until they no longer surface and, in Lake Victoria, live permanently at depths of over 40 m (Gee 1969).

HABITAT AND DISTRIBUTION

Members of the Clariidae are found distributed throughout Africa from the Umtamvuna River (Crass 1966) and the Orange River in the south, to the Nile in the north; through the Near East, especially Syria, and into India and the Far East (Hora 1937). They are now also found in Hawaii and in southern Florida in the continental U.S.A. Figure 1 shows the world distribution of Clariidae.

The major habitat for *Clarias* spp. is warm shallow tropical pools, although several species have readapted to deep lacustrine conditions (viz. *Xenoclarias* spp.). In Lake Kariba the catfish was found to be more common at 2 m depth than at 5 m, and more common at 5 m than at 10 m (Mitchell, pers. comm.). The catfish is one of the most widely distributed fish in Africa. The major limiting factors to distribution are low temperatures (Pardue 1970) and salinity (Clay, in press). They have been found above and below waterfalls (Bell-Cross 1968), in pans or pools that were previously dry and have no inlet (Das 1927), and in drying river pools with only mud as a medium of support (Donnelly 1973).

Although a minimum temperature of less than 10°C is generally fatal to *C. batrachus* (Pardue 1970), a diurnal range of between 13.5 and 27.5°C (Donnelly 1973) does not seem to have any adverse effects. This tolerance of a wide diurnal range of temperatures is another strong asset for fish culture operations where the shallow ponds often can exhibit great daily fluctuations in temperature.

Salinity trials on *C. lazera* indicate that the fish cannot survive salinities over 15 ppt and, as such, estuaries of more than 50% seawater would provide an effective barrier to distribution (Clay, in press).

Their aerial respiration and ability to move over land are major adaptations allowing such wide distribution. In India, when pools are disappearing in the dry season, *C. batrachus* bury themselves in the moist mud below the surface (Das 1927). This gave "rise to the superstition [then] current amongst the peasantry that they have been rained from the clouds...as they appear in great numbers after the monsoons" begin (Das 1927). Whether these fish truly aestivate must still be proven; Donnelly (1973) observed *C. gariepinus* in drying pools in Africa and found that, even when only a thick liquid mud remains, the fish were still actively breathing. In this case predators reduced the numbers of *Clarias* spp. as the pool dried up; the major predators besides man in Africa are crocodiles (Hipple 1946), the fish eagle (Donnelly 1973; Jubb 1952; Junor 1967) and marabou storks (Donnelly 1973).

An unusual adaptation to habitat desiccation in the African catfish was observed in Natal, R.S.A. Clements (1974) observed several *Clarias* spp. in a small river pool to cross a short sand bar and jump into a "den". The den had an opening about 15 cm in diameter and contained water below ground level. This retreat to a den utilizing ground water is one method of survival when the surface water dries up. Unlike *C. batrachus* described earlier, *C. gariepinus* and its near relative the vundu *(Heterobranchus longifilis)* are only partially adapted to habitat desiccation. When conditions become too severe, these fish often try to leave the local environment, usually by means of terrestrial locomotion.

Much work has been done on movement of Clariidae over land (Boulenger 1907; Das 1927; Johnels 1957; Welman 1948). The spines are kept rigid during "walking" by a T-joint that locks into position (Nawar 1954, 1955). Johnels (1957) filmed this locomotion as a rolling movement using the spines alternately as levers to heave the body from the ground (Fig. 4 shows a sample of this motion). Sandon (1950) records *C. lazera* as moving over 200 m from a swamp to a river in the Sudan. Although terrestrial locomotion has been observed and filmed, the author has observed that it cannot successfully take place over dry ground and Bell-Cross (pers. comm.) noted that it generally involves a downhill slithering type of walk.

REPRODUCTION

The reproductive biology of a fish being considered for culture is perhaps the second most important factor after the study of feeding and growth. Many researchers have studied the reproduction of the Asian catfish (Khan 1972; Sidthimunka et al. 1966; Sidthimunka and Ekuru 1959; Wanichakorn 1962) and the African catfish (Carey and Bell-Cross 1967; Corbet 1960; Holl 1966; Nawar and Yoakim 1962; Van der Lingen 1965; Van der Waal 1974). The Asian catfish *(C. batrachus)* appears to be similar in breeding habits to the American catfish *(Ictalurus punctatus)* by breeding in dens, while the African catfish *(C. gariepinus)* more closely resembles the cultured carp *(Cyprinus carpio)* and breeds on flooded vegetation. Neither method poses any serious problems to culturists that cannot be solved by some basic research.

The African catfish was first investigated in East Africa by Greenwood (1955). He followed a spawning run in a flooded low lying area adjacent to a stream. During the time of this spawning run, he artificially fertilized some ova and found that they hatched within 23-25 h at 24°C. In later studies Greenwood (1957) found the ova hatched in nature between 36 and 42 h after fertilization. Holl (1968) found *C. gariepinus* in Central Africa had similar breeding habits and the same size ova (1.5-2.0 mm diameter) as the previously described *C. gariepinus*, and that it required 40-48 h at a temperature of 20-24°C for hatching to occur.

Fig. 4. The muscular contraction of the Clariidae in swimming (A) and terrestrial (B) locomotion (white shading indicates muscular contraction).

The larvae ranged in size from 3.5-4.0 mm and both Holl (1968) and Greenwood (1955) report the larvae swim actively and have a ventral sucker for resting on the substrate. Pham (1975) found the larvae of *C. lazera* when cultured must be aerated for at least the first 12 days. Bowmaker (1973), in his description of the development of a Clariid (thought to be *Heterobranchus longifilis*), records they were almost continuously quiescent, resting on the bottom. These larvae were larger than *Clarias* spp. (5.3 mm at hatching + 8 h), although the fry were approximately the same size in later development. All the above workers indicate the extreme rapidity of development in the first 50 h. This may be an adaptive feature to compensate for the short-lived nature of the flooded nursery grounds. Greenwood (1955) found the young fry could withstand diurnal temperature variations of 22-34°C. The yolk sac is completely resorbed 6-8 days after hatching when the fry begin feeding on zooplankton and insect larvae. Pham (1975) found the larvae reached 15-30 mm in 15 days in culture conditions when feeding on zooplankton.

The breeding behaviour of *C. gariepinus* was studied in aquaria by Van de Waal (1974). The spawning was found to occur at intervals, each emission of ova being made up of a small cloud of eggs followed by some gas bubbles. Van der Waal (1974) observed that "while swimming actively together, the male ... folds himself around [the female's] head and body for a period of from 15-30 sec. During the last few seconds, the male's body presses hard laterally on the female's head, shivering continuously. At this stage the female's body also starts to contract ... [and] in a sudden movement the female bends her body sideways, often pushing her face into the substrate while the male moves backwards ... still folded around the female's body and still shivering. Immediately a small cloud of ova is ejected ... ." The eggs were widely distributed over the substrate by the vigorous flicking of the female's tail. A large adhesive disc attaches the eggs to the substrate within about 60 sec. In a natural situation this adhesive nature permits wide distribution and thus adequate oxygenation of the ova. Similar behaviour was observed in the field with *C. lazera* in Israel (Clay, in press).

Unfortunately the adhesive nature of the eggs makes artificial fertilization of large numbers difficult as they tend to clump together and high mortalities result. Work needs to be carried out on artificial stripping, fertilization, and hatching of the African catfish to facilitate the production of the millions of fry required for culture. Clay (in press) has tried using the method described for carp ova (also sticky by nature) and found it successful in preventing clumping of ova.

Under natural conditions a serious problem of fry mortality among the African catfishes has been found. It is generally attributed to cannabalism, both parents and larger fry preying upon fry. This cannabalism has been a factor leading to Micha's (1975) failure to achieve survival of fry in densities over 1450 individuals per ha. After hatching, the fry of *C. batrachus* have also been found to be cannabalistic, so intensive feeding and frequent grading are required to prevent mass mortalities in culture conditions.

The ova of *C. macrocephalus* are slightly smaller in size (1.3-1.6 mm) (Sidthimunka 1972) than the ova of the African catfish although Khan (1972) found *C. batrachus* ova in India averaged 1.8 mm in diameter, with the fry hatching out at 5.8 mm. *C. batrachus* of Thailand spawn in deep water, preparing a horizontal hole or den in the bank about 20 cm in diameter and 25 cm deep. The eggs are deposited and stick to the vegetation and soil surface, and hatch in less than 20 hours at 25-37°C. *C. macrocephalus* spawn in water less than 30 cm deep (Tongsanga et al. 1963) and build their 'redd' or nest about 30 cm in diameter and 5-8 cm deep on the grassy bottom. After the eggs are deposited and stick to the substrate the male then guards the nest while the female remains close by.

Most authors indicate that, although the breeding season for both the African and Asian catfish may vary as to the month of occurrence, it nearly always coincides with the onset of the rainy season (Bowmaker 1973; Greenwood 1957; Holl 1968; Sidthimunka 1972). The author observed that *C. lazera* in Israel spawn at the end of the rainy winter season as the weather warms but before the winter flood waters recede. This latter observation may indicate that temperature is the major stimulus dependent upon sufficient water. All researchers have emphasized the short duration of their observed respective spawning runs. Bowmaker (1973) describes *C. gariepinus* from Lake Kariba as going up the rapids in "gay abandon" with no concern of damage.

Bowmaker (1973) had the unique opportunity of observing (through the clear water beneath a Salvinia mat in the Mwenda estuary of Lake Kariba) that *C. gariepinus* gathered before the floods and lay quietly on the bottom in 1-2 m of water. The suddenness and short duration of the run reported in Lake Victoria (Greenwood 1956) indicate this may also be the case there.

Seasonal maturity in summer rainfall areas follows a definite pattern as shown in Fig. 5. Until immediately before the rains few fish are truly "ripe" - yet within 12 h of a heavy rain most fish are "running ripe" (Bowmaker 1973). Both Greenwood (1957) and Holl (1968) have remarked on the "one shot" nature of the spawning run; Bowmaker (1973) found the ova diameter of *C. gariepinus* had a unimodal distribution, this being indicative of a single spawner. More recently it has been found that there is a second peak in the ova distribution at about 300 microns - too small to find by conventional



Fig. 5. Estimated percent of adult population being mature at different times of the year in tropical regions of seasonal summer rainfall.

washing and hardening techniques (Clay, in press). This latter find indicates that the 'one shot' nature of the spawning may be more attributable to environmental conditions than the reproductive biology of the tropical catfish. Idyll (1969) has remarked that *C. batrachus* spawns through much of the year in Florida. This may be because seasonal rains do not have quite the same regularity there that is found in parts of Asia and Africa.

The size at which a catfish reaches maturity seems to vary with location. Pham (1975) found cultured fish ready to breed at 8 mo of age. Table 4 gives some examples of lengths at first maturity and fecundity of ripe females. Bowmaker (1973) has called *Clarias* spp. an 'average fecund fish' and Greenwood (1955) has mentioned the high mortality in shallow flood plains for catfish fry. These two factors make it hard to understand how *Clarias* spp. can often make up over 50% by weight of the commercial catch throughout East and Central Africa. One possible explanation for this, aside from the ruggedness and large size of the Clariidae, is the possibility of multiple spawning. Pham (1975) found that artificially induced spawning could be repeated in 3 or 4 mo.

Greenwood (1957) found small fry in the lake long after the single spawning run and postulated that this meant either spawning was not confined to the streams and rivers during the floods and occurs in the lake as well, or the growth rate was low in the lake, in the order of 5 mm per month. Although the latter interpretation was favoured, this is difficult to accept in view of recent work on growth rates. Corbet (1960) has mentioned that young *C. gariepinus* (3-14 cm) were found along rocky wave-washed shores of Lake Victoria. This is in an environment with similar vigorous water movements to the temporary streams where spawning (normally) takes place. In Lake Kariba, Bowmaker (1973) found ripe fish away from the rivers at the time of breeding. From Matthes' (1960) statement that *C. gariepinus* of Lake Tanzania breed in the bordering swamps and marshes, it is not a far (evolutionary) step for the catfish to begin breeding on vegetated flooded shores of man-made reservoirs. It can thus be assumed that African catfish may in some lakes be made up of two separate breeding populations, those invading flooded streams and pans and those breeding on newly flooded shore lines or even rocky wave-washed coasts, another indication of the adaptability of this fish.

For culture purposes, leaving *Clarias* spp. to breed on their own in special breeding ponds is an expensive and often risky business, as breeding may not occur. Present-day moves for nearly all fish culture operations are towards induced breeding by chemical injection. This usually involves the injection of a pituitary extract or some substitute into an already ripe fish (see Table 5). Khan (1972) has found mashing testes and stripping ripe eggs into the sperm suspension was very successful. Problems did occur though, in that the eggs were non-floating and sticky by nature and clumping resulted. Khosa and Chandrasekhar (1972) found that a weekly intramuscular injection of 12.5 micrograms of copper acetate in ½ ml of 0.75% saline for 3 mo caused formation of many enlarged yolky eggs when compared to control fish. More experimental work will determine optimum chemical levels to stimulate such maturation and optimum hormonal levels to allow artificial stripping. This should provide the information which will result in the production of a continuous supply of fry. This is more desirable than a sudden natural production during the normal breeding season.

In male fish with the gonads surgically removed, regeneration of the testes was found to occur and a large change in the histological structure of the pituitary was noted with little or no change in the thyroid (Lehri 1966). This leaves little doubt that the pituitary is of major importance in maturation and spawning of the tropical catfish. The location of the pituitary gland is shown in Fig. 3. It is located on the ventral side of the brain attached to the infundibulum and can be found by removing the upper skull plates (supraoccipital) and exposing the brain. If tweezers are used to break the notocord, the brain can be lifted up and forward from the posterior end and the small creamy coloured ovoid body of the pituitary gland will be found beneath.

Lehri (1967, 1968) divided both the male and female sexual cycles into five phases: a) resting phase (3 mo), b) early maturing phase (4 mo), c) advanced maturing phase (2 mo), d)

Table 4. Table showing the length at earliest maturity and fecundity of the Clariidae.

Species	Location	Sex	Length at maturity (cm)	Length/Fecundity	Author
C. gariepinus	Lake Victoria, Uganda	Female		50-75/ 32-48,000	Greenwood (1957)
C. gariepinus	Lake Tanzania	Female		/ ± 4,000	Matthes (1960)
C. gariepinus	Katanga (Zaire)	Female		60-70/ 30-40,000	Coureur & Spaas (1956)
C. gariepinus	Lake Kariba, Rhodesia	Male Female	53 44	/ 1,450/cm	Bowmaker (1973)
C. gariepinus	Mazoe Dam, Rhodesia	Male Female	43 38		Van der Lingen (1965)
C. gariepinus	Lake McIlwaine, Rhodesia	Male Female	45 39	45-60/ 20-70,000	Munro (1965)
C. gariepinus	Lake McIlwaine, Rhodesia	Male Female	25 28	30-70/ 4-72,000	Clay (in press)
C. gariepinus	Oliphants River, R.S.A.	Male Female	40 40	-	Van der Waal (1970)
C. lazera	Bangui, C.A.R.	Male Female	7-10 mo[a] 7-10 mo[a]		Micha (1975)
C. lazera	Bouake, Ivory Coast	Male Female	8 mo[a] 8 mo[a]		Pham (1975)
C. lazera	Lakes Edko & Mataria, Egypt	Female		35 ave/ 38,000	Nawar & Yoakim (1962)
C. senegalensis	Lake Nuguna, Ghana	Male Female	32 32		Thomas (1966)
C. batrachus	[Bangeladesh]	Female		32 ave/ 11,612	Mookerjee & Mazumdar (1950)
C. macrocephalus	Thailand	Male Female	22 22		Tongsanga et al. (1963)

[a]No size was given, fish were held in fish ponds.

Table 5. Dosage quantities of various chemicals and hormones to stimulate spawning in ripe Clariidae.

Species	Injection	Quantity/Dose	Reaction time (hours)	Author
C. batrachus	Pituitary extract	0.5 ml / 1 gland	8-10	Khan (1972)
C. macrocephalus	Pituitary extract	1 ml / 2.6 mg gland 1 ml / 3.9 mg gland	14-16 14-16	Tongsanga et al. (1963)
C. lazera	Desoxycorticosterone	5 mg Doca/ 100 g fish	12	Micha (pers. comm.)
C. lazera	Desoxycorticosterone	5 mg Doca		Pham (1975)
C. gariepinus	Pituitary extract	1 ml / 1 gland	15	Van der Waal (1974)

functional maturity phase (2½ mo), e) spent phase (½ mo). From this work it is possible to estimate the stage of sexual development of a population by back counting the months to the next rainy season or from the last spawning season.

The initial groundwork by fish culturists for controlled breeding has been completed. What is now needed is practical, applied work in the field of induced and controlled maturation. In effect, a systematic breeding program, with a series of local variety trials, is the only major requirement to solve many of the problems of fry porduction for catfish culture.

FEEDING AND DIGESTION

Clarias gariepinus has been described as an omnivorous scavenger by Jubb (1967). The term "omnivorous" in this case does not make it clear whether plant material makes up any significant proportion of the dietary items. The feeding habits of *Clarias* spp. have been studied in many differing habitats, giving varying results (see Table 6).

Angelopulo (1947) suggested that the short gut of *C. gariepinus* indicated a carnivorous natural history. Thomas (1966) felt that *C. senegalensis* must be carnivorous as there were no pyloric caeca in which cellulose digesting microorganisms could live. From a fish culture point of view, the most important query is not whether *Clarias* spp. can digest cellulose but whether they can use plant proteins. Poll (1964) found *Clarias* sp. do ingest plant material, and Cockson and Bourne (1972) listed plant material in the diets of Lake Chilwa *C. gariepinus*. Munro (1965) found that in certain locations *C. gariepinus* in Lake McIlwaine feed on large amounts of plant material. Many authors accept at the outset of their study that *Clarias* spp. are indiscriminant feeders and the presence of large quantities of non-animal material, which is of little food value to a carnivorous species, must be regarded as incidental (Groenwald 1964). It is on this basic point (the ability of *Clarias* spp. to utilize plant material) that the entire potential for future culture of the Clariidae rests.

In Thailand, due to high popular demand and large quantities of relatively cheap low quality marine trash fish, *Clarias* spp. culture has become an important industry in the last 20 yr. Even in this case, it will be necessary to utilize other fish feeds when technology finds alternative uses for this previously cheap animal protein feed. In Bangladesh it was found that over a 6-mo period *C. batrachus* lost weight when fed a total plant diet (Mookerjee and Mazumdar 1950). Preliminary experiments on plant diets have been carried out in Egyptian fish ponds (Iman et al. 1970). Unfortunately confusion over wet weight-dry weight conversions makes the data inconclusive, but this work does indicate *C. lazera* can survive and grow on a diet composed of plant materials and whatever natural food is available in the ponds.

Generally speaking, surveys show a change in diet with size of fish. A suitable diet must maintain the health of the fish, allow flesh production, and in a fish culture situation allow a profit with or without supplemental feeding. Fish culturists in developing countries are interested in fish about 25 cm (250 g) or less in size. For the initial growth of this size catfish, the most important natural foods are insects and zooplankton; a vegetable substitute needs to be found to allow supplemental feeding.

Yanni (1972) found that adding olive oil (an unsaturated fat) to the diet of *C. lazera* increased the fat and reduced the water balance in almost all tissues; when added to the diet in excess, the fat tends to be laid down as unsaturated fat in the fat bodies. This information is useful, as it indicates *Clarias* spp. can utilize small amounts of fat as a concentrated energy source in supplemental feeding for pond culture. This would result in less protein

Table 6. Feeding habits of the African catfish *(Clarias gariepinus)* from various locations.

Survey location	Season	Fish length (cm)	Food types (% volume) (%* frequency) plankton	fish	insects	molluscs	non-animal	Author
Barberspan, TVL., RSA			++	++	+		+	Groenewald (1964)
Barberspan, TVL., RSA		0-12	11%					Schoonbee (1969)
		40+	22%					
Vaal River, TVL., RSA			+	+	++		+	Groenewald (1964)
Oukskei River, TVL., RSA			++		+		++	
Incomati River, TVL., RSA	Summer	35	1%	69%	6%	8%	13%	Gaigher (1969)
	Winter	35		17%	24%	27%	9%	
Lake Sibayi, Natal, RSA			15%	30%	12%	5%	7%	Minshull (1969)
Wankie National Park, Rhodesia					+++			Weir (1972)
Savory Dam, Rhodesia	Summer	3	++					Holl (1968)
		3-8			++			
Ngondoma Dam, Rhodesia			30%*	12%*	25%*	4%*	12%*	Toots (1969a)
Nyambuia Dam, Rhodesia					35%*	4%*	55%*	Toots (1969b)
Cleveland Dam, Rhodesia				11%*	30%*		24%*	Toots (1968)
Lake McIlwaine, Rhodesia		1.2-7.6	+	+	++			Munro (1967)
		20-40	4%	3%	85%	8%	+	Munro (1965)
		40-60	21%	2%	55%	18%	+	
		60+	65%	1%	14%	9%	10%	
		5-30		++				Murray (1975)
		30+	++	+				
	Summer			35%*	35%*	15%*		Clay (in press)
	Winter		22%*	34%*	43%*	13%*	40%*	
Lake Kyle, Rhodesia	Summer	20+		15%*	29%*	26%*		
	Winter	20+	19%*	36%*	31%*	27%*	4%*	
Mazoe Dam, Rhodesia	Summer			10%*	50%*			
	Winter				39%*		13%*	
Savory Dam, Rhodesia	Summer	0-20			100%*			
		20-50	6%*	4%*	50%*		14%*	
		50+	10%*	10%*	40%*	12%*	14%*	
	Winter	0-20	50%*	16%*	32%*	8%*		
		20-50	51%*	17%*	35%*		13%*	
		50+	43%*	15%*	7%*	7%*	7%*	
Lake Kariba, Rhodesia/Zambia			++					Bowmaker (1973)
Kafue Flood Plan, Zambia				++	++			Carey (1971)

Table 6. (cont'd)

Survey location	Season	Fish length (cm)	Food types (% volume) (%* frequency) plankton	fish	insects	molluscs	non-animal	Author
Lake Chilwa, Malawi			+	+			+	Cockson & Bourne (1972)
Lake Malawi, Malawi				++				Fryer & Iles (1955)
Lake Ruka, Tanzania			+	+	++	+	+++	Mann (1964)
Lake Victoria, Uganda					90%			Corbet (1957)
Lake Victoria, Uganda		1-1.2	+		+			Greenwood (1957)
		24-90	2%	75%	23%	1%		Corbet (1961)
			+	++	+	+		Graham (1929)
Bugungu Stream, Uganda		0-15	9%	45%	44%			Corbet (1961)
Victoria Nile, Uganda		10-20	++		++			
Lake Nuguna, Ghana			5%	2%	75%		7%	Thomas (1966)

Note: + indicates the relative importance of feeding when the data is not compatible with the table.

being broken down for metabolic energy and more being available for tissue growth. This should lower the total protein input required for catfish production.

The rate of gastric digestion is loosely related to the time and size of the meal (Palmer 1973). The calorific value of a meal of fresh fish was found to change from 4.8 Kcal g^{-1} dry weight at ingestion to 4.1 Kcal g^{-1} at defaecation (Palmer 1973). This indicates that the fat of the eaten fish is being utilized in preference to the carbohydrates and/or proteins (these have a base calorific value of about 4.4 Kcal g^{-1} dry weight). As it can be assumed that fish protein will be highly digestible for *Clarias* spp., it follows that *C. gariepinus* must utilize fats relatively efficiently while not as readily digesting carbohydrates.

During starvation fish tend to utilize their carbohydrate stores (glycogen) first and afterwards draw on their fat (lipid) reserves (Love 1970). Kiermeir (1939) showed that, in starvation, fish which were sluggish tended to maintain very constant blood glucose levels, while active fish showed a gradual decrease. Yanni (1964) and Al-Gauhari (1958) found that little or no change in blood glucose levels occurred in *C. lazera* after 5-7 mo starvation. Thus it can be assumed that *Clarias* spp. fit into the category of sluggish fish (Love 1970), which might indicate a greater amount of their food energy will go into flesh production and less into activity.

The pH of the stomach of *C. gariepinus* is just slightly alkaline (pH 7-8) when empty, but becomes acidic (pH 3-4.5) when food is ingested. The intestine is generally slightly acidic (pH 5-6) (Palmer 1973). Cockson and Bourne (1972) found protease in the stomach and anterior intestine of *C. gariepinus* with an optimum enzyme activity level occurring at a pH of 4-6. The enzyme activity was found to be four times as great in the stomach as in the intestine. Most of the protease was found to be pepsin with a little trypsin; these enzymes are very general in nature and indicate a strong likelihood of digestion of all proteins with no specialization for the digestion of either plant or animal protein. Pepsin is found in both herbivorous and carnivorous mammals where it acts through the hydrolysis of proteins, especially among the amino acids, phenylalanine, tryptophan, tryosine, and leucine (White et al. 1968).

Cockson and Bourne (1972) also found amylase produced and located in the anterior and posterior intestine of *C. gariepinus*. Amylase has an optimum activity at a pH between 4.5 and 7 (White et al. 1968). As this shows that starch digestion cannot begin until the food is in the intestine, *Clarias* spp. would appear to be inefficient at starch (carbohydrate) digestion.

Palmer (1973) found that the percent inorganic material in the food increased from ingestion to defaecation with the greatest change occurring between the intestine and the rectum. This indicates that absorption is mainly of organic matter and occurs largely in the intestine as would be expected.

Clarias spp. has few predators. Crocodiles (Cott 1954; Hipple 1946) and predatory birds (Donnelly 1973; Junor 1967) are the only significant natural predators besides man. This may be the reason why cannibalism is often a problem in dense *Clarias* spp. populations. Cloudsley-Thompson (1959) put forward a theory that cannibalism is often a means of keeping population density down and conserving protein within a species.

Many surveys have indicated the importance of filter feeding habits of *Clarias* spp. (Table 6). Although juvenile catfish (<3 cm) were known to feed on zooplankton, it was often thought that adult fish (>30 cm) were more ontogenetically adapted to filter feeding than young juvenile fish (between 3 and 30 cm). This was because it was thought that the number and length of the gill rakers increased and the spacing decreased with length (Alexander 1966; Gilchrist and Thompson 1913; Jubb 1967; Thomas 1966). Murray (1975) found that the gill raker spacing in fact increases with length in *C. gariepinus*. Although the young *C. gariepinus* (3-30 cm) did not filter feed in Lake McIlwaine, under laboratory conditions they did successfully filter feed on plankton. This physical adaptation of *Clarias* spp. under 30 cm to filter feed is important when deciding on species selection for warm water pond culture, especially where temperatures warrant fertilization of the water.

Food selection in nature is variable; the Clariidae family tend to be opportunistic feeders. In Lake Victoria, Corbet (1956) found *C. gariepinus* so unspecialized in feeding that, when *Simulium* spp. which made up 90% of the diet were removed by insecticides (DDT), the fish changed to a diet of plants, molluscs, etc. Weir (1972) found *C. gariepinus* in pools in Wankie National Park, Rhodesia, feeding exclusively on the only food available, insects. Munro (1965) found that during the lunar emergence of chironomids *C. gariepinus* fed exclusively on these insects. Similar exclusive feeding patterns were noted by the author at Lake Kyle, Rhodesia, with termite flights after the heavy rains, and in Sand River Lake, Swaziland, during the seasonal bivalve (Lammellibranchia) "spawning run" into shallow water. Groenwald (1964) found *C. gariepinus* diet to be greatly dependent on food availability in different environments; thus the catfish becomes a piscivore when there is an abundance of fish available. In Zambia, Carey (1967) and Tait (1967) found *Clarias* sp. to be a major factor in

fish predation, especially in the Kafue River system, in the absence of the tiger fish (*Hydrocynus vittatus*). Generally, it can be stated that juvenile Clariidae have been found to be filter feeders at less than 3 cm in length (Corbet 1961; Munro 1967). Thereafter they tend to feed on almost anything, with a preference for animal protein.

Even when food may be in short supply *Clarias* spp. have one alternative adaptive feature. Social or pack hunting was observed and recorded by several authors (Anon. 1975; Donnelly 1966; Pooley 1972). During this activity, *C. gariepinus* of 75 cm and over form a long crescent-shaped line and drive small fish ahead of them by making a violent commotion within the water. They drive the prey fish along the shore until they reach a small bay or inlet which they can effectively seal off. They then commence to feed.

Thomas (1966) stated that the head and body adaptations of *C. senegalensis* indicated a sluggish bottom dweller; in Lake Nungua, Ghana, the catfish he observed were predominantly benthic feeders. Groenwald (1964) states that several features of the head region of *C. gariepinus*, such as size of mouth and bands of small (cardiform) teeth, are adaptations for manipulating large prey; the catfish he observed were largely piscivorous. Gilchrist and Thompson (1913) state that, for this latter species, the large number of fine, closely set gill rakers is clearly an anatomical feature of a plankton feeder. With such variable adaptations and catholic feeding habits, it is unlikely that food would ever provide a serious environmental barrier to the dispersal (Corbet 1961) or growth of these fish.

GROWTH AND PRODUCTION

The growth of the Clariidae has been the subject of much speculation but little detailed study. Due to their wide range of feeding habits, they have been repeatedly mentioned as having a great potential for culture (Brown 1955; Couveur and Spaas 1956; Symes 1954). This potential can only be realized if their growth rate is rapid and efficient under culture conditions. Little work has been done in this field and what has been done is very contradictory (see Table 7). The basic study (of production) that is so important for the utilization of this family of fishes for aquaculture has been sadly neglected.

The two very high growth rates of *C. gariepinus* found by Munro (1965) and Van der Waal (1973), shown in Table 7, appear to be rather overly optimistic. Munro (1965) utilized monthly frequency peaks in commercial gillnets to predict the growth of modal groups. With fish caught by gilling (the more regular way of a gillnet working) this is difficult (Holt 1963) and since *Clarias* spp. have a tendency to tangle their spines in gillnets of nearly any size, it is virtually impossible to accurately use this method. It is also possible that he "lost" the first year or two as the catfish would be too small to be adequately sampled by the commercial fisheries smallest nets (50 mm stretched mesh). Van der Waal (1974) used spines to age his fish. The initial growth rate he found for the first year is extremely high, especially when considering the growth rate in second and subsequent years. There is a possibility that, as was pointed out by Marzolf (1955), the lumen of the spine of the catfishes enlarges with age, thus reabsorbing the surrounding bone and with it the first ring. Corbet (1960) observed *C. gariepinus* in Lake Victoria to be between 6 and 12 mo old at 11 cm. This tallies well with the observations by Greenwood (1955) and Holl (1968) for wild fish (see Fig. 6).



Fig. 6. Growth of juvenile *Clarias gariepinus* in east and central Africa (from data of Corbet 1960; Greenwood 1957; Holl 1968).

Catfish under conditions of culture have a much more rapid growth rate. Sidthimunka (1972) found cultured *C. macrocephalus* in Thailand had reached a marketable size of 250 g (approximately 30.0 cm) in 4-5 mo, an average of about 55 g month^{-1}. Van der Waal (1970) achieved similar growth rates for *C. gariepinus* in the Republic of South Africa. By using pelleted feed, the fish reached 300 g in a 5-mo growing season (60 g month^{-1}). Mortimer (1964) reared *C. gariepinus* in fish ponds in Zambia and in a 6-mo growing season the fish grew from 675 to 1666 g, an increase of over 160 g month^{-1}. During the colder months, Mortimer grew fish from 50 to 300 g in 5 mo (50 g month^{-1}).

Table 7. Sample field growth rates of the Clariidae in the tropics.

Species	Ageing technique[a]	Location	Sex	Total length (cm) at end of time (Days) 0	7	15	60	120	270	(Years) I	II	III	IV	Author
C. batrachus	Obs.	(Bangeladesh)		.58	1.0									Mookerjee and Mazumdar (1950)
C. lazera	Vert.	Egypt	M							19.1	29.5	38.1	46.7	El Bolock (1972)
			F							19.3	29.8	37.2	44.8	
C. lazera	Obs.	C.A.R.				3[b]								Pham (1975)
C. gariepinus	Obs.	Uganda		.4	.8									Greenwood (1955)
C. gariepinus	Vert.	Zambia								10.0[c]	15.0[c]	20.7[c]	29.0[c]	Pivnicka (1974)
C. gariepinus	Obs.	Rhodesia							13					Bowmaker (1973)
C. gariepinus	CCC	Rhodesia					4.3	6.2						Holl (1968)
			M							30.0	47.0	60.0	72.0	Munro (1965)
			F							27.0	37.0	50.0	58.0	
C. gariepinus	Spine	R.S.A.								40.0	44.0	50.0	60.0	Van der Waal (1973)

[a]Abreviations are: Obs. - Observation
Vert. - Vertebrae ageing
Spine - Spine sections
CCC - Commercial gillnet catch curves

[b]Average of range of 2.5-3.5 cm given by author

[c]Length corrected to total length as original paper had them expressed as standard lengths

Feeding *Clarias* spp. in culture is another aspect requiring a great deal of future work. Little has been done on this subject even though *Clarias* spp. are widely cultured in some countries. Micha (pers. comm.) reported obtaining yields of 16-18 tons ha^{-1} in the Central African Republic (CAR) feeding *C. lazera* on cottonseed cake and brewers' waste. Van der Waal (1970) in the Republic of South Africa (RSA) found production in ponds with no feeding to be less than 70 kg ha^{-1}, with a mortality rate of 93%. In fish ponds with supplemental feeding of pelleted food, production was found to be 3.2 tons ha^{-1}, with a 90% mortality. The conversion rate of these latter fish was found to be between 5:1 and 6:1 (dry pellets:live weight fish). This does not compare very favourably with the 6:1 conversion ratio (wet weight food:live weight fish) by Sidthimunka (1972) in Thailand with catfish fed on marine trash fish.

Khan (1972) noted that cannibalism occurs among the fry if there is much size variation. The problem of cannibalism, a probable reason for Van der Waal's (1970) very high mortalities, is very common with *Clarias* spp. of southern Africa. It is another topic that requires much work to determine its true significance to culture.

The culture of *Clarias* spp. both alone and in polyculture with *Tilapia* spp. (Symes 1954) and other species requires "variety trials" in many locations before it can be offered to farmers and/or companies as a feasible proposition. This type of work is generally very time consuming, although not necessarily expensive. Until this is done, the many well-documented exotic species will continue to lure potential culturists away from indigenous species. Although some of the needed trials have begun in francophone Africa, it is imperative that the necessary trials should begin throughout Africa and Asia as soon as possible to allow the fullest utilization of indigenous fish for culture.

ACKNOWLEDGMENTS

I would like to thank Dr. Alan Bowmaker of the Division of Biological Sciences, University of Rhodesia for commenting on the initial manuscript and encouraging its completion. The tedium of compiling and coding the bibliography was shared by my wife Heather, who also acted as editor and critic in the later stages of production.

[1] ABDEL-MAGID,A.M. (1971)THE ABILITY OF CLARIAS LAZERA(PISCES) TO SURVIVE WITHOUT AIR BREATHING. J.ZOOL.LOND., 163(1):63-72.

[2] ABDEL-MAGID,A.M. BABIKER,M.M. (1975) OXYGEN CONSUMPTION AND RESPIRATION BEHAVIOUR OF THREE NILE FISHES. HYDROBIOLOGIA,46(4):359-367.

[3] ABOLARIN,M.O. (1971)A NEW SPECIES OF HENNUGUYA (MYXOSPORIDA,PROTOZOA) FROM WEST AFRICAN CATFISH, CLAZIAS LAZERA VAL., WITH A REVIEW OF THE GENUS HENNUGUYA THELON,1892. AFR.J.TROP.HYDROBIOL.FISH.1:93-105.

[4] ABOUL-ELA,I. (1973)THE EMBRYONIC AND LARVAL DEVELOPMENT OF THE NILE CATFISH, CLARIAS LAZERA, CUV.AND VAL. PROC. SEVENTH ARAB SCIENCE CONG.,CAIRO,EGYPT.

[5] ABOUL-ELA,I. AMER,F.I. EL BOLOCK,A.R. (1973) STUDIES ON SPAWNING AND SPAWNING BEHAVIOUR OF CLARIAS LAZERA CUV.AND VAL. IN FISH FARMS OF THE A.R.EGYPT.BULL.ZOOL.SOC.EGYPT,25:25-33

[6] ACCAME MURATORI,R. (1974)[THAILAND:TRADITIONS IN FISH CULTURE].RIV.ITAL. PISCIC.ITTIOPATOL.,9(1):5-6. (IN ITALIAN).

[7] ADEROUNMU,E.A. ADENIYI,F. (1972) CESTODES IN FISH FROM A POND AT ILE-IFE,NIGERIA. AFR.J.TROP.HYDROBIOL. FISH.,2(2):151-156.

[8] AFINOWI,M.A. (1973)SURVEILLANCE OF RESOURCES.ANN.REPORT DEPT.FISH., NIGERIA:76-81.

[9] AGER,L.A. (1975)INVASION OF A RENOVATED POND BY WALKING CATFISH, CLARIAS BATRACHUS (LINNAEUS), AND OTHER SPECIES. Q.J.FLA.ACAD.SCI. 38(3):183-191.

[10] AGRAWAL,V.P. RAJBANSHI,V.K. (1965) A COMPARATIVE STUDY OF GUSTATORY CUTANEOUS SENSE ORGANS OF A FEW FRESH-WATER FISHES. J.ZOOL.SOC. INDIA 17:1-7.

[11] AGRAWAL,V.P., VERMA,S.R. (1966) THE DIGESTIVE PHYSIOLOGY OF THE STOMACH OF A FRESH WATER TELEOST. ZOOL.JBH(ALLG.ZOOL.), 72(2/3):309-315

[12] AGRAWAL,V.P. SWARUP,R. (1967) THE CIRCULATORY SYSTEM OF CLARIAS BATRACHU LINN. AGRA UNIV.J.RES. 16:89-98.

[13] AGRAWAL,V.P. TYAGI ,A,P. (1969) FOOD,FEEDING HABITS AND ALIMENTARY CANAL OF FRESH WATER FISHES OF MUZAFFARNAGAR.AGRA.UNIV.J. RES(S)XVIII(1):15-28.

[14] AGRAWAL,V.P. (1975)SEASONAL VARIATION IN THE ACTIVITY OF DIGESTIVE ENZYMES IN 3 TELEOST FISHES.IND.SCI.CONG.ASSOC. PROC.62:157-158.

[15] AGRAWAL,V.P., SASTRY,K.V., KAUSHAB,S.K.S. (1976) DIGESTIVE ENZYMES OF 3 TELEOST FISHES.ACTA PHYSIOL ACAD.SCI.HUNGARICAE.46(2):93-98.

[16] AHL,E. (1934)[MORE ON THE FISHES OF LAKE TOBA IN SUMATRA.] SITZBER.GES.NATURF.FR.BERLIN.1934 (4-7):235-238.

[17] AHSAN,S.N., AHSAN,H. (1974) DEGENERATIVE CHANGES IN THE TESTES OF CLARIAS BATRACHUS CAUSED BY CADMIUM CHLORIDE.IND.J.ZOOT. 15(1):39-43.

[18] AHSAN,S.N., AHSAN,J. (1975) CHANGES IN LIVER GLYCOGEN IN STARVED AND NORMALLY FED GROWTH HORMONE TREATED CLARIAS BATRACHUS.ANN. ZOOL.(AGRA,INDIA).11(3):53-58.

[19] ALEXANDER,R.MCN. (1964)THE STRUCTURE OF THE WEBERIAN APPARATUS IN SILURIDS.PROC.ZOOL.SOC.LOND.142:419-440.

[20] ALEXANDER,R.MCN. (1966)STRUCTURE AND FUNCTION IN THE CATFISH.J.ZOOL., LOND.148:88-152.

[21] ALFRED,E.R. (1964)NOTES ON A COLLECTION OF FRESHWATER FISHES FROM

PENANG .BULL.NAT.MUS.SINGAPORE 32:143-154.

[22] ALFRED,E.R. (1966)THE FRESH-WATER FISHES OF SINGAPORE. ZOOL.VERH., LEIDEN 78:1-68.

[23] ALLEN,R.M. (1964)FISHERIES STATISTICS (NATURAL WATERS).REPORT OF MINISTRY OF LANDS AND NATURAL RESOURCES.PP94.

[24] ALLEN,R.M. (1965)MWERU-LUAPULA: FISHERIES STATISTICS-1963. FISH.RES.BULL.ZAMBIA 1963-1964.PP75-96.

[25] ALLEN,R.M. (1965)A PRELIMINARY REPORT OF DEPARTMENTAL NETTINGS IN LAKE KARIBA.FISH.RES.BULL.ZAMBIA.1963-1964. PP116-142.

[26] AL-GAUHARI,A.E.I. (1958)ON THE BLOOD SUGAR IN CLARIAS LAZERA.C&V. Z.VERGL PHYSIOL.41:26-34.

[27] AL-GAUHARI,A.E.I. (1960)THE MORPHOLOGY OF THE PANCREAS IN TELEOSTEI WITH SPECIAL REFERENCE TO THE NILE FISH CLARIAS LAZERA (C&V). ZOOL.ANZ. 165:63-70.

[28] AL-HUSSAINI,A.H. KHOLY,A.A. (1954) ON THE FUNCTIONAL MORPHOLOGY OF THE ALIMENTARY TRACT OF SOME OMNIVOROUS TELEOST FISH.PROC. EGYPT.ACAD.SCI.9:7-39.

[29] AL-HUSSAINI,A.H. LUFTY,R.G. (1958) ON THE MICROSCOPIC ANATOMY OF THE SKIN OF FISHES. I. THE HISTOLOGY OF THE SKIN OF SOME SILURIDS OF THE NILE. AIN.SHAMS. SCI.BULL. CAIRO 3: 215-263.

[30] AMATYAKUL,C. (1970)RICE FIELD FISH CULTURE AND CLARIAS CULTURE IN THAILAND, FAO FISH.CULT.BULL 2(3):10.

[31] AMATYAKUL,C. (1972)PARASITES OF POND RAISED CLARIAS IN THAILAND. FAO AQUACULTURE BULL. 5(1):16.

[32] ANGELOPULO,V. (1947)THE ANATOMY OF CLARIAS GARIEPINUS(BURCHELL).M.SC. THESIS,UNIVERSITY OF PRETORIA,RSA.65PP.

[33] ANON (1969)COUNTRY REPORT:VIETNAM.IN:PROC.OF THE INTERNAT. SEMINAR ON POSSIBILITIES AND PROBLEMS OF FISHERIES DEVELOPMENT IN SOUTHEAST ASIA.BERLIN, SEPT. 1968:310-315.

[34] ANON (1969)RESTRICTIONS ON WALKING CATFISH PROPOSED. COMML FISH. REV.,31(10):6.

[35] ANON (1970)'WALKING'CATFISH POPULAR IN THAILAND. AM.FISH FARMER,1:27.

[36] ANON (1970)SCIENTIFIC WORK OF THE ORGANIZATION. ANN.REP. EAST AFR.FISH.RES.ORG.1969:5-11.

[37] ANON (1971)FAO'S CURRENT FIELD PROJECTS IN AQUACULTURE. FAO AQUACULT.BULL.3(2):17-19.

[38] ANON (1975)BARBEL FEAST.WILD RHODESIA.NO.6:42.

[39] APRIETO,V.L. (1974)EARLY DEVELOPMENT OF CLARIAS MACROCEPHALUS GUNTER REARED IN THE LABORATORY (PISCES:CLARIIDAE).DEP. MAR.FISH.,UNIV.PHILIPPINES,QUEZON CITY,TECH.REP.NO 3, PP11.

[40] ARAI,R. HIRANO,H. (1974) FIRST RECORD OF THE CLARIID CATFISH,CLARIAS FUSCUS,FROM JAPAN. JAP.J.ICTHYOL.21(2):53-60.

[41] ARAMBOURG,C. [(1947)CONTRIBUTION TO THE GEOLOGIC AND PALEONTOLOGIC STUDY OF LAKE RUDOLPH BASIN AND THE LOWER OMO VALLEY I. PALEONTOLOGY.'MUS.NAT.HIST.NAT.PARIS, MISSION SCI. DE L'OMO 1932-1933 1(3):469-489.

[42] ARNOLD,J.P. AHL,E. (1936) [FRESHWATER FISH].BUSTAV WENZEL & SOHN, BRAUNSCHWEG DRUCK GUSTAV SOHN.GERMANY.PP.241-242 (IN GERMAN)

[43] BAILEY,R.G. (1966)THE DAM FISHERIES OF TANZANIA.E.AFR.AGRIC.&FOR.J. XXXII(1):1-15.

[44] BAILEY,R.G. (1969)THE NON-CICHLID FISHES OF THE EAST FLOWING RIVERS OF TANZANIA,EAST AFRICA.REV.ZOOL.BOT.AFR.,LXXX-1-2:170-199.

[45] BALON,E.K. (1972)POSSIBLE FISH STOCK SIZE ASSESSMENT AND AVAILABLE PRODUCTION SURVEY AS DEVELOPED ON LAKE KARIBA.AFR.J.TROP.HYDROBIOL.FISH.2(1):45-73

[46] BALON,E.K. (1973)RESULTS OF FISH POPULATION SIZE ASSESSMENTS IN LAKE KARIBA COVES(ZAMBIA),A DECADE AFTER THEIR CREATION.IN:MAN MADE LAKES:THEIR PROBLEMS AND ENVIRONMENTAL EFFECTS.GEOPHYS.MONO.SER.17:149-158

[47] BALON,E.K. COCHE,A.G.(ED) (1974) LAKE KARIBA A MAN - MADE TROPICAL ECOSYSTEM IN CENTRAL AFRICA:FISH PRODUCTION OF A TROPICAL ECOSYSTEM.DR.JUNK,THE HAGUE,1974.

[48] BANERJEE,D.K. CHATTERJEA,J.B. (1963) VITAMIN B-12 CONTENT OF SOME ARTICLES OF INDIAN DIETS AND THE EFFECT OF THE COOKING ON IT.BR.J. NUTR.17:385-389

[49] BANERJEE,D.K. CHATTERJEA,J.B. (1963) VITAMIN B-12 CONTENT OF INDIAN FISHES AND THE EFFECT OF BOILING ON ITS AVAILIBILITY.BULL. CALCUTTA SCH.TROP.MED.HYG.11: -12.

[50] BANERJEE,D.K. CHATTERJEA,J.B. (1963) EFFECT OF COOKING ON THE AVAILABILITY OF FOLIC ACID IN INDIAN FISHES.BULL.CALCUTTA SCH TROP.MED. HYG.11:49-50.

[51] BANERJEE,T.K. MITTAL,A.K. (1975) HISTOCHEMISTRY AND THE FUNCTIONAL ORGANIZATION OF THE SKIN OF A 'LIVE-FISH' CLARIAS BATRACHUS (LINN) (CLARIIDAE,PISCES) . MIKROSKOPIE 31(11-12):333-349

[52] BANERJEE,T.K., MITTAL,A.K. (1975) HISTOCHEMISTRY AND THE FUNCTIONAL ORGANIZATION OF SKIN OF A LIVE FISH CLARIAS BATRACHUS CLARIIDAE PISCES.MIKROSKOPIE.31(11-12):333-349.

[53] BANO,Y. (1975)VARIATIONS IN THE CHEMICAL COMPOSITION OF DIFFERENT SECTIONS OF THE FLESH OF CLARIAS BATRACHUS.IND.J.ZOOL.3(1-2):39-42.

[54] BANO,Y. (1977)SEASONAL VARIATIONS IN THE BIOLOGICAL COMPOSITION OF CLARIAS BATRACHUS.PROC.IND.ACAD.SCI.SECTION-B. 85(3):147-155.

[55] BARDACH,I.E. RYTHERJ.H. (1968) THE STATUS AND POTENTIAL OF AQUACULTURE, PARTICULARLY FISH CULTURE.US.DEPT.COMM.,NAT.TECH. INFO.SER.PUB.,PB177768.SPRINGFIELD,VA.PP.225.

[56] BARD,J. (1975)[PROGRESS IN THE TECHNOLOGY OF FISH CULTURE]. NOTES ET DOCUMENTS SUR LA PECHE ET LA PISCICULTURE. CENTRE TECHNIQUE FORESTIER TROPICAL. NOUVELLE SERIE NO 10:31-419.(IN SPANISH).

[57] BARD,J. DE KIMPE,P. LESSENT,P. (1975) [NEW FISHES AND NEW METHODS OF CULTIVATION IN AFRICA].IN:FAO/CIFA SYMPOSIUM ON AQUACULTURE IN AFRICA. CIFA/75/SR2. PP7.(IN FRENCH).

[58] BARNARD,K.H. (1943)REVISION OF THE FRESHWATER FISHES OF THE S.W.CAPE REGION.ANN.SA.MUS.XXXVI.

[59] BAYLIS,H.A. (1923)REPORT ON A COLLECTION OF PARASITIC NEMATODES MAINLY FROM EGYPT.PART 111.CAMALLANIDAE,ETC. WITH A NOTE ON PROBSTMAYRIA AND AN APPENDIX ON ACANTHOCEPHALA.PARASITOLOGY,15(1):24-38.

[60] BAYLIS,H.A. (1928)SOME PARASITIC WORMS MAINLY FROM FISHES FROM LAKE TANGANYIKA.ANN.MAG.NAT.HIST.SER.10,1(4): 552-562.

[61] BAYLIS,H.A. (1930)MISSION SAHARIENNE ALGIERAS-DRAPER,1927-1928. PARASITIC NEMATODES.BULL.MUS.NATN.HIST.NAT.PARIS 2E SERIE,2(1):117-130.

[62] BAYLIS,H.A. (1940)ON A FURTHER COLLECTION OF PARASITIC WORMS FROM THE BELGIAN CONGO. ANN. MAG.NAT.HIST., SER. 29(36):401-417.

[63] BEATTY,D.M.F. (1971)CASH SURPLUSES OF FISHING UNITS ON LAKE MWERU IN 1966.FISH.RES.BULL.ZAMBIA.5:351-377.

[64] BEAUCHAMP,R.S.A. (1960)THE COMMERCIAL VALUE OF STUDIES ON FISH BEHAVIOUR WITH PARTICULAR REFERENCE TO LAKE VICTORIA. PROC. INDO-PAC.FISH.COUNC.,8,SECT.3:97-99.

[65] BELL-CROSS,G. (1960)OBSERVATIONS ON THE MOVEMENTS OF FISH IN A FISH LADDER IN NORTHERN RHODESIA.IN:THIRD SYMPOSIUM ON HYDROBIOLOGY AND INLAND FISHERIES-PROBLEMS OF MAJOR LAKES, 1960.CSA.PUB.NO.63:13-125

[66] BELL-CROSS,G. (1961)SUNKEN LAKES IN THE NDOLA DISTRICT.NORTH.RHOD.J. 4:433-439.

[67] BELL-CROSS,G. (1965)ADDITIONS AND AMENDMENTS TO THE CHECKLIST OF THE FISHES OF ZAMBIA. PUKU 3:29-43.

[68] BELL-CROSS,G. (1965)MOVEMENT OF FISH ACROSS THE CONGO ZAMBEZI WATER SHED IN THE MWINILUNGA DISTRICT OF NORTHERN RHODESIA. PROC.CENTRAL AFR. SC.CONGRESS 1963. PERGAMMON PRESS,LONDON.

[69] BELL-CROSS,G. (1968)THE DISTRIBUTION OF FISHES IN CENTRAL AFRICA.FISH RES.BULL.,ZAMBIA.4:3-20.

[70] BELL-CROSS,G. (1968)PHYSICAL BARRIERS SEPARATING THE FISHES OF THE KAFUE AND MIDDLE ZAMBEZI RIVER SYSTEMS.FISH.RES. BULL.,ZAMBIA.4:97-98.

[71] BELL-CROSS,G. (1971)WEIR FISHING ON THE CENTRAL BAROTSE FLOOD PLAIN IN ZAMBIA.FISH.RES.BULL.ZAMBIA(5):331-340.

[72] BELL-CROSS,G. (1974)A FISHERIES SURVEY OF THE UPPER ZAMBEZI RIVER SYSTEM.OCC.NATN.MUS.RHOD.B5(5):279-338.

[73] BELSARE,D.K., MURTHY,P.S.R. (1968) THE EFFECTS OF HYPOPHYSECTOMY AND GOITROGEN TREATMENT ON OVARIAN COMPENSATORY HYPERTROPHY IN CLARIAS BATRACHUS L. ADVAN.ABSTR,CONTRIB.FISH AQUATIC SCI. INDIA.11(4):30.

[74] BELSARE,D.K. BELSARE,S.G. MURTHY,P.S.R. (1970) HYPOTHALAMO-HYPOPHYSIAL NUROSECRETORY SYSTEM IN NORMAL AND HYPOPHSECTOMISED CATFISH, CLARIAS BATRACHUS L. ACTA ZOOL., STOCKH. 51:219-228

[75] BELSARE,D.K. (1973)ON THE EVOLUTION OF TESTICULAR ENDOCRINE TISSUE IN SOME TELEOSTS. Z.MIKROSK.ANAT.FORSCH.87(5/6) :610-618.

[76] BELSARE,D.K. (1973)COMPARATIVE ANATOMY AND HISTOLOGY OF THE CORPUSCLES OF STANNIUS IN TELEOSTS. Z.MIKROSK. ANAT.FORSCH.,87(4):445-456.

[77] BELSARE,D.K. (1974)STUDIES ON THE DEVELOPMENT OF ENDOCRINE GLANDS IN FISHES.III MORPHOGENESIS OF THE ENDOCRINE PANCREAS IN CLARIAS BATRACHUS L. Z. MIKROSK-ANAT.FORSCH., LEIPZIG. 88(6):981-986.

[78] BELSARE ,D.K. (1974)MORPHOLOGICAL TYPES OF THE PREOPTICO-NEURO-HYPO-PHYSIAL NEUROSECRETORY PATHWAYS IN SOME TELEOSTS. Z.MIKROSK .ANAT.FORSCH.,LEIPZIG. 98(6):1029-1035.

[79] BELSARE,D.K. (1974)STUDIES ON THE DEVELOPMENT OF ENDOCRINE GLANDS IN FISHES.IV.DEVELOPMENT OF THE GONADS IN CLARIAS BATRACHUS.ZOOL.JAHRB.,ABT.ANAT.ONTOG. TIERRE,93(2):165-174.

[80] BELSARE,D.K. (1974)STUDIES ON THE DEVELOPMENT OF ENDOCRINE GLANDS IN FISHES. II DEVELOPMENT OF THE PINEAL ORGAN IN CLARIAS BATRACHUS (L.) ANAT.ANZ.,136(3):269-273.

[81] BELSARE,D.K. (1975)HEMATOLOGICAL STUDIES IN TELEOSTS PART I A STUDY OF BLOOD IN SOME FRESH WATER TELEOSTS.ZOOL. POLONIAE.25(2/3):55-63.

[82] BELSARE,D.K. (1975)COMMENT ON THE OVARIAN ENDOCRINE TISSUE IN TELEOST SPECIES.ZOOL.POLONIAE.25(2/3):5-11.

[83] BELSARE,D.K. (1975)COMPARATIVE HISTO MORPHOLOGY OF THE MAUTHNER CELLS IN SOME FRESHWATER TELEOSTS. ZEITSCHRIFT FUER MIKROS.ANAT.FORSC.89(5):804-814.

[84] BELSARE,D.K. (1976)STUDIES ON THE DEVELOPMENT OF ENDOCRINE GLANDS IN FISHES. V. DEVELOPMENT OF THE THYROID GLAND IN

TWO SPECIES OF TELEOSTS. MORPHOL.JAHRB., 122(1):
118-122.

[85] BELSARE,D.K., (1976)
BELSARE,S.D.
LIVER PHOSPHOLIPID DISTRIBUTION IN HYPOPHYSECTOM-
IZED CATFISH CLARIAS BATRACHUS AND HETEROPNEUSTES
FOSSILIS.ENDOKRINOLOGIE.67(3):365-368.

[86] BERGSTRAND,E. (1970)
CORDONE,A.J.
EXPLORATORY BOTTOM TRAWLING IN LAKE VICTORIA.ANN
REP.EAFRO.,(1969):42-52.

[87] BERGSTRAND,E. (1971)[A FISHERY PROJECT IN LAKE VICTORIA-NYANZA],
ZOOL.REV. 33(1-2):3-16. (IN SWEDISH)

[88] BERGSTRAND,E. (1971)
CORDONE,A.J.
EXPLORATORY BOTTOM TRAWLING IN LAKE VICTORIA.
AFR.J.TROP.HYDROBIOL.FISH.,1(1):13-23.

[89] BEVERLEY-BURTON,M. (1962)SOME TREMATODES FROM CLARIAS SPP. IN THE
RHODESIAS, INCLUDING ALLOCREADIUM MAZOENSIS N.SP
AND EUMASENIA BANGWEULENSIS N.SP, AND COMMENTS
ON THE SPECIES OF THE GENUS ORIENTOCREADIUM
TUBANGUI,1931. PROC. HELMINTH. SOC.WASH,
29(2):103-115.

[90] BEVERLEY-BURTON,M. (1963)A NEW STRIGEID DIPLOSTOMUS (TYLODELPHYS)
MASHONENSE N.SP.(TREMATODA)DIPLOSTOMATIDAE
FROM THE GREY HERON, ARDEA CINEREA L. IN
SOUTHERN RHODESIA WITH AN EXPERIMENTAL
DEMONSTRATION OF PART OF THE LIFE CYCLE.
REVUE ZOOL. BOT. AFR, 68(3/4):291-308.

[91] BHATT,S.D. (1973)FURTHER STUDIES ON THE HISTOLOGY OF THE ISLETS
OF LANGERHANS IN A FRESH WATER TELEOST,CLARIAS
BATRACHUS(L.).ZOOL.BEITR.19(1):1-8.

[92] BHATTACHARYA,S. (1973)
GUPTA,P.D.
MUKHERJEE,S.
A COMPARATIVE STUDY OF HEAD AND TAIL KIDNEY
PEROXIDASE IN A FISH (CLARIAS BATRACHUS). COMP.
BIOCHEM.PHYSIOL.(BIOCHEM.),44B(3):693-700.

[93] BHATTACHARYA,S. (1975)
MUKHERJEE,S.
BHATTACHARYA,S.
TOXIC EFFECTS OF ENDRIN ON HEPATOPANCREAS OF THE
TELEOST FISH, CLARIAS BATRACHUS (LINN). INDIAN J.
EXP.BIOL.,13(2):185-186.

[94] BHATTACHARYA,S., (1975)
MUKERJEE,S.
VARIATIONS IN THE KIDNEY PEROXIDASE ACTIVITY IN
FISHES EXPOSED TO SOME INDUSTRIAL POLLUTANTS.IND.
SCI.CONG.ASSOC.PROC.62:157.

[95] BHATT,S.D., (1976)
KHANNA,S.S.
HISTO PATHOLOGY OF THE ENDOCRINE PANCREAS OF A
FRESH WATER FISH CLARIAS BATRACHUS EFFECTS OF
HYDROCORTISONE AND THYROXINE ADMINISTRATION.ACTA
BIOL.ACAD.SCI.HUNGARICAE.27(1):25-35.

[96] BISHOP,W.W. , (1970)
CHAPMAN,G.R.
EARLY PLIOCENE SEDIMENTS AND FOSSILS
FROM THE NORTHERN KENYA RIFT VALLEY.
NATURE. 226:914-918.

[97] BISHOP,W.W. (1971)
CHAPMAN,G.R.
HILL,A.
MILLER,J.A.
SUCCESSION OF CAINOZOIC VERTEBRATE ASSEMBLAGES
FROM THE NORTHERN KENYA RIFT VALLEY. NATURE,LOND.
233:389-394.

[98] BLEEKER,M.P. (1862)[ICTHYLOGICAL ATLAS OF THE DUTCH EAST INDIES.
VOL.II.SILUROIDES,CHAROIDES,AND
HETEROBRANCHOIDES.] DUTCH COLONIAL GOVERNMENT
AMSTERDAM.PP112.(IN FRENCH)

[99] BOAKE,B. (1864)ON THE AIR BREATHING FISH OF CEYLON.JOURN.OF THE
CEYLON BRANCH R.ASIATIC SOC.:128-142.

[100] BOESEMAN,M. (1975)SOME FAUNISTIC AND NOMENCLATURAL NOTATIONS

CONCERNING FISHES FROM THE NETHERLANDS AND THE NEARBY NORTH SEA. ZOOL. BIJDRAGEN.1975(17):44-47.

[101] BOSE, K.C. (1959)PINEAL ORGAN IN FISHES. CURR SCI.,28(2): 76-7.

[102] BOULENGER,G.A. (1886)ON NEW FISHES FROM THE LOWER CONGO.ANN.N.H. 55(XIX):148-149.

[103] BOULENGER,G.A. (1896)DESCRIPTIONS OF NEW FISHES FROM THE UPPER CONGO.ANN.NAT.HIST.(6)XVIII:309-311.

[104] BOULENGER,G.A. (1898)REPORT ON A COLLECTION OF FISHES MADE BY J.E. S.MOORE IN LAKE TANGANYIKA DURING HIS EXPEDITION 1895-1896.WITH AN APPENDIX BY J.E.MOORE. TR.Z.S. XV:1-30.

[105] BOULENGER,G.A. (1899)[FAUNA MATERIAL FROM THE CONGO.NEW FISHES FROM THE CONGO.PART V.CYPRINIDS,SILURIDS,CYPRINIDONTS ACANTHOPTERIDS].ANN.MUS.CONGO.ZOOL.I:97-128. (IN FRENCH)

[106] BOULENGER,G.A. (1899)EXHIBITION OF LIVING SPECIMENS OF THE HARMUT CLARIAS LAZERA,C.&V. P.Z.S.P718.

[107] BOULENGER,G.A. (1900)REVISION OF THE CLARIIDAE.P.Z.SOC.LOND.(2):1062-1097.

[108] BOULENGER,G.A. (1900)[FAUNA MATERIAL FROM THE CONGO.NEW FISHES FROM THE CONGO.PART VI.MOMYRIDS,CHARACIDS,CYPRINIDS, SILURIDS].ANN.MUS.CONGO,ZOOL.I:129-164. (IN FRENCH)

[109] BOULENGER,G.A. (1900)LIST OF FISHES COLLECTED BY MR.J.S.BUDGETT IN THE RIVER GAMBIA.P.Z.S. :511-516.

[110] BOULENGER,G.A. (1901)DIAGNOSES OF FOUR NEW FISHES DISCOVERED BY MR.J. E.S.MOORE IN LAKES ALBERT AND ALBERT EDWARD.ANN. NAT.HIST.(7)VII:12-14.

[111] BOULENGER,G.A. (1902)CONTRIBUTIONS TO THE ICHTHYOLOGY OF THE CONGO.I. ON SOME NEW FISHES FROM THE FRENCH CONGO.P.Z.S.: 234-237.

[112] BOULENGER,G.A. (1902)DESCRIPTIONS OF NEW FISHES FROM A COLLECTION MADE BY MR.E.DEGEN IN ABYSSINIA.ANN.NAT.HIST.(7) IX:421-439.

[113] BOULENGER,G.A. (1902)[ADDITIONS TO THE FISH FAUNA OF THE CONGO BASIN]. ANN.MUS.CONGO.ZOOL.II:19-57.(IN FRENCH)

[114] BOULENGER,G.A. (1903)DESCRIPTION OF A NEW SILURID FISH OF THE GENUS CLARIAS FROM BRITISH CENTRAL AFRICA. ANN.NAT. HIST.(7)XI:362-363.

[115] BOULENGER,G.A. (1903)DESCRIPTIONS OF NEW FRESHWATER FISHES FROM SOUTHERN CAMEROON.ANN.NAT.HIST.(7) XI:435-441.

[116] BOULENGER,G.A. (1904)ON THE TYPE SPECIMEN OF A WEST AFRICAN FISH CLARIAS LAEVICEPS GILL.P.Z.S.1904,(4):200-201.

[117] BOULENGER,G.A. (1905)A LIST OF FRESHWATER FISHES OF AFRICA. ANN.NAT. HIST. (7) XVI:36-60.

[118] BOULENGER,G.A. (1906)DESCRIPTION OF A NEW FISH OF THE GENUS CLARIAS FROM UGANDA. ANN.MAG.NAT.HIST.,LONDON 17:569

[119] BOULENGER,G.A. (1906)DESCRIPTIONS OF NEW FISHES DISCOVERED BY MR E. DEGEN IN LAKE VICTORIA. ANN.MAG.NAT.HIST.,LONDON 17 :433-452

[120] BOULENGER,G.A. (1907)DESCRIPTIONS OF THREE NEW FRESHWATER FISHES DISCOVERED BY MR G.L. BATES IN SOUTHERN CAMEROON. ANN.MAG.NAT.HIST.,LONDON,20:50-52

[121] BOULENGER,G.A. (1907)A REVISION OF THE AFRICAN SILURID FISHES OF THE SUB FAMILY CLARIIDAE. PROC.ZOOL.SOC.,LONDON : 1062-1097.

[122] BOULENGER,G.A. (1907)THE FISHES OF THE NILE.HUGH REES,LOND,PP278-292.

[123] BOULENGER,G.A. (1908)NOTE ON CLARIAS CAPENSIS C&V. ANN.NATAL GOVT. MUS 1(3):237-239.

[124] BOULENGER,G.A. (1909)DESCRIPTIONS OF NEW FRESHWATER FISHES DISCOVERED BY MR . G.L.BATES IN SOUTH CAMEROON. ANN.MAG.NAT. HIST. LONDON. 4:186-188.

[125] BOULENGER,G.A. (1911)ON A COLLECTION OF FISHES FROM THE LAKE NGAMI BASIN,BECHUANALAND. LONDON.TRANS.ZOOL.SOC.

18:399-418

[126] BOULENGER,G.A. (1911)DESCRIPTIONS OF THREE NEW FRESHWATER FISHES DISCOVERED BY MR G.L. BATES IN SOUTH CAMEROON. ANN.MAG.NAT.HIST.,LONDON.8(8):372-373.

[127] BOULENGER,G.A. (1911)CATALOGUE OF THE FRESHWATER FISHES OF AFRICA IN THE BRITISH MUSEUM (NATURAL HISTORY).VOL II. LONDON. PP 529.

[128] BOULENGER,G.A. (1911)DESCRIPTION OF A NEW AFRICAN FISH OF THE GENUS. CLARIAS FROM LAKE RUKWA.ANN.MAG.NAT.HIST.(8),8:54

[129] BOULENGER,G.A. (1913)DESCRIPTIONS OF FOUR NEW FISHES DISCOVERED BY MR G.L. BATES IN THE NYONG RIVER,SOUTH CAMEROON. ANN.MAG.NAT.HIST.,LONDON.12:67-70.

[130] BOULENGER,G.A. (1915)[DESCRIPTIONS OF NEW FISHES .II MORMYRIDES, KNERIIDES, CHARACINIDES, CYPRINIDES, SILURIDES.] BRUXELLES REV.ZOOL.AFRICAINE 4:162-171(IN FRENCH)

[131] BOULENGER,G.A. (1919)FRESH WATER FISHES OF AFRICA.2:221-278.

[132] BOULENGER,G.A. (1920)[FISHES CAUGHT IN THE BELGIAN CONGO ON THE EXPEDITION OF DR C.CHRISTY.] ANN.MUS. DU CONGO BELGE SER 1(2) FASC 4:1-38 (IN FRENCH).

[133] BOURN,D.M. (1974)THE FEEDING OF THREE COMMERCIALLY IMPORTANT FISH SPECIES IN LAKE CHILWA,MALAWI. AFR.J.TROP. HYDROBIOL.FISH.3(2):135-145.

[134] BOWMAKER,A.P. (1960)A REPORT ON THE LAKE KARIBA AREA AND THE ZAMBESI RIVER PRIOR TO INUNDATION: THE INITIAL EFFECTS OF INUNDATION WITH PARTICULAR REFERENCE TO THE FISHERIES. IN:REPORT ON THE TRAINING CENTRE ON FISHERY SURVEYS FOR THE COUNTRIES OF THE AFRICAN REGION AT TANGA, TANGANYIKA VOLII ESSAYS ON THE STATUS OF FISHERIES DEVELOPMENT IN SELECTED COUNTRIES AND TERRITORIES OF AFRICA.FAO/UN. PP100-128.

[135] BOWMAKER,A.P. (1963)CORMORANT PREDATION ON TWO CENTRAL AFRICAN LAKES. THE OSTRICH.XXXIV(1):2-26

[136] BOWMAKER,A.P. (1964)RESEARCH RESULTS, LAKE BANGWEULU . REP.JT.FISH.RES ORG.N.RHOD.NYASALAND NO.11(1961):6-20.

[137] BOWMAKER,A.P. (1968)SOME UPPER CONGO FISH WHICH OFFER A MEANS OF BIOLOGICAL CONTROL OF THE SNAIL VECTORS OF BILHARZIASIS. PROC.TRANS.RHOD. SCI. ASSOC.(52):28-37.

[138] BOWMAKER,A.P. (1968)PRELIMINARY OBSERVATIONS ON SOME ASPECTS OF THE BIOLOGY OF THE SINAMWENDA ESTUARY,LAKE KARIBA PROC.TRANS.RHOD.SCI.ASSOC.(53):3-8.

[139] BOWMAKER,A.P. (1969)FISH POPULATION FLUCUATIONS ASSOCIATED WITH ANADROMESIS AT MWENDA RIVER MOUTH, LAKE KARIBA. NEWS LETT.LIMNOL.SOC.SOUTH.AFR.NO.13(SUPP.):45-49

[140] BOWMAKER,A.P. (1971)POTAMODROMESIS IN THE MWENDA RIVER,LAKE KARIBA. SYMPOSIUM ON MAN-MADE LAKES,KNOXVILLE,TENNESSE, 1971.

[141] BOWMAKER,A.P. (1973)POTAMODROMESIS IN THE MWENDA RIVER,LAKE KARIBA. IN:MAN MADE LAKES:THEIR PROBLEMS AND ENVIRONMENT-ALL EFFECTS.GEOPHYS.MONO.SER.17:159-164.

[142] BOWMAKER,A.P. (1973)AN HYDROBIOLOGICAL STUDY OF THE MWENDA RIVER AND ITS MOUTH,LAKE KARIBA.PH.D.THESIS.UNIV.OF WITWATERSRAND.RSA.PP576.

[143] BRAGANCA,B.DE M. (1944)NICOTINIC ACID CONTENT OF FISH.ANN.BIOCHEM.EXP. MED.4:41-44.

[144] BRONCHART-LANNOYE,M. THINES,G. (1975) [INFLUENCE OF SOME PHYSICAL FACTORS ON THE BEHAVIOUR OF FISHES WITH A DOUBLE SYSTEM OF RESPIRATION.]BULL.ACAD.R.BEG.,SCI .,61(5):524-542 (IN FRENCH) .

[145] BROWN,J.M. (1955)REPORT OF THE WORK AT THE FISH CULTURE FARM FOR THE YEAR 1955.REVIEW OF KENYA FISHERIES.NAIROBI.

[146] BRUTON,M.N. (1976)ON THE SIZE REACHED BY CLARIAS GARIEPINUS.J.LIMNOL SOC.STHN.AFR.2(2):57-58.

[147] BUNNASOPITH,T. (1959)[HANDLING OF THE FRY OF PLA DUK DAN,CLARIAS BATRACHUS]. THESIS UNIV.KASETSART, THAILAND. (IN THAI).

[148] CAMBRAY,J.A., JUBB,R.A. (1977) DISPERSAL OF FISHES VIA THE ORANGE-FISH TUNNEL, SOUTH AFRICA.J.LIMNOL.SOC.STHN.AFR.3(1):33-36.

[149] CAMPANA-ROUGET,Y. (1961)[NEMATODES OF FISH] EXPLO. HYDROBIOL. LACS KIVU, EDOUARD ET ALBERT (1952-54). BRUSSELLS, 3(4): 1-61. (IN FRENCH).

[150] CAREY,T.G. BELL-CROSS,B. (1967) BREEDING SEASONS AND QUANTITATIVE DATA ON GONADS AND OVA FOR CERTAIN FISH SPECIES.FISH.RES.BULL. ZAMBIA.(3):12-22.

[151] CAREY,T.G. (1967)FISH POPULATIONS IN LAGOON AND RIVERINE ENVIRONMENTS.FISH.RES.BULL.ZAMBIA.(3):9-12.

[152] CAREY,T.G. (1971)HYDROBIOLOGICAL SURVEY OF THE KAFUE FLOOD PLAIN. FISH.RES.BULL.ZAMBIA:245-295.

[153] CARREON,J.A. (1973)INDUCED BREEDING OF CLARIAS. FAO AQUACULT.BULL. 5(2):3

[154] CARREON,J.A. VENTURA,R.F. AMAZAN,G.J. (1973) NOTES ON THE INDUCED BREEDING OF CLARIAS MACROCEPHALUS GUNTHER. AQUACULTURE 2(1):5-16.

[155] CARREON,J.A. ESTOCAPIO,F.A. ENDEREZ,E.M. (1976) RECOMMENDED PROCEEDURES FOR INDUCED SPAWNING AND FINGERLING PRODUCTION OF CLARIAS MACROCEPHALUS GUNTER.AQUACULTURE,8:269-281.

[156] CARTER,G.S. (1931)AQUATIC AND AERIAL RESPIRATION IN ANIMALS. BIO. REV.6:1-35.

[157] CARTER,G.S. BEADLE,L.C. (1931) THE FAUNA OF THE SWAMPS OF THE PARAGUAYAN CHACO IN RELATION TO ITS ENVIRONMENT. II. RESPIRATORY ADAPTATION IN FISHES. J.LINN.SOC.LOND.ZOOL.37 :327-368.

[158] CARTER,G.S. (1957)AIR BREATHING.IN:THE PHYSIOLOGY OF FISHES.VOL.I. M.R. BROWN ED. ACADEMIC PRESS. PP65-79.

[159] CATCHY NGAKOUDOU,A. (1975)[PRESENT STATE OF AQUACULTURE IN THE CENTRAL AFRICAN REPUBLIC].IN:FAO/CIFA SYMPOSIUM ON AQUACULTURE IN AFRICA,GHANA,30SEPT. 1975, FAO-CIFA--75/SC-6. PP14. (IN FRENCH).

[160] CHAKRABARTE,J. SAHARYA,R. BELSARE,D.K. (1973) STRUCTURE OF THE GALL BLADDER IN SOME FRESHWATER TELEOSTS. Z.MIKROSK.ANAT.FORSCH .87(1):23-32.

[161] CHANDRASEKHAR,N. (1959)MULTIPLE HAEMOGLOBINS IN FISH. NATURE, LOND. 184(4699): 1652-3.

[162] CHANDRASEKHAR,K. (1961)THE STRUCTURE OF THE KIDNEY OF SOME TELEOSTEAN FISHES.REC.INDIAN MUS.59:479-495

[163] CHANDRASEKHAR,K. KHOSA,D. (1972) EFFECT OF MAMMALIAN ESTROGENS AND PROGESTERONE ON THE HYPOTHALAMO-ADENOHYPOPHYSIAL COMPLEX OF OVARIECTOMISED CATFISH CLARIAS BATRACHUS (LINN.).PROC. INDIAN ACAD.SCI(B) 76(6):240-250.

[164] CHANDRASEKHAR,K., KHOSA,D. (1973) EFFECT OF ELECTRICAL STIMULATION OF THE OLFACTORY MUCOSA ON THE HYPOTHALAMIC NEURO SECRETORY SYSTEM AND ITS IMPACT ON THE OVARIES OF 2 FRESHWATER TELEOSTEAN FISHES CLARIAS BATRACHUS AND HETEROPNEUSTES FOSSILIS.IND.J.ZOOT.14(2):81-88.

[165] CHANDHURI,H. (1976)PLANT SOURCE MATERIAL AS FOOD FOR CLARIAS BATRACHUS. FAO AQUACULT.BULL.8(1):4.

[166] CHAPMAN,D.W. MILLER,W.H. DUDLEY,R.G. (1971)

SCULLY,R.J. ECOLOGY OF THE FISHES IN THE KAFUE RIVER.CENTRAL FISHERIES RESEARCH INSTITUTE,ZAMBIA.FS:SF/ZAM 11 TECH.REP.2:1-66.

[167] CHATTERJI,R.C. (1933)ON THE TREMATODE PARASITES OF A RANGOON SILUROID FISH CLARIAS BATRACHUS (LINN.1785).BULL. ACAD.SCI.ALLAHABAD.3:33-40.

[168] CHITRAY,B.B. SAXENA,D.B. (1962) FUNCTIONAL ANATOMY OF THE DIGESTIVE ORGANS OF FRESH-WATER FISHES OF INDIA: HETEROPNEUSTES FOSSILIS (BLOCH) AND CLARIAS BATRACHUS (LINNAEUS) PART I. MORPHOLOGY. VESTN.CESKOSL.ZOOL. SPOLEC 26:148-159.

[169] CLAY,J.D. (1972)A PRELIMINARY SURVEY ON THE FISHERIES OF SANDRIVER LAKE,1972.OCC.PAP.NO.1,FISH.SECT.MIN. OF AGRIC.SWAZILAND.PP11.

[170] CLAY,D. (1976)AN INVESTIGATION INTO THE DISTRIBUTION OF FISHES IN SWAZILAND.REV.ZOOL.AFR.90(3):547-558.

[171] CLAY,D. (1977)BIOLOGY OF THE TROPICAL CATFISH(FAMILY: CLARIIDAE) WITH SPECIAL EMPHASIS ON ITS SUITABILITY FOR CULTURE (INCLUDING A BIBLIOGRAPHY OF THE CLARIIDAE AND RELATED TOPICS).FISH.MAR.SERV.MS REP.(IN PRESS).

[172] CLAY,D. (1977)PRELIMINARY OBSERVATIONS ON SALINITY TOLERANCE OF CLARIAS LAZERA FROM ISRAEL.BAMIDGEH.29(3):102-109

[173] CLAY,D. (1978)SEXUAL MATURITY AND FECUNDITY OF THE AFRICAN CATFISH (CLARIAS GARIEPINUS) WITH AN OBSERVATI ON THE SPAWNING BEHAVIOUR OF THE NILE CATFISH (CLARIAS LAZERA).J.LINN.SOC.(IN PRESS).

[174] CLEMENTS,B.N. (1968)AESTIVATING BARBEL. LAMMERGEYER.8:55-56.

[175] CLOUDSLEY-THOMPSON,J.L. (1959)THE SIGNIFICANCE OF CANNIBALISM.NEW SCIENTIST, 5(130):1068-1070.

[176] COBB,R. (1971)WALKING CATFISH.AQUARIUM,N.J.5(2):45-48.

[177] COCHE,A.G. (1968)DESCRIPTION OF PHYSIO-CHEMICAL ASPECTS OF LAKE KARIBA,AN IMPOUNDMENT IN ZAMBIA-RHODESIA. FISH. RES.BULL.,ZAMBIA.5(200-267.

[178] COCKSON,A. (1972)NOTES ON THE ANATOMY,HISTOLOGY,AND HISTOCHEMISTRY OF THE RESPIRATORY TREE OF CLARIAS MOSSAMBICUS. ZOOL.BEITR.18(1):101-108.

[179] COCKSON,A. BOURNE,D. (1972) ENZYMES IN THE DIGESTIVE TRACT OF TWO SPECIES OF EURYHALINE FISH. COMP. BIOCHEM. PHYSIOL. (PHYSIOL.), 41 (4A):715-718.

[180] COKE,M. (1968)DEPTH DISTRIBUTION OF FISH ON A BUSH CLEARED AREA OF LAKE KARIBA, CENTRAL AFRICA. TRANS. AMER.FISH.SOC.97(4):460-465.

[181] CORBET,P.S. (1957)SOME EFFECTS OF SIMULIUM CONTROL BY INSECTICIDE ON THE FEEDING HABITS OF INSECTIVOROUS FISHES. IN:SECOND SYMPOSIUM ON AFRICAN HYDROBIOLOGY AND INLAND FISHERIES,1956.CSA.PUB.NO.25:81.

[182] CORBET,P.S. (1958)EFFECTS OF SIMULIUM CONTROL ON INSECTIVOROUS FISHES. NATURE,LONDON.181:570-571.

[183] CORBET,P.S. (1958)SOME EFFECTS OF DDT ON THE FAUNA OF THE VICTORIA NILE. REV. ZOOL. BOT. AFR. 57:73-95

[184] CORBET,P.S. (1959)THE FOOD ON NON-CICHLID FISHES IN LAKE VICTORIA. ANN.REP.EAFFRO.(1958):21-37.

[185] CORBET,P.S. (1960)BREEDING SITES OF NON-CICHLID FISHES OF LAKE VICTORIA.NATURE.LOND.4737:616-617.

[186] CORBET,P.S. (1961)THE FOOD OF NON-CICHLID FISHES IN THE LAKE VICTORIA BASIN,WITH REMARKS ON THEIR EVOLUTION AND ADAPTATION TO LACUSTRINE CONDITIONS.PROC.ZOOL SOC.LOND. 136:1-101.

[187] CORBET,S.A. GREEN,J. GRIFFITH,J. BETNEY,E. (1973) ECOLOGICAL STUDIES ON CRATER LAKES IN WEST

CAMEROON LAKES KOTTO AND MBOANDONG.J.ZOOL.170 (3):309-324.

[188] CORDONE,A.J. KUDHONGANIA,B.W. (1970) STOCK EVALUATION AND ASSESSMENT. REP.E.A.F.F.R.O. :5-14.

[189] COTT,H.B. (1954)THE STATUS OF THE NILE CROCIDILE IN UGANDA. UGANDA J.18:1-12.

[190] COUREUR,J. SPAAS,J. (1957) [CONTRIBUTIONS TO THE BIOLOGY OF SEVERAL VORACIOUS FISH IN PONDS].IN:SECOND SYMPOSIUM ON AFRICAN HYDROBIOLOGY AND INLAND FISHERIES,1956. CSA.PUB.NO.28:103-107.(IN FRENCH)

[191] COURTENAY,W.R. OGILVIE,V.E. (1971) SPECIES POLLUTION. INTRODUCED ANIMALS AND THE BALANCE OF NATURE.ANIM.KINGD.74(2):22-28.

[192] COURTNAY,W.R. MILEY,W.W. (1975) RANGE EXPANSION AND ENVIRONMENTAL IMPRESS OF THE INTRODUCED WALKING CATFISH IN THE UNITED STATES. ENVIRON.CONSERV.2(2):145-148.

[193] CRASS,R.S. (1960)NOTES ON THE FRESHWATER FISHES OF NATAL WITH DESCRIPTIONS OF FOUR NEW SPECIES.ANN.NATAL.MUS. XIV(3):405-458.

[194] CRASS,R.S. (1966)FEATURES OF FRESHWATER FISH DISTRIBUTION IN NATAL AND A DISCUSSION OF CONTROLLING FACTORS.NEWS LETT LIMNOL.SOC.SOUTH AFR.7:31-35.

[195] CRASS, R.S. (1966)FRESHWATER FISH FAMILIES. NEWSL. LIMNOL. SOC. STH. AFR. NO.2:16-25.

[196] DAGET,J. (1954)THE FISH OF THE UPPER NIGER. MEM.INST.FRANC.AFR. NOIRE 36:1-391.

[197] DAGET,J. (1957)[SECOND PART. THE FISH . IN: THE WATERS AND FISH OF THE UPPER VOLTA.] BLANC,M. AND DAGET,J. MEM. INST.FRANC AFR.N. 50:95-169 (IN FRENCH).

[198] DAGET,J. (1961)[REMAINS OF FISH IN QUATERNARY SAHARA.] BULL.INST FRANC. AFR. NOIRE 23A: 182-191 (IN FRENCH)

[199] DAGET,J. (1961)[THE NATIONAL PARK NIOKOLA-KOBAI PART II. FISH.] MEM.INST.FRANC.AFR. NOIRE 62:327-361 (IN FRENCH).

[200] DAGET,J. (1962)[FISH OF FONTA DIALON AND OF LOWER GUINEA].MEM. INST.FR.AFR.NOIRE.65:1-210.(IN FRENCH)

[201] DAGET,J. ILTIS,A. (1965) [FISH FROM THE IVORY COAST (FRESHWATER AND BRACKISH).] MEM.INST.FRANC.AFR.NOIRE 74:1-385 (IN FRENCH)

[202] DAGET,J. PLANQUETTE,P. (1967) [ON SEVERAL FISHES OF THE IVORY COAST WITH A DESCRIPTION OF A NEW SPECIES, CLARIAS CAMOTTEI N.SP. (PISCES:SILURIFORMES, CLARIIDAE).]BULL.MUS. NA[N.HIST.NAT.PARIS (2)39:278-281.(IN FRENCH).

[203] DAGET, J. (1968)[A CONTRIBUTION TO THE STUDY OF FRESHWATER 4.FISH.].BULL. INST. FONDAM. AFR. NOIRE (A), 30(4):1582-9. (IN FRENCH)

[204] DAGET,J. (1969)[ON THE PRESENCE OF CLARIAS ALBOPUNCTATUS (PISCES, CLARIIDAE) IN THE TCHAD BASIN.] CAH.OFF.RECH.SCI. TECH.OUTRE-MER (SER.HYDROBIOL.)3(2):74-79.(IN FRENCH).

[205] DAMRONGRAT,S. KETSUNCHAI,P. (1962) [AN EXPERIMENT ON FEEDING OF FRY OF CLARIAS BATRACHUS WITH VARIOUS KINDS OF FOOD].IN:TECH. REP. OF BANGKHEN FISHERIES STATION,BANGKOK. (IN THAI).

[206] DAS,B.K. (1940)NATURE AND CAUSES OF EVOLUTION AND ADAPTATION OF THE AIR BREATHING FISHES. 27TH INDIAN SCI. CONG.27(2):215-260.

[207] DAS,B.K. (1945)LETHAL TEMPERATURES TO SOME AIR-BREATHING AND NON AIR-BREATHING FISHES OF INDIA.SCI.CULT.CALCUTTA 11:4.

[208] DAS,S.M. (1962)ON A COLLECTION OF FISHES FROM CEYLON WITH A DISCUSSION ON THEIR IMPORTANCE IN ZOOGEOGRAPHY. ICHTHYOLOGICA,KANPOR.1:1-12.

[209] DAS,S.M. (1962)ON A COLLECTION OF FISHES FROM CEYLON WITH A DISCUSSION ON THEIR IMPORTANCE IN ZOOGEOGRAPHY. ICTHYOLOICA, KANPUR 1: 1-12

[210] DAS, B.K. (1927)THE BIONOMICS OF CERTAIN AIR-BREATHING FISHES OF INDIA, TOGETHER WITH AN ACCOUNT OF THE DEVELOPMENT OF THEIR AIR BREATHING ORGANS. PHILOS. TRANS. B216:183-219.

[211] DATTA,N .C . SAHA,A.K. BAIDYA,A. (1975) COMPARATIVE STUDY OF OSTEOCRANIUM AND WEBERIAN APPARATUS OF CLARIAS BATRACHUS (LINN.) AND HETEROPNEUSTES FOSSILIS (BL.) (PISCES). ZOOL.ANZ. 195(5-6):374-386.

[212] DATTA,N.C., SAHA,A.K., BAIDYA,A. (1975) COMPARATIVE STUDY OF OSTEO CRANIUM AND WEBERIAN APPARATUS OF CLARIAS BATRACHUS AND HETEROPNEUSTES FOSSILIS.IND.SCI.CONG.ASSOC.PROC.62:204.

[213] DAVID,L. (1935)[THE DEVELOPMENT AND DISTRIBUTION OF THE CLARIIDAE].REV.ZOOL.BOT.AFR.28(1-2):78-147.(IN GERMAN)

[214] DAVID,L. POLL,M. (1937) [CONTRIBUTIONS TO THE ICHTHYOLOGICAL FAUNA OF THE BELGIAN CONGO.] ANN.MUS .CONGO (1)3:189-294. (IN FRENCH).

[215] DAYAL,J. (1938)STUDIES ON THE TREMATODE PARASITES OF FISHES.A NEW TREMATODE NEOGANADA BARABANKIAE (NOV.GEN. NOV.SPEC.) FROM CLARIAS BATRACHUS.PROC.IND.ACAD. SCI.SECT.B.7:132-137.

[216] DAYAL,J. GUPTA,S.P. (1953) A NEW TREMATODE GUAHATIANA BATRACHIING.N.SP. FROM THE INTESTINE OF A FRESHWATER FISH, CLARIAS BATRACHUS (LINN) OF THE SUB-FAMILY HAPLOCLADINAE ODHNER, 1911 PROC.40.INDIAN SC.CONG.(LUCKNOW 1953),PART III,SECT.7,PP190.

[217] DAY,F. (1868)OBSERVATIONS ON SOME OF THE FRESHWATER FISHES OF INDIA. PROC.ZOOL.SOC.LOND:274-288.

[218] DAY,F. (1871)ON THE FRESHWATER SILURIDS OF INDIA AND BURMA. P.Z.S:703-721

[219] DAY,F. (1877)GEOGRAPHICAL DISTRIBUTION OF INDIAN FRESHWATER FISHES.II.THE SILURIDAE.J.LINN.SOC.XIII:338-353.

[220] DAY,F. (1877)ON AMPHIBIOUS AND MIGRATORY FISHES OF ASIA.J.LINN SOC.XIII:198-214

[221] DAY,F. (1888)FISHES OF INDIA. TEXT SUPPL.LOND.PP.485.

[222] DEHADRAI,P.V. BANERJI,S.R. THAKUR,N.K. DAS,N.K. (1973) SEXUAL DIMORPHISM IN CERTAIN AIR BREATHING TELEOSTS. J.INLAND FISH.SOC.INDIA,BARRACKPORE, 5:71-77.

[223] DEHADRAI,P.V. (1976)INDUCED SPAWNING OF CLARIAS BATRACHUS. FAO AQUACULT.BULL.8(1):4.

[224] DENNY,P. (1969)LAKES OF SOUTH WESTERN UGANDA.I.PHYSICAL AND CHEMICAL STUDIES ON LAKE BUNYONYI. FRESHWAT. BIOL.2(2):143-158.

[225] DERANIYAGALA,P.E.P. (1932)THE NEMATOGNATHA OF CEYLON.PART I.CEYLON J.SCI. SECT.B.16:271-289.

[226] DERANIYAGALA,P.E.P. (1956)GIGANTISM IN THE CATFISH CLARIAS TEYSMANNI BRACHYSOMA (GUNTHER). SPOLIA ZEYLAN. 28:17-18.

[227] DERANIYAGALA,P.E.P. (1958)THREE NEW CYPRINOIDS,A NEW CATFISH AND VARIATION AMONG SOME CYPRINOIDS AND AN ANABANTOID OF CEYLON SPOLIA ZEYLAN.28(2):129-138.

[228] DEVARAJ,K.V. (1972)RESTING AND LOCOMOTORY BEHAVIOUR OF CLARIAS BATRACHUS(LINN.).CURR.SCI.41(7):268-269.

[229] DEVARAJ,K.V.
VARGUESE,T.J.
SATHANARAYANA RAO,G.P. (1972) INDUCED BREEDING OF THE FRESHWATER CATFISH CLARIAS BATRACHUS (LINN) BY USING PITUITARY GLANDS OF MARINE CATFISH. CURR.SCI. 41(24): 858-870.

[230] DE BONT,A.F. (1969)THE STATUS OF LIMNOLOGICAL KNOWLEDGE OF THE CONGO BASIN. IN: IBP REPORT OF THE REGIONAL MEETING OF HYDROBIOLOGISTS IN TROPICAL AFRICA. KAMPALA, UGANDA,20-28 MAY 1968:15-31.

[231] DE KIMPE,P. (1971)FEEDS FOR TILAPIA AND CATFISH. FAO AQUACULT.BULL 3(4):4.

[232] DE KIMPE,P. (1972)GROWTH AND PRODUCTION OF CLARIAS. FAO AQUACULT. BULL. 4(4):5.

[233] DE KIMPE,P.
MICHA,J.C. (1974) FIRST GUIDELINES FOR THE CULTURE OF CLARIAS LAZERA IN CENTRAL AFRICA. AQUACULTURE 4:227-248.

[234] DE KIMPE,P.
MICHA,J.C. (1975) [FIRST INFORMATION FOR THE REARING OF CLARIAS LAZERA IN CENTRAL AFRICA.] NOTES DOC.PECHE PISCIC. (NOUV.SER.) 11:1-35. (IN FRENCH).

[235] DE KIMPE,P. (1975)AQUACULTURE ECONOMICS IN TROPICAL AFRICA. IN: FAO/CIFA SYMPOSIUM ON AQUACULTURE IN AFRICA, GHANA,30 SEPT. 1975, FAO-CIFA--75/SR-3, PP15

[236] DE WITT,J.W. (1973)LAKE NASSER FISHERIES SURVEY AND TRAINING PROGRAMME,A.R.EGYPT. FAO-FI-DP/EGY-66/558/10, PP56.

[237] DIXIT,V.P. (1970)HISTOPHYSIOLOGICAL STUDIES OF THE INTERRENAL GLAND IN CLARIAS BATRACHUS LINN. ACTA ANAT. 77(2):310-318.

[238] DIXIT,V.P. (1970)EFFECT OF THYROID INHIBITORS ON THE THYROTROPHS AND THE HYPOTHALOMO-HYPOPHSIAL NEUROSECRETORY SYSTEM OF CLARIAS BATRACHUS L. INDIAN J.EXP.BIOL. 8:98-101

[239] DIXIT,V.P. (1970)NEUROSECRETION AND FEEDBACK MECHANISM IN CLARIAS BATRACHUS LINN.OVARIECTOMY AND ADMINISTRATION OF EXOGENOUS SEX HORMONES. CELLULE 68:213-221.

[240] DIXIT, V.P. (1971)THE KARYOMETRIC RESPONSE OF CAUDAL NEUROSECRETORY CELLS IN CLARIAS BATRACHUS TO SEX STEROIDS. GEN. COMP. ENDOCRINOL. 17(3):561-563.

[241] DIXIT,V.P. (1976)KARYOMETRIC STUDIES OF THE NUCLEUS PREOPTICUS IN FISH CLARIAS BATRACHUS AND TOAD BUFO ANDERSONII. ACTA.ANAT.95(2):182-189.

[242] DJHA,J.
TIWARY,N.P.
DANDOTIA,O.P. (1975) EFFECT OF STARVATION ON FAT CONTENT IN THE AIR BREATHING ORGANS OF AMPHIBIOUS FISHES ANABAS TESTUDINEUS AND CLARIAS BATRACHUS.J.BIOL.SCI. 18(2):81-84.

[243] DOBSON,G.E. (1874)NOTES ON THE RESPIRATION OF SOME SPECIES OF INDIAN FRESH WATER FISHES.PROC.ZOOL.SOC.LOND. :312-321.

[244] DONNELLY,B.G. (1966)SHOALING,COMMUNICATION AND SOCIAL HUNTING IN THE CATFISH CLARIAS NGAMENSIS. PISCATOR, (67): 54-55.

[245] DONNELLY,B.G.
CAULTON,M.S. (1969) A POSSIBLE TECHNIQUE FOR AGE DETERMINATIONS IN SOUTHERN AFRICA SILUROID FISHES. NEWSL. LIMNOL. SOC.S.AFR.12:13-15.

[246] DONNELLY,B.G.
RUSHWORTH,D. (1970) TOUR REPORT - CHIZARIRA GAME RESERVE.(MIMEO.

REPORT)INTERNAL REPORT OF DEPT.NAT.PARKS & WDLF. MAN.RHODESIA.PP5.

[247] DONNELLY,B.G. (1973)ASPECTS OF BEHAVIOUR IN THE CATFISH,CLARIAS GARIEPINUS(PISCES:CLARIIDAE) DURING PERIODS OF HABITAT DESICCATION. ARNOLDIA 6(9):1-7.

[248] DUBALE,M.S. GEORGE,C.J. (1941) A NOTE ON THE DEGENERATION OF GILLS IN AIR BREATHING FISHES. J.UNIV.BOMBAY(BIOL.),10(3) :80-85.

[249] DUBALE,M.S. (1951)A COMPARATIVE STUDY OF THE EXTENT OF GILL SURFACE IN SOME REPRESENTATIVE INDIAN FISHES AND ITS BEARING ON THE ORIGIN OF THE AIR BREATHING HABIT. J.UNIV.BOMBAY(BIOL.),29(5):90-102.

[250] DUBALE,M.S. (1955)PAUCITY OF OXYGEN IN THE MEDIUM AS A FACTOR IN THE EVOLUTION OF AIR BREATHING HABIT AMONG TELEOSTS. J.MAHARAJA SAYAJIRAO UNIV.BARODA, 4:63-72.

[251] DUBALE,M.S. (1959)A COMPARATIVE STUDY OF THE OXYGEN CARRYING CAPACITY OF THE BLOOD IN WATER AND AIR BREATHING TELEOSTS.J.MORPH.PHYSIOL.6(1):48-54.

[252] DUBALE,M,S. CHRISTIAN,E.G. (1964) FUNCTIONAL ANATOMY OF THE PELVIC FINS OF CERTAIN INDIAN FRESH - WATER SILUROIDS. J.UNIV. BOMBAY (B),32(3/5):104-116.

[253] DUBIOS, G. (1930)[TWO NEW SPECIES OF CLINOSTOMIDAE.] BULL. SOC. NEU CHATEL. SCI.NAT, 54:61-72. (IN FRENCH)

[254] DUTTA,S.K. (1924)ON A PECULIAR DISPOSITION OF THE LIVER AND THE KIDNEY IN THE FISH GENERA CLARIAS AND SACCOBRANCHUS. J.ASIAT.SOC.BENG. CALCUTTA 19:111-120.

[255] EISAWY,A. EL BOLOCK,A.R. (1975) STATUS OF AQUACULTURE IN THE ARAB REPUBLIC OF EGYPT. IN: FAO/CIFA SYMPOSIUM ON AQUACULTURE IN AFRICA. FAO/CIFA/75/SC 11. PP10.

[256] ELLIOTT,O. (1968)THE INLAND FISHERIES OF NIGERIA. M.SC. THESIS, UNIVERSITY OF WASHINGTON,USA.PP100.

[257] ELLIOTT,O.O. (1975)BIOLOGICAL OBSERVATIONS ON SOME SPECIES USED FOR AQUACULTURE IN NIGERIA.IN:FAO/CIFA SYMPOSIUM ON AQUACULTURE IN AFRICA,GHANA,30 SEPT 1975, FAO-CIFA--MK/SE.PP12.

[258] ELMOSSALAMI,E. SHERIF, I.H. (1964) ENCYSTED METACERCARIA IN THE MUSCLES OF CLARIAS LAZERA (ARMOUT CATFISH).CEYLON VET.J.12(3):50-53

[259] EL BOLOCK,A.R. KOURA, R. (1960) OBSERVATIONS ON AGE, GROWTH AND FEEDING HABITS OF CLARIAS LAZERA, CUVIER AND VALENCIENNES IN BARRAGE EXPERIMENTAL PONDS. NOTES HYDROBIOL. DEP.MINIST.AGRIC. U.A.R. (56):1-17.

[260] EL BOLOCK,A.R. (1969)CATFISH CULTURE. FAO FISH.CULT.BULL. 2(1):7.

[261] EL BOLOCK,A.R. (1972)THE USE OF VERTEBRAE FOR DETERMINING AGE AND GROWTH OF THE NILE CATFISH CLARIAS LAZERA(CUV& VAL.) IN: THE A.R.EGYPT.BULL.INST.OCEANOGR.FISH. UAR.2:53-82.

[262] EL BOLOCK,A.R. (1973)BREEDING AND CULTURE OF CLARIAS. FAO AQUACULT. BULL. 5(3-4):4.

[263] EL BOLOCK,A.R. (1975)STUDIES ON THE BIOLOGY AND CULTURE OF CLARIAS LAZERA IN EGYPTIAN PONDS. PH.D. THESIS.CAIRO.

[264] EL BOLOCK,A.R. (1975)REARING OF THE NILE CATFISH,CLARIAS LAZERA,TO MARKETABLE SIZE IN EGYPTIAN EXPERIMENTAL PONDS. IN: FAO/CIFA SYMPOSIUM ON AQUACULTURE IN AFRICA, GHANA,30SEPT. 1975, FAO-CIFA--75/SE-9, PP8.

[265] EL BOLOCK,A.R. EL SAMAGAWI,D. (1975) SOME DISEASES OF CULTIVATED FISHES IN EGYPT. IN:FAO/CIFA SYMPOSIUM ON AQUACULTURE IN AFRICA.

GHANA,30 SEPT.1975,FAO-CIFA/75/SE 14. PP4.

[266] EL ILA,A.
EL BOLOCK,A.R. (1972)
ON THE BIOLOGY AND CULTURE OF CLARIAS LAZERA (CUV.&VAL.)IN EGYPTIAN PONDS. SPAWNING AND DEVELOPMENT STAGES.

[267] ENAMI, M. (1955)MELANOPHORE CONCENTRATING HORMONE (MCH) OF POSSIBLE HYPOTHALAMIC ORIGIN IN THE CATFISH PARASILURUS. SCIENCE 121:36-37.

[268] ENSLIN,J.M.N. (1966)[A COMPARATIVE STUDY OF THE FEEDING HABITS OF CERTAIN FISH IN BARBERSPAN, TRANSVAAL]. M.SC. THESIS, POTCHEFSTROON UNIVERSITY FOR CHRISTIAN HIGHER EDUCATION. (IN AFRICAANS).

[269] EVERETT,G.V. (1971)SAMPLING FISH STOCKS IN THE KAFUE RIVER. FISH. RES. BULL. ZAMBIA,5:297-304.

[270] EVERETT,G.V. (1974)AN ANALYSIS OF THE 1970 COMMERCIAL FISH CATCH IN THREE AREAS OF THE KAFUE FLOODPLAIN.AFR.J.TROP. HYDROBIOL.FISH.3(2):147-159

[271] FAINE,A. (1953)[A CONTRIBUTION TO THE STUDY OF LARVAL FORMS OF TREMATODES FROM THE BELGIAN CONGO AND, ESPECIALLY THE LARVA OF SCHISTOSOMA MANSOMI.] MEM.INST.R.COLON.BELGE SECT.SCI.NAT.MED.22(5), :312.(IN FRENCH).

[272] FAO/UN. (1966)REPORT TO THE GOVERNMENT OF MALAWI ON FISH HANDLING,PRESERVATION AND DISTRIBUTION. BASED ON THE WORK OF Y.C.GILBERG.REP.FAO/UNDP(TA) (2224):PP52.

[273] FAO/UN. (1968)REPORT TO THE GOVERNMENT OF ZAMBIA ON FISHERY DEVELOPMENT IN THE CENTRAL BAROTSE FLOOD PLAIN. BASED ON THE WORK OF D.W.KELLEY.REP.FAO/UNDP(TA) (2554).PP83.

[274] FISCHER,J.G. (1883)[ABOUT A FEW REPTILES,AMPHIBIANS,AND FISHES, IN THE NATURAL HISTORY MUSEUM.]J.B.HAMB. I:1-39.(IN GERMAN).

[275] FISCHTHAL,J.H.
KUNTZ,R.E. (1959)
TREMATODE PARASITES OF FISHES FROM EGYPT. PART I. BASIDIODISCUS ECTORCHIS N.GEN.N.SP. AND SANDONIA SUDANENSIS MACLELLAND, 1957 (PARARAMPHLISTOMIDAE). PROC. HELMINTH SOC. WASH. 26(1): 32-37.

[276] FISCHTHAL,J.H.
KUNTZ,R.E. (1963)
TREMATODE PARASITES OF FISHES FROM EGYPT. PART V11. ORIENTOCREADIUM BATRACHOIDES TUBANGUI, 1931 (PLAGIORCHIDIDEA) FROM CLARIAS LAZERA, WITH A REVIEW OF THE GENUS AND RELATED FORMS. J.PARASIT. 49(3):451-464.

[277] FISCHTHAL,J.H.
THOMAS, J.D. (1968)
DIGENETIC TREMATODES OF SOME FRESHWATER AND MARINE FISHES FROM GHANA. PROC. HELMINTH SOC. WASH. 35(2):126-140.

[278] FISCHTHAL,J.H.
THOMAS,J.D. (1972)
DIGENETIC TREMATODES OF FISHES FROM THE VOLTA RIVER DRAINAGE SYSTEM PRIOR TO THE CONSTRUCTION OF THE VOLTA DAM AT AKOSOMBO IN MAY 1964.J. HELMIN.XLVI(1):91-106.

[279] FISCHTHAL,J.H (1973)THREE DIGENETIC TREMATODES OF CLARIAS MOSSAMBICUS PETERS(CLARIIDAE) FROM ETHIOPIA. PROC.HELMIN.SOC. WASH.40(1):166-167.

[280] FISH,G.R. (1956)SOME ASPECTS OF THE RESPIRATION OF SIX SPECIES OF FISH FROM UGANDA. J.EXP.BIOL.33:186-195.

[281] FLORKIN,M.(ED)
SCHEER,B.T.(ED) (1974)
CHEMICAL ZOOLOGY:DEUTEROSTOMIANS,CYCLOSTOMES, AND FISHES VOL VIII.ACADEMIC PRESS.LONDON.PP 682.

[282] FOWLER,H.W. (1915)NOTES ON NEMATOGNATHOUS FISHES. PHILADELPHIA PROC.ACAD.NAT.SCI. 67:203-243.

[283] FOWLER,H.W. (1930)THE FRESHWATER FISHES OBTAINED BY THE GRAY

AFRICAN EXPEDITION - 1929.WITH NOTES ON OTHER SPECIES IN THE ACADEMY COLLECTION.PROC.ACAD.NAT. SCI.PHILAD.82 :27-83.

[284] FOWLER,H.W. (1936)ZOOLOGICAL RESULTS OF THE GEORGE VANDERBILT AFRICAN EXPEDITION 1934 PART III- THE FRESHWATER FISHES. PROC.ACAD.NAT.SCI. PHILAD. 88:243-335.

[285] FOWLER,H.W. (1937)ZOOLOGICAL RESULTS OF THE THIRD DE SCHAUENSEE SIAMESE EXPEDITION. PART VIII. FISHES OBTAINED IN 1936. PROC.ACAD.NAT.SCI. PHILAD. 89:125-264

[286] FOWLER,H.W. (1949)RESULTS OF THE TWO CARPENTER AFRICAN EXPEDITIONS, 1946-1948.PI.II-THE FISHES.PROC.ACAD.NAT.SCI. PHILAD.101.:233-275.

[287] FRANCIS,R.C. (1974)RELATIONSHIP OF FISHING MORTALITY TO NATURAL MORTALITY AT THE LEVEL OF MAXIMUM SUSTAINABLE YIELD UNDER THE LOGISTIC PRODUCTION MODEL.J.FISH. RES.BD.CAN.31(9):1539-1542.

[288] FRECHKOP,S. (1954)[SOME GLANDS ADJACENT TO THE PECTORAL SPINES OF THE SILURID CLARIAS]BULL.MUS. HIST. NAT. BELGE. 30(36):1-7.(IN FRENCH)

[289] FRENKEL S.A.
IAKOVLEV,V.N. (1971)
[AMPHIBIAN FISH.] PRIRODA, MOSK. 5:123-125. (IN RUSSIAN)

[290] FRYER,G.
ILES,T.D. (1955)
PREDATION PRESSURE AND EVOLUTION IN LAKE NYASA. NATURE 176:470.

[291] FRYER,G. (1959)THE TROPHIC INTERRELATIONSHIP AND ECOLOGY OF SOME LITTORAL COMMUNITIES OF LAKE NYASA WITH SPECIAL REFERENCE TO THE FISHES,AND A DISCUSSION OF THE EVOLUTION OF A GROUP OF ROCK FREQUENTING CICHLIDAE. PROC.ZOOL.SOC.LOND.132:153-281.

[292] FRYER, G. (1959)A REPORT ON THE PARASITIC COPEPODA AND BRANCHIURA OF THE FISHES OF LAKE BANGWEULU (NORTHERN RHODESIA). PROC.ZOOL.SOC.LOND.132: 517 -550.

[293] FRYER,G. (1968)THE PARASITIC CRUSTACEA OF AFRICAN FRESHWATER FISHES:THEIR BIOLOGY AND DISTRIBUTION.J.ZOOL.156 (1):35-43.

[294] FUHRMANN,O.
BAER,J.G. (1925)
ZOOLOGICAL RESULTS OF THE THIRD TANGANYIKA EXPEDITION CONDUCTED BY DR.W.A.CUNNINGTON, 1904-1905. REPORT ON THE CESTODA. PROC.ZOOL. SOC. LOND. PART 1. : 79-100.

[295] FUKUDA,Y. (1971)THE REVIEW OF AIR BREATHING ORGANS OF FISHES, WITH SPECIAL REFERENCE TO SNAKEHEAD FISH, CHANA ARGUS (CANTOR). ZOOL.MAG.TOKYO ZOOL.SOC. 80(6):181-190.

[296] FUNAIOLI,U.
RONCATI,R. (1964)
[TRADITIONAL FISHES IN THE LOWER UEBI SCEBELI (SOMALIA)].RIV.AGR.SUBTROP. TROP. 58(4/6):250-72 (IN ITALIAN)

[297] FURTADO,J.I. (1963)A NEW CARYPHYLLAEID CESTODE,LYTOCESTUS PARVULUS SP.NOV.FROM A MALAYAN CATFISH. ANN.MAG.NAT.HIST. 6(62):97-106.

[298] FURTADO,J.I. (1973)INCIDENCE OF SOME HELMINTH PARASITES IN THE MALAYSIAN CATFISH CLARIAS BATRACHUS(LINN.). VERH.INT.VER.THEOR.ANGEW.LIMNOL.18(3):1674-1685

[299] GABIE,V. (1965)PROBLEMS ASSOCIATED WITH DISTRIBUTION OF FRESHWATER FISHES IN SOUTH AFRICA.S.AFR.J.SCI.61: 383-391.

[300] GAIGHER,I.G. (1969)[ASPECTS REGARDING THE ECOLOGY,GEOGRAPHY,AND TAXONOMY OF THE FRESHWATER FISH OF THE LIMPOPO AND INCOMATI RIVER SYSTEMS].D.SC.THESIS AFRIKAANS UNIVERSITY OF THE RAND,RSA. PP226-229. (IN AFRIKAANS)

[301] GAIGHER,I.G. (1972)

POTT,R.MCC. A CHECKLIST OF INDIGENOUS FISHES IN THE EAST FLOWING RIVERS OF THE TRANSVAAL.NEWS LETT.LIMNOL. SOC.SOUTH AFR.18:26-32.

[302] GEE,J.M. (1970)NOTES ON THE GENUS XENOCLARIAS(GREENWOOD). ANN. REP.EAST AFR.FISH.RES.ORGAN.1969:28-34

[303] GEORGE,T.T. (1975)INTRODUCTION AND TRANSPLANTATION OF CULTIVABLE SPECIES INTO AFRICA. IN:FAO/CIFA SYMPOSIUM ON AQUACULTURE IN AFRICA. CIFA/75/SR 7. PP25.

[304] GHOSH,E. (1934)AN EXPERIMENTAL STUDY OF THE ASPHYXIATION OF SOME AIR BREATHING FISHES OF BENGAL.J.PROC. ASIATIC SOC.BENGAL,29:327-332.

[305] GHOSH,A. KAR,A.B. (1952) SEASONAL CHANGES IN THE GONADS OF THE COMMON INDIAN CATFISH HETEROPNEUSTES FOSSILIS(BLOCH). PROC.ZOOL.SOC.BENG.5:29-50.

[306] GHOSH,A. KONAR,S.K. (1973) EFFECTS OF PHOSPHAMIDON ON THE ACTIVITY OF THE DIGESTIVE ENZYME INVERTASE IN THE CATFISH CLARIAS BATRACHUS(LINN.). J.INLAND FISH SOC. INDIA,BARRACKPORE,5:129-130.

[307] GIDEIRI,ABU,Y.B. (1961)FISHES OF THE BLUE NILE BETWEEN KHARTOUM AND ROSEIRES.REV.ZOOL.BOT.AFR.LXXV1(3/4):345-348.

[308] GILCHRIST,J.D. THOMPSON,W.W. (1913) THE FRESH WATER FISH OF SOUTH AFRICA.ANN.S.A.MUS. (11):321-463.

[309] GOEL,K.A. (1966)SOUND PRODUCTION IN CLARIAS BATRACHUS. COPEIA,(3 :622-624.

[310] GOEL,K.A. SASTRY,K.V. (1973) DISTRIBUTION OF THE ALKALINE PHOSPHATASE IN THE DIGESTIVE SYSTEM OF A FEW TELEOST FISHES. ACTA HISTOCHEM. 47(1):8-14.

[311] GOHAR,H.A.F. LATIF,A.F.A. (1963) GASTRIC AND INTESTINAL ABSORPTION OF OLIVE OIL AND OLEIC ACID IN CLARIAS LAZERA C&V. PUBL.MAR. BIOL.STA. AL-GHARDAGA 12:43-64.

[312] GOSWAMI,O.B. JAIN,S.P. KHANNA,S.S. (1971) ON THE MORPHOLOGY OF THE TELEOSTEAN BRANCHIOCEPHALIC VENEOUS SYSTEM. PROC.INDIAN ACAD. SCI.(B)73:15-19.

[313] GOSWAMI,O.B. JAIN,S.P. KHANNA,S.S. (1971) ON THE VISCERAL ARTERIAL SYSTEM OF SOME TELEOSTS PROC.INDIAN ACAD.SCI.(B)73(2):58-64.

[314] GOUGH,P.B. (1973)INVESTIGATION INTO THE PRACTICAL APPLICATION OF FISH FARMING.TRIBAL TRUST LAND DEVELOPMENT CORPORATION LIMITED,RHODESIA PP6.

[315] GRAHAM,M. (1929)THE VICTORIA NYANZA AND ITS FISHERIES.CROWN AGENTS FOR THE COLONIES.LONDON.PP .

[316] GRAHAM,P. (1964)DESTRUCTION OF BIRDS AND OTHER WILDLIFE, BY DIELDREX SPRAYING AGAINST TSETSE FLY IN BECHUANALAND. ARNOLDIA. 1(10):1-4.

[317] GRAHAM,J.B. (1971)AERIAL VISION IN AMPHIBIOUS FISHES. FAUNA (RANCHO MIRAGE,CALIF.) NO.3:14-23.

[318] GREENWOOD,P.H. (1950)QUATERNARY FISH-FOSSILS. EXPLOR. PARC NAT. ALBERT MISS. DE BRAVEOURT 4:1-80.

[319] GREENWOOD,P.H. (1955)REPRODUCTION IN THE CATFISH CLARIAS MOSSAMBICUS PETERS. NATURE,LOND.176(4480):516-518

[320] GREENWOOD,P.H. (1956)A NEW SPECIES OF CLARIALLABES(PISCES,CLARIIDAE) FROM THE NILE.PROC.ZOOL.SOC.LOND.127:555-564.

[321] GREENWOOD,P.H. (1956)THE FISHES OF UGANDA .UGANDA SOCIETY,KAMPALA.

[322] GREENWOOD,P.H. (1957)FISH REMAINS FROM THE MUMBA CAVE LAKE EYASI. IN: LEHMANN U. EINE JUNGPLEISTOZANE WIRBELTIER FAUNA AUS OSTAFRIKA. MITT.GEOL.(ST.) INST.HAMB. 26:125-130.

[323] GREENWOOD,P.H. (1957)THE REPRODUCTION OF CLARIAS MOSSAMBICUS IN LAKE VICTORIA.IN:SECOND SYMPOSIUM ON AFRICAN HYDROBIOLOGY AND INLAND FISHEREIS,1956.CSA.PUB. NO.25:77-78.

[324] GREENWOOD,P.H. (1958)A NEW GENUS AND SPECIES OF CATFISHES(PISCES(CLARIIDAE FROM THE DEEPER WATERS OF LAKE VICTORIA ANN.MAG.NAT.HIST.(13)1:321-325.

[325] GREENWOOD,P.H. (1961)A REVISION OF THE GENUS DINOPTERUS BLGR.(PISCES, CLARIIDAE)WITH NOTES ON THE COMPARATIVE ANATOMY OF THE SUPRA-BRANCHIAL ORGANS IN THE CLARIIDAE. BULL.BRIT.MUS.(NAT.HIST.)7(4):217-241.

[326] GREENWOOD,P.H. (1963)A COLLECTION OF FISHES FROM THE ASWA RIVER DRAINAGE SYSTEM,UGANDA.PROC.ZOOL.SOC.LOND.140(1): 61-74.

[327] GREENWOOD,P.H. (1973)
HOWES,G.J. FISH REMAINS.IN:EXCAVATIONS IN THE NECROPOLIS OF SALAMIS III. SALAMIS 5.PUBL.DEPT.ANTIQUITUS, NICOSIA CYPRUS:259-267.

[328] GREENWOOD,P.H. (1974)REVIEW OF CENOZOIC FRESHWATER FISH FAUNAS IN AFRICA.ANNALS OF THE GEOG.SURVEY OF EGYPT, IV: 211-232.

[329] GREEN,J. (1976)
CORBET,S.A.
WATTS,E.
OEY BIAUW LAN ECOLOGICAL STUDIES ON INDONESIAN LAKES AND RESTRATIFICATION OF RANU LAMONGAN. J.ZOOL.,LOND. 180:315-354.

[330] GROENEWALD,A.A.V.J. (1961)CLARIAS,THE FRESHWATER BARBEL.FAUNA FLORA.NO.12: 47-55.

[331] GROENEWALD,A.A.V.J. (1964)THE ROLE OF EUTROPIUS DEPRESSIROSTRIS PETERS (PISCES:SCHILBEIDAE) AS A PREDITOR IN A LOWVELD IMPOUNDMENT IN THE TRANSVAAL. HYDROBIOLOGIA 23:267 273.

[332] GROENEWALD,A.A.V.J. (1964)OBSERVATIONS ON THE FOOD HABITS OF CLARIAS GARIEPINUS(BURCHELL),THE SOUTH AFRICAN FRESHWATER BARBEL(PISCES,CLARIIDAE)IN THE TRANSVAAL. HYDROBIOLOGIA,23(1/2):287-291.

[333] GROVER,J.H. (1976)
RECOMETA,R.D.
DUREZA,V.A. PRODUCTION AND GROWTH OF MILKFISH, COMMON CARP,AND CATFISH IN FERTILIZED FRESH WATER PONDS.KALIKSAN. 5(2):193-206.

[334] GUIGLIA,D. (1935)PESCI SPEDIZIONE ZOOLOGICA DEL MARCHESE SAVERIO PATRIZI NEL BASSO GIUBA E NELL'OITRE-GIUBA. ANN.MUS.STOR.NAT.GENOVA 58,1935:27-49.

[335] GULLAND,J.A. (1961)
HARDING, D. THE SELECTION OF CLARIAS MOSSAMBICUS (PETERS) BY NYLON GILLNETS.J.CONS.PERM.INT.EXPLOR.MER. 26(2): 215-222.

[336] GUNTERT,H. (1938)BESCHREIBUNG EINIGER UNBESTIMMTER CLARIIDEN MIT DREI NEUEN ARTEN AUS DER NATURHISTORISCHEN MUSEUM IN WIEN. ZOOL.ANZ.LEIPZIG. 124:289-303.

[337] GUNTHER,A. (1864)CATALOGUE OF THE FISHES OF THE BRITISH MUSEUM. VOL.V.BRITISH MUSEUM,LONDON.

[338] GUNTHER,A. (1867)NEW FISHES FROM THE GABOON AND GOLD COAST. ANN. AND MAG.NAT.HIST.XX.: 110-117.

[339] GUNTHER,A. (1896)REPORT ON A COLLECTION OF REPTILES AND FISHES MADE BY MISS.M.H.KINGSLEY,DURING HER TRAVELS ON THE OGOOUE RIVER AND IN OLD CALABAR. ANN.NAT.HIST (6)XVIII: 261-285.

[340] GUNTHER,A. (1896)REPORT ON A COLLECTION OF FISHES MADE BY DR. A. DONALDSON SMITH DURING HIS EXPEDITION TO LAKE RUDOLF. P.Z.S. PP217-224

[341] GUNTHER,A. (1902)LAST ACCOUNT OF FISHES COLLECTED BY MR.R.B.N. WALKER,C.M.Z.S.ON THE GOLD COAST.P.Z.S.II,

PP.330-339.

[342] GUPTA,S.P. (1951)TREMATODE PARASITES OF INDIAN FISHES.THREE NEW TREMATODES OF THE SUB-FAMILY LEPTOPHALLINAE DAYAL,1938 FROM FRESHWATER FISHES OF U.P.INDIA. IND.J.HELM.3:29-40.

[343] GUSSEV,A.V. (1961)[A VIVIPAROUS MONOGENETIC TREMATODE FROM FRESHWATER BASINS OF AFRICA.] DOK .AKAD. NAUK SSSR, 136(1-6):177-179(IN RUSSIAN).

[344] HADDADIN,M.J. ALAWI,A.A. (1974) DEAD POPULATIONS OF FISH IN THE RIVERS JORDAN AND ZARQA. BIOL.CONSERV.6(3):215-216.

[345] HAGENDOORN,H. WIEME,R. (1975) PRELIMINARY RESULTS CONCERNING THE CULTURE OF CLARIAS LAZERA IN CAMEROON. IN: FAO/CIFA SYMPOSIUM ON AQUACULTURE IN AFRICA,GHANA, 30 SEPT. 1975, FAO-CIFA--75/SE-10, PP6.

[346] HAIDER,S. SATHYANESAN,A.G. (1973) AUTORADIOGRAPHIC LOCALISATION OF TRIIODOTHYRONINE SENSITIVE AREA IN THE LATERAL HYPOTHALMIC REGION OF A TELEOST FISH. CURR.SCI . 42(6):204-205.

[347] HAIDER,S. SATHYANESAN,A.G. (1975) ANTAGONISTIC RESPONSE OF CHLORORPROMAZINE AND FORMALIN IN EFFECTING QUANTITATIVE CHANGES IN THE NEUROSECRETORY MATERIAL IN THE TELEOST CLARIAS BATRACHUS (L).ACTA.ANAT. 92(4):630-637.

[348] HAIDER,S., SATHYANESAN,A.G. (1975) ALCOHOL INHIBITION OF FORMALIN INDUCED DEPLETION OF NEURO SECRETORY MATERIAL IN THE TELEOST CLARIAS BATRACHUS.ENDOKRINOLOGIE.65(3):313-319.

[349] HAIDER,S. (1975)INVOLVEMENT OF CHLORPROMAZINE IN COMPENSATING THE FORMALIN INDUCED DEPLETION OF NEURO SECRETORY MATERIAL IN THE TELEOST CLARIAS BATRACHUS.IND.SCI. CONG.ASSOC.PROC.62:174-175.

[350] HAIDER,S. SATHYANESAN,A.G. (1975) ON THE SIMILARITY BETWEEN THE CORTICOTROPHS AND PARS INTERMEDIA CELLS IN A TELEOST FISH CLARIAS BATRACHUS.EXPERIENTIA.31(4):423-424.

[351] HANEK,G. FURTADO,J.I. (1973) GYRODACTYLUS FERNANDOI SP.N.(MONOGENEA: GYRODACTYLINAE)FROM CLARIAS BATRACHUS L. IN MALAYSIA. ZOOL.ANZ.191(3/4):196-198.

[352] HARDING,D. (1960)PRELIMINARY OBSERVATIONS ON THE EFFECTS OF SEASONAL CHANGE IN WATER CONDITIONS ON THE FISHERY IN THE BANGWEULU AREA OF NORTHERN RHODESIA.IN:THIRD SYMPOSIUM ON HYDROBIOLOGY AND INLAND FISHERIES-PROBLEMS OF MAJOR LAKES,1960. CSA,PUB,63:86-96.

[353] HARDING,D. (1964)HYDROLOGY AND FISHERIES IN LAKE KARIBA.VERU. INTERNAT.VEREIN.LIMNOL.XV:139-149.

[354] HART,P.J.B. PITCHER,T.J. (1969) FIELD TRIALS OF FISH MARKING USING A JET INNOCULATOR.J.FISH.BIOL.1:383-385.

[355] HASNAIN,A.U. SIDDIQUI,A.Q. (1974) COMPARATIVE ELECTROPHORETIC STUDIES ON THE TISSUE PROTEINS OF SOME CATFISHES.MAHASAGAR 7(1&2):53-59.

[356] HASTINGS,W.H. (1973)[REGIONAL PROJECT ON RESEARCH AND FISHERIES DEVELOPMENT(CAMEROON-CENTRAL AFRICAN REPUBLIC-GABON-CONGO,PEOPLE,S REP.).EXPERIENCE RELATED TO THE PREPARATION OF FISH FEED AND THEIR FEEDING.REPORT PREPARED FOR THE REGIONAL PROJECT]. FAO/UN, FAO-FI-DP/RAF--66/054/1:1-24. (IN FRENCH).

[357] HASTINGS,W.H. (1975)FORMULA FOR EXTENSIVE CULTURE.IN:VOL 3.FOODS FOR THE EXPANDING WORLD.PROC.NINTH INTERNATIONAL

CONFERENCE OF NUTRITION,MEXICO CITY,SEPT.3-9,1972. PP208-214.

[358] HATTINGH,J. (1972)OBSERVATIONS ON THE BLOOD PHYSIOLOGY OF FIVE SOUTH AFRICAN FRESHWATER FISH. J.FISH BIOL. 4:555-563.

[359] HATTINGH,J. (1973)SOME BLOOD PARAMETERS OF THE YELLOWFISH(BARBUS HOLUBI) AND THE BARBEL(CLARIAS GARIEPINUS)ZOOL. AFR.8(1):35-39.

[360] HATTINGH,J. (1973)A STUDY OF SOME BLOOD CONSTITUENTS OF TWO SPECIES OF MUDFISH .COMP.BIOCHEM.PHYSIOL.

[361] HATTINGH,J. (1974)PLASMA PROTEINS OF BARBUS HOLUBI AND CLARIAS GARIEPINUS (TELEOSTEI).ZOOL.AFR.9(2):185-193.

[362] HATTINGH,J. (1975)HEPARIN AND ETHYLENEDIAMINE TETRA-ACETATE AS ANTICOAGULANTS FOR FISH BLOOD. PFLUGERS ARCH. 355(4):347-352.

[363] HERRE,A.W.C.T. (1924)DISTRIBUTION OF THE TRUE FRESHWATER FISHES IN THE PHILIPPINES-II THE PHILIPPINE LABYRINTHICI, CLARIIDAE AND SILURIDAE. PHILIPPINE J.SCI. MANILA P.I. 24:683-707.

[364] HICKLING,C.F. (1961)TROPICAL INLAND FISHERIES.CAMELOT PRESS,LONDON. PP287.

[365] HILGENDORF F. (1905)[FISH OF GERMAN AND BRITISH EAST AFRICA.] ZOOL. JAHRB.SYST. XXII : 405-420 (IN GERMAN)

[366] HILL,T.K. PARDUE,G.B. SMITH,B.W. (1970) AN EVALUATION OF SEVERAL MARKS ON CHANNEL CATFISH,ICTALURUS PUNCTATUS.PROC.24TH.ANN.CONF. SE.GAME AND FISH COMM.1970.

[367] HIPPLE,E.VON (1946)STOMACH CONTENTS OF CROCODILES.UGANDA J.10:148-149.

[368] HOLEY,D.T. BELSARE,D.K. (1973) MORPHOLOGY OF THE OLFACTORY SAC IN SOME TELEOSTS Z.MIKROSK. ANAT.FORSCH.87(4):463-469.

[369] HOLLY,M. (1927)[MORE FISHES FROM THE CAMEROON.]SITZ.BER.AK.. WISS. WIEN.ABT. 1(136):421-427.(IN GERMAN).

[370] HOLLY,M. (1930)[ON THE NOMENCLATURE OF CLARIAS CAMERONENSIS HOLLY.] ZOOL.ANZ.LEIPZIG 90:112-113 (IN GERMAN).

[371] HOLLY,M. (1930)[ON SYNONYMY OF THE CLARIAS CAMERONENSIS,HOLLY.] ZOOL.ANZ.LEIPZIG.90:112-113. (IN GERMAN)

[372] HOLL,E.A. (1966)SOME NOTES ON THE BREEDING OF BARBEL CLARIAS GARIEPINUS (BURCHELL) IN RHODESIA NEWSL. LIMNOL.SOC. STH. AFR.NO.7:38-41

[373] HOLL,E.A. (1968)NOTES ON THE SPAWNING BEHAVIOUR OF THE BARBEL CLARIAS GARIEPINUS BURCHELL IN RHODESIA. ZOOL AFR 3(2):185-188.

[374] HORA,S.L. (1933)RESPIRATION IN FISHES. J.BOMBAY NAT.HIST.SOC. 36(3):538-560.

[375] HORA,S.L. (1934)RAINS OF FISHES IN INDIA.J.ASIATIC SOC.BENGAL. N.S. 59(1):95-110.

[376] HORA,S.L. (1934)TRADE IN LIVE FISH (JIOL MACHH) IN CALCUTTA. J.AS.SOC.BENGAL (NS),30:1-15.

[377] HORA,S.L. (1935)PHYSIOLOGY,BIONOMICS AND EVOLUTION OF THE AIR BREATHING FISHES OF INDIA.TRANS. NAT.INST.SCI. INDIA 1(1-16

[378] HORA,S.L. (1936)SILURIOD FISHES OF INDIA,BURMA,AND CEYLON.IV. FISHES OF THE GENUS CLARIAS.REC.IND.MUS.38:347-351.

[379] HORA,S.L. (1937)GEOGRAPHICAL DISTRIBUTION OF INDIAN FRESHWATER FISHES AND ITS BEARING ON THE PROBABLE CONNECTION BETWEEN INDIA AND THE ADJACENT COUNTRIES.CURR. SCI.5(7):351-356.

[380] HORA,S.L. (1941)SILUROID FISHES OF INDIA, BURMA AND CEYLON XII. A FURTHER NOTE ON FISHES OF THE GENUS CLARIAS GRONOVIUS.REC.INDIAN MUS. CALCUTTA. 43:97-115.

[381] HORA,S.L. (1941)

GUPTA,J.C.
NOTES ON THE MALAYAN FISHES IN THE COLLECTION OF THE RAFFLES MUSEUM SINGAPORE. BULL.RAFFLES MUS. SINGAPORE. 17:12-43.

[382] HORA,S.L.
(1941)SILURIOD FISHES OF INDIA,BURMA,AND CEYLON.XII. A FURTHER NOTE ON FISHES OF THE GENUS CLARIAS. REC.IND.MUS.42:112-115.

[383] HORA,S.L.
(1949)SYSTEMATIC POSITION OF SILURIDS IN HAMILTONS GANGETIC FISHES .REC.INDIAN MUS.CALCUTTA 46 : 63-72.

[384] HORA,S.L.
(1951)FISH CULTURE IN RICE FIELDS.CURR.SCI.(BANGALORE INDIA).(20)7:171-173

[385] HORA,S.L.
MENON,A.G.K.
(1952)
DISTRIBUTION OF CLARIID FISHES IN SPACE AND TIME. PROC.INDIAN SCI.CONGR. 38(4):17-18 (ABSTRACT)

[386] HORA,S.L.
(1955)HANDBOOK OF FISHCULTURE OF THE INDOPACIFIC REGION FAO,ROME.PP264.

[387] HORA,S.L.
PILLAY,T.V.R.
(1962)
HANDBOOK ON FISH CULTURE IN THE INDO-PACIFIC REGION.FAO FISHERIES BIOLOGY TECHNICAL PAPER NO. 14:204.

[388] HUBRECHT,A.A.W.
(1880)ON A COLLECTION OF FISHES FROM THE ST PAULS RIVER LIBERIA, WITH DESCRIPTIONS OF THREE NEW SPECIES. NOTES LEYD.MUS.III.PP.66-71

[389] HUET,M.
(1974)TEXTBOOK OF FISH CULTURE :BREEDING AND CULTIVATION OF FISH.FISHING NEWS (BOOKS)LTD. LONDON.PP436

[390] HUGHES,G.M.
MUNSHI,J.S.D.
(1973)
NATURE OF AIR BREATHING ORGANS OF THE INDIAN FISHES CHANNA,AMPHINOUS,CLARIAS AND SACCOBRANCHUS AS SHOWN BY ELECTRON MICROSCOPY. J.ZOOL.LONDON,170(2):245-270.

[391] HUISMAN,E.A.
(1976)ASPECTS OF FISH CULTURE AND FISH BREEDING.MISC. PAP.LANDBOUWHOGESCH WAGENINGEN.13:1-69.

[392] IBRAHIM,K.H.
(1975)PROGRESS AND PRESENT STATE OF AQUACULTURE IN TANZANIA. IN: FAO/CIFA SYMPOSIUM ON AQUACULTURE IN AFRICA GHANA,30 SEPT.1975, FAO-CIFA--75/SC-2. PP14.

[393] IDRASENA,H.H.A.
(1970)LIMNOLOGICAL AND FRESHWATER FISHERIES DEVELOPMENT WORK IN CEYLON. PROC. IBP/UNESCO REG.MEET.INLAND WAT.BIOL.SOUTHEAST ASIA,KUALA LUMPUR + MALACCA,5TH-11TH MAY,1969, PP45-47.

[394] IDYLL,C.
(1969)NEW FLORIDA RESIDENT THE WALKING CATFISH. NAT. GEOGRAPHIC,135:847-851.

[395] IDYLL,C.
(1970)[A NEW RESIDENT IN FLORIDA] ZIVA,PRAGUE 18:231-232 (IN CZECH)

[396] IMAN,A.
(1970)REARING OF CATFISH IN ARTIFICIAL PONDS AND BIOLOGICAL CHANGES INDUCED BY LOW DOSE TOTAL BODY GAMMA IRRADIATION. M SC THESIS, FACULTY OF SCIENCE,CAIRO UNIVERSITY.

[397] IMAN,A.E.
ROUSHDY,H.M.
PHILISTEEN,A.
(1970)
FEEDING OF CATFISH CLARIAS LAZERA IN EXPERIMENTAL PONDS. BULL.INST.OCEANOGR.FISH. UAR ,(1):205-222

[398] INGER,R.F.
(1952)WALKING FISHES OF SOUTHEASTERN ASIA TRAVEL ON LAND. CHICAGO NAT.HIST.MUS.BULL.2319:4-5.

[399] INGER,R.F.
KONG,C.P.
(1962)
THE FRESHWATER FISHES OF NORTH BORNEO. FIELDIANA (ZOOL.) 45: 1-268.

[400] INGLIS,W.G.
OGDEN,C.G.
(1965)
CHANAUDUS CHABAUDI GEN. ET SP. NOV. FROM A FRESHWATER FISH IN SIERRA LEONE (NEMATODA, SEURATOIDEA.) REVUE ZOOL. BOT. AFR,71(1/2):171 -176

[401] INOUE,K.
SWEGAN,S.
(1970)
ECONOMIC SURVEY ON CATFISH CULTURE IN SUPHANBURI PROVINCE,THAILAND.THAI.FISH.GAZ. 23(2):251-269.

[402] INTENGAN,C.L.
ALEJO,L.G.
CONCEPCION,I.
CORPUS,V.A.
SALID,R.D.
DEL ROSARIO,I.
GOMES,R.
HENSON,J. (1956)
COMPOSITION OF PHILIPPINE FOODS V.PHILIPP.J.SCI. 85:203-213.

[403] IRVINE,F.R. (1947)THE FISH AND FISHERIES OF THE GOLDCOAST. CROWN AGENTS. LONDON. PP325.

[404] ISHAK,M.M.,
KHALIL,S.R.,
ABDELMALIK,W.E.Y. (1977)
DISTRIBUTION AND TISSUE RETENTION OF CESIUM-134 AND COBALT-60 IN THE NILE CATFISH CATFISH CLARIAS LAZERA.HYDROBIOLOGIA.54(1):41-48.

[405] JACKSON,P.B.N. (1955)A NEW FISH OF THE GENUS CLARIAS GRONOV FROM LAKE NYASA, WITH NOTES ON THE DISTRIBUTION OF THE CLARIIDAE AND OTHER CATFISHES IN THE LAKE. PROC.ZOOL.SOC.LONDON. 125:681-684.

[406] JACKSON,P.B.N. (1959)A REVISION OF THE CLARIID CATFISHES OF NYSALAND WITH DESCRIPTIONS OF A NEW GENUS AND SEVEN NEW SPECIES.PROC.ZOOL.SOC.LOND.132:109-128.

[407] JACKSON,P.B.N. (1960)ICHTHYOLOGY-FISH OF THE MIDDLE ZAMBEZI.KARIBA STUDIES.MANCHESTER UNIVERSITY PRESS.

[408] JACKSON,P.B.N. (1961)THE FISHES OF NORTHERN RHODESIA.GOVT.PRINTER, LUSAKA:140.

[409] JACKSON,P.B.N. (1961)CHECKLIST OF THE FISHES OF NYASALAND.OCC.PAPERS. NAT.MUS.RHODESIA.3:552-557.

[410] JACKSON,P.B.N. (1962)ECOLOGICAL FACTORS AFFECTING THE DISTRIBUTION OF FRESHWATER FISHES IN TROPICAL SOUTHERN AFRICA.ANN.CAPE PROV.MUS.11:223-228.

[411] JACKSON,P.B.N. (1970)THE AFRICAN GREAT LAKE FISHERIES:PAST,PRESENT,AND FUTURE.AFR.J.TROP.HYDROBIOL.FISH.1(1): 35-49.

[412] JAFRI,A.K.
SHAFFI,S.A. (1975)
EFFECT OF ASPHYXIATION ON THE SERUM AND KIDNEY ALKALINE PHOSPHATASE ACTIVITY IN CLARIAS BATRACHUS(LINN).BROTERIA.44(1-2):59-62.

[413] JAISWAL,A.G.
BELSARE,DK (1974)
REGENERATION OF CAUDAL NEUROSECRETORY SYSTEM IN THE CATFISH, CLARIAS BATRACHUS L. Z.MIKROSK.ANAT. FORSCH.,LEIPZIG. 88(6):987-996 .

[414] JAISWAL,A.G.,
BELSARE,D.K. (1975)
EFFECT OF ELECTRICAL STIMULATION OF THE OLFACTORY TRACT AND THE CAUDAL SPINAL CORD ON THE DAHLGREN CELLS IN CLARIAS BATRACHUS.ZEITSCHRIFT FUER MIKROS ANAT.FORSC.89(3):410-414.

[415] JAISWAL,A.G.
BELSARE,D.K. (1975)
ON THE MALFORMED CAUDAL NEUROSECRETORY SYSTEM IN CLARIAS BATRACHUS L. CURR.SCI.44(4):129-130.

[416] JANICKI,C. (1926)CESTODES S.STR. AUS FISCHEN UND AMPHIBIEN. RESULTS OF THE SWEDISH ZOOLOGICAL EXPEDITION TO EGYPT AND THE WHITE NILE, 1901,5(27B):1-58.

[417] JOCQUE,R. (1977)A STUDY OF CLARIAS SENEGALENSIS IN THE REGION OF LAKE KOSSOU IVORY COAST.HYDROBIOLOGIA.54(1):49-66.

[418] JOHNELS,A.G. (1957)THE MODE OF TERRESTRIAL LOCOMOTION IN CLARIAS. OIKOS, 8(2): 122-129

[419] JOHNSEN,S. (1926)ON A COLLECTION OF FISHES FROM THE WHITE NILE AND THE YEI RIVER (WITH NOTES ON THE SKULL OF ALABENCHELYS AND CLARIAS). BERGENS MUS.AARB 10:22

[420] JORDAN,D.S. (1900)FISHES. PP403-404.

[421] JORDAN,D.S.
EVERMANN,B.W. (1902)
NOTES ON A COLLECTION OF FISHES FROM THE ISLAND FORMOSA.P.U.S.MUS.XXV: 315-368.

[422] JORDAN,K. (1936)DR KARL JORDAN'S EXPEDITION TO SOUTH WEST AFRICA AND ANGOLA. NARRATIVE. NOVIT.ZOOL.TRING 40:17-62

[423] JORDAN,J. (1976)THE INFLUENCE OF BODY WEIGHT ON GAS EXCHANGE IN THE AIR BREATHING FISH, CLARIAS BATRACHUS. COMP. BIOCHEM.PHYSIOL.(A),53(3):305-310.

[424] JOY,K.P. SATHYANESAN,A.G. (1976) DISTRIBUTION OF MONO AMINE OXIDASE IN RELATION TO THE HYPOTHALMO HYPOPHYSEAL SYSTEM OF THE TELEOST CLARIAS BATRACHUS.J.NEURAL TRANSMISSION. 39(1-2): 47-62.

[425] JUBB,R.A. (1952)NOTES ON FRESHWATER FISHES OF SOUTHERN RHODESIA: FAMILY CLARIIDAE.RHOD.AGRIC.J.49(1):3-5.

[426] JUBB,R.A. (1953)FISH LADDERS AND THE MIGRATION OF FRESHWATER FISHES.RHOD.AGRIC.J.50(2):68-69.

[427] JUBB,R.A. 1950,AND TO BAROTSELAND,1952.OCC.PAP.NAT.MUS.S. RHOD.3(22B):177-189.
(1958)A CAVE DWELLING CAT-FISH CLARIAS CAVERNICOLA, TREWAVAS 1936, IN SOUTH WEST AFRICA.PISCATOR 12 (43):56-57.

[428] JUBB,R.A. (1958)A PRELIMINARY REPORT ON THE COLLECTIONS OF FRESHWATER FISH MADE BY THE BERNARD CARP EXPEDITIONS TO THE CAPRIVI STRIP,1949,LOWER SABI

[429] JUBB,R.A. (1960)FRESHWATER FISHES OF THE CAPE PROVINCE.CAPE. PROV.MUS.4:1-72.

[430] JUBB,R.A. (1961)AN ILLUSTRATED GUIDE TO THE FRESHWATER FISHES OF THE ZAMBEZI RIVER,LAKE KARIBA,PUNGWE,SABIE,LUNDI AND LIMPOPO RIVERS.BULAWAYO,STUART MANNING.

[431] JUBB,R.A. (1964)A NEW SPECIES OF CLARIALLABES (PISCES,CLARIIDAE) FROM THE UPPER ZAMBEZI RIVER.ANN.MAG.NAT.HIST. (13) 7: 393-395.

[432] JUBB,R.A. (1967)THE FRESHWATER FISHES OF SOUTHERN AFRICA.BALKEMA, CAPETOWN. PP .

[433] JUBB,R.A. GAIGHER,I.G. (1971) CHECKLIST OF THE FISHES OF BOTSWANA. ARNOLDIA (RHODESIA) 5(7):1-22.

[434] JULIANO,R.O. (1968)THE GROSS MORPHOLOGY OF THE GAS BLADDERS OF SOME LAGUNA BAY FISHES. PHILIPPINE J.FISH 8(1):1 -21.

[435] JUNOR,F.J.R. (1968)BRIEF OBSERVATIONS ON THE WEIGHT OF FISH THAT CAN BE CARRIED BY THE FISH EAGLE,HALIAETUS VOCIFER.PISCATOR,72:22-23.

[436] KALK,M. (1969)A REPORT ON THE LAKE CHILWA LIMNOLOGICAL PROGRAMME IN MALAWI. IN: IBP REPORT OF THE REGIONAL MEETING OF HYDROBIOLOGISTS IN TROPICAL AFRICA. KAMPALA,UGANDA,20-28 MAY,1968:PP88-91.

[437] KAK,A.B. GHOSH,A. (1950) RESPONSES OF THE GENITAL SYSTEM AND THE URINOGENITAL PAPILLA OF THE FEMALE CATFISH, HETEROPNEUSTES FOSSILS (BLOCH) TO SEX HORMONES. ROUX.ARCH.ENTW.MECH.ORGAN.144 PP 257-264.

[438] KEILHACK,L. (1910)[ABOUT A FEW FISH COLLECTED BY DR H. MONKE IN DUALA (CAMEROON).]BERLIN MITT.ZOOL.MUS. 5:117-124.(IN GERMAN)

[439] KEIRMEIR,A. (1939)ON THE BLOOD SUGAR OF FRESHWATER FISH.Z.VERGL. PHYSIOL.27(460-491.

[440] KHALIL,L.F. (1961)ON A NEW TREMATODE, ORIENTOCREADIUM LAZERI N.SP. FROM A FRESHWATER FISH, CLARIAS LAZERA, IN THE SUDAN.J.HELMINTH.35:259-62.

[441] KHALIL,L.F. (1963)ON DIPLOSTOMUM TREGENNA, THE DIPLOSTOMUM STAGE OF DIPLOSTOMUM TREGENNA NAZMI GOHAR, 1932, WITH AN EXPERIMENTAL DEMONSTRATION OF PART OF THE LIFE CYCLE. J.HELMINTH, 37(3):199-206.

[442] KHALIL,L.F. (1963)ON SOME PROTEOCEPHALID CESTODES FROM FRESH WATER FISHES IN THE SUDAN.J.HELMINTH, 37(4): 307-318.

[443] KHALIL,L.F. (1969)STUDIES ON THE HELMINTH PARASITES OF

FRESHWATER FISHES OF THE SUDAN, J.ZOOL. LOND. 158:143-170.

[444] KHALIL,L.F. (1971)CHECKLIST OF THE HELMINTH PARASITES OF AFRICAN FRESHWATER FISHES.TECH.COMM.NO.42.COMM.AGRIC. BUR.UK.PP80.

[445] KHALIL,L.F. (1972)AFROMACRODEROIDES LAZERA GEN.ET.SP.NOV. (ALLOCREADIIDAE(WALLINIINAE),A NEW DIGENEAN FROM THE AFRICAN FRESHWATER FISH CLARIAS LAZERA.J. HELMIN.XLV1(4):314-344.

[446] KHANNA,S.S. GOSWAMI,O.B. (1965) STUDIES ON THE TELEOSTEAN HEART. AGRA.UNIV.J.RES. 14(1):105-113.

[447] KHANNA,S.S. (1966)HISTOLOGY OF THE DIGESTIVE TRACT OF A TELEOST, CLARIAS BATRACHUS(LINN).J.ZOOL.SOC.INDIA. 16:53-58

[448] KHANNA,S.S. SINGH,H.R. (1967) ON THE MORPHOLOGICAL PECULIARITIES OF THE VALVULA CEREBELLI IN SOME TELEOSTS. ZOOL. ANZ .178(5/6):413-419.

[449] KHANNA,S.S. MEHROTRA,B.K. (1968) HISTOLOGY OF THE ISLETS OF LANGERHANS IN NORMAL AND ALLOXAN TREATED FRESHWATER INDIAN TELEOST, CLARIAS BATRACHUS.ZOOL.BEITR.14:487-497.

[450] KHANNA,S.S. MEHROTRA,B.K. (1969) EFFECT OF INSULIN AND GLUCOSE ON THE BETA CELLS OF THE PANCREATIC ISLETS IN FRESHWATER TELEOST, CLARIAS BATRACHUS.ACTA ZOOL.,STOKH.50:91-95.

[451] KHANNA,S.S. BHATT,S.D. (1974) HISTOPATHOLOGY OF THE ENDOCRINE PANCREAS OF A FRESHWATER FISH,CLARIAS BATRACHUS(L.)VI EFFECT OF COBALT CHLORIDE ADMINISTRATION. VERH. ANAT. GES.JENA ANAT.ANZ.RD.136(1/2):157-164.

[452] KHANNA,S.S. BHATT,S.D. (1975) HISTOPATHOLOGY OF THE ENDOCRINE PANCREAS OF A FRESH-WATER FISH, CLARIAS BATRACHUS (L).5. EFFECT OF A EPINEPHRINE ADMINISTRATION. ACTA.ANAT.,93(3): 351-360.

[453] KHANNA,S.S., BHATT,S.D. (1975) HISTO PHYSIOLOGY OF THE ENDOCRINE PANCREAS OF A FRESH WATER FISH CLARIAS BATRACHUS PART 7:THE EFFECT OF PANCREATECTOMY AND INSULIN.J.AN.MORP.PHY 22(1):27-37.

[454] KHAN,S.H. QAYYAIM,A. (1969) DIFFERENTIAL BLOOD CELL COUNTS OF FOUR SPECIES OF FRESHWATER AIR-BREATHING FISHES. PROC.INDIAN ACAD. SCI. 69B:29-35.

[455] KHAN,H.A. (1972)INDUCED BREEDING OF AIR-BREATHING FISHES.IND. FARM.22(4):44-45.

[456] KHAN,S.H. SIDDIQUI,Q.A. HASNAIN,A. (1972) THE HYPOTONIC FRAGILITY OF SOME FISH ERYTHROCYTES. PROC.INDIAN ACAD.SCI.75(B)265-270.

[457] KHAN,H.A., MUKHOPADHYAY,S.K. (1975) PRODUCTION OF STOCKING MATERIAL OF SOME AIR BREATHING FISHES BY HYPOPHYSATION.J.INL.FISH.SOC. INDIA.7:156-161.

[458] KHOSA,D. CHANDRASEKHAR,K. (1972) EFFECT OF COPPER ACETATE AND ASPHALT ON GONADAL ACTIVITIES AND THE CORRELATED CHANGES IN THE PRE-OPTIC NUCLEUS OF TWO GENERA OF TELEOSTEAN FISHES, CLARIAS BATRACHUS(LINN.) AND OPHIOCEPHALUS PUNCTA TUS(BLOCH).PROC.OF IND.ACAD.SCI.7:B)NO6:229-239

[459] KLOKE,C.W. POTAROS,M. (1975) THE TECHNOLOGY AND ECONOMICS OF CATFISH (CLARIAS SPP.) FARMING IN THAILAND. FAO.IPFC. OCC.PAP. 1975/2.

[460] KNER,R. (1867)[THE CRUISE OF THE AUSTRIAN FRIGATE 'NOVARA' AROUND THE WORLD IN THE YEARS 1857,1858,1859.] ZOOLOGISCHER THEIL. FISCHE. 3.ABTHEILUNG.WEIN, :275-433.

[461] KNUDSEN,E.I. (1977)DISTINCT AUDITORY AND LATERAL LINE NUCLEI IN THE MIDBRAIN OF CATFISHES.J.COMP.NEUROLOGY.173(3):417-431.

[462] KONIKOFF,M. (1975)NIGERIA. FEASIBILITY OF CAGE CULTURE AND OTHER AQUACULTURE SCHEMES AT KAINJI LAKE. FAO. KAINJI LAKE RES . PROJECT, FAO FI:NIR- 66/524/18.PP17.

[463] KOSSWIG,C. (1969)NEW CONTRIBUTIONS TO THE ZOOGEOGRAPHY OF FRESH WATER FISH OF ASIA MINOR, BASED ON COLLECTIONS MADE BETWEEN 1964-1967.ISRAEL J.ZOOL.18:249-254.

[464] KOUMANS,F.P. (1949)ON SOME FOSSIL FISH REMAINS FROM JAVA. ZOOL. MEDED.LEIDEN 30(5):77-82.

[465] KRISHNAMOORTHI,K.P. ABDULAPPA,M.K. JAGANNATH,RAO,A.V. (1976)

PRODUCTIVITY OF CLARIAS BATRACHUS IN SEWAGE FERTILIZED FISH PONDS.IND.J.ENVIR.HEALTH 18(4): 292-298.

[466] KRUGER,E.J. (1971)[AN INVESTIGATION INTO THE ECOLOGY AND FEEDING HABITS OF CERTAIN FISH SPECIES IN THE LOSKOPDAM]. D.SC. THESIS POTCHEFSTROON UNIVERSITY FOR CHRISTIAN HIGHER EDUCATION. (IN AFRICAANS).

[467] KULKARN,R.B., BELSARE,D.K. (1975)

HISTO MORPHOLOGICAL OBSERVATIONS ON THE LIVER IN SOME FRESH WATER TELEOSTS.ZEITSCHRIFT FUER MIKROS. ANAT.FORSC.89(5):922-933.

[468] KUMAR,A. (1954)A CRANIAL MORPHOLOGY OF THE SKULL OF CLARIAS BATRACHUS. CURR.SCI. 23:406-407.

[469] KY,H. (1968)[NEW SPECIES OF MONOGENEA FROM FRESHWATER FISHES OF NORTH VIETNAM.PART1 :PARAZITOLOGIYA,2(4):297 -301. (IN RUSSIAN)

[470] KY,H. (1971)[SOME SPECIES OF NEMATODA FROM FRESH WATER FISHES OF NORTH VIETNAM.] PARAZITOLOGIYA,5(3):241-250. (IN RUSSIAN)

[471] LACHNER,E.A. ROBINS,C.R. COURTENAY,JUN,W.R. (1970)

EXOTIC FISHES AND OTHER AQUATIC ORGANISMS INTRODUCED INTO NORTH AMERICA. SMITHSON.CONTRIB. ZOOL.59:1-29.

[472] LADIGES,W. ET AL (1958)

[THE TYPES AND KINDS OF FISH COLLECTIONS OF THE HAMBURG ZOOLOGICAL INSTITUTE AND ZOOLOGICAL MUSEUMS.] MITT HAMBURG ZOOL.MUS. 56:155-167.

[473] LAGLER,K.F. KAPETSKY,J.M. STEWART,D.J. (1971)

THE FISHERIES OF THE KAFUE RIVER FLATS, ZAMBIA,IN RELATION TO THE KAFUE GORGE DAM. FINAL REPORT. CENTRAL FISHERIES RESEARCH INSTITUTE, CHILANGA, ZAMBIA. REPORT PREPARED FOR THE FOOD AND AGRICULTURE ORGANIZATION OF THE UNITED NATIONS (ACTING AS EXECUTIVE AGENCY FOR THE UNITED NATIONS DEVELOPMENT PROGRAMME). ORA PROJECT 365020, FI:SF/ZAM11 TECH.REP.1.PP161.

[474] LAHIRI,S.B., KAPOOR,B.G. (1975)

A REPORT ON THE EXISTENCE OF SMALL PIT ORGANS IN THE CATFISHES CLARIAS BATRACHUS AND HETEROPNEUSTES FOSSILIS.GEOBIOS(JODHPUR).2(2/3):86-87.

[475] LAL,C. (1965)TWO NEW SPECIES OF GENUS PROCAMALLANUS BAYLIS,1923 FROM NORTH INDIA.LABDEV 3:199 200

[476] LAL,C. (1966)TWO NEW NEMATODES OF THE GENUS SPINITECTUS FOURMENT,1883,FROM CATFISH OF NORTH

INDIA. LABDEV 4:121-123

[477] LAMBERT,J.G. (1958)[SILURID AND CYPRINOID FISHES COLLECTED IN FRENCH GUINEA, WITH A DESCRIPTION OF A NEW SPECIES OF MICROSYNODONTIS.] REV.ZOOL.BOT.AFR. 57:39-56 (IN FRENCH)

[478] LAUZANNE,L. (1975)THE SELECTION OF PREY IN 3 MALACOPHAGUS FISHES OF LAKE CHAD,CHAD.CAHIERS ORSTOM,SER.HYDROBIOL.9(1): 3-7.

[479] LAZARD,J. (1973)[ASSESSMENT OF THE FISHERIES IN THE NOMBOLO RESERVOIR].NOTES DOC.PECHE.PISCIC.(NOUV.SER.). 6:52-62. (IN FRENCH)

[480] LEHRI,G.K. (1964)AN INSTANCE OF HERMAPHRODITISM IN CATFISH, CLARIAS BATRACHUS (LINN).CURR.SCI.33:529-530.

[481] LEHRI,G.K. (1966)STUDIES ON THE PITUITARY GLAND IN RELATION TO THE GONADAL CYCLE IN CLARIAS BATRACHUS.(LINN.). NATURWISSENSCHE 53(1):20-21.

[482] LEHRI,G.K. (1966)ENDOCRINE ACTIVITY OF GONADECTOMIZED FISH,CLARIAS BATRACHUS(LINN).NATURWISSENSCHAFTEN,53(15):390.

[483] LEHRI,G.K. (1966)THE PITUITARY GLAND OF CATFISH,CLARIAS BATRACHUS. COPEIA,(4):810-818.

[484] LEHRI,G.K. (1967)THE ANNUAL CYCLE IN THE TESTIS OF THE CATFISH CLARIAS BATRACHUS L. ACTA ANAT.67(1):135-154.

[485] LEHRI,G.K. (1968)CYCLICAL CHANGES IN THE OVARY OF THE CATFISH C. BATRACHUS L. ACTA ANAT. ,69(1):105-124.

[486] LEHRI,G.K. (1970)CORRELATIVE CYCLICAL CHANGES IN THE PITUITARY GLAND AND THE TESTIS OF A CATFISH CLARIAS BATRACHUS L. MIKROSKOPIE 26:50-56.

[487] LELEK,A. (1968)THE VERTICAL DISTRIBUTION OF FISHES IN THE EBO STREAM AND NOTES TO THE FISH OCCURRENCE IN LAKE BOSUMTWI,ASHANTI,GHANA.ZOOL.LISTY.17(3):245-252

[488] LERDPRASERT,N. (1964)THE CULTURE OF CLARIAS FRY IN FLOATING BASKET. THAI FISH.GAZ.13(4).

[489] LERICHE,M. (1901)[CONTRIBUTIONS TO THE STUDY OF FOSSIL SILURIDS]. ANN.S.CGEOL.NORD XXX:165-175.(IN FRENCH)

[490] LE ROUX,P.
STEYN, L. (1968)
FISHES OF THE TRANSVAAL.S.A.BREWERIES INST. JOHANNESBURG,S.A.

[491] LIBOSVARSKY,J.
LUSK,S.
EL-SEDFY,H.M. (1972)
FISHERY SURVEY CARRIED OUT AT LAKE BORULLUS,A.R. EGYPT,IN THE SPRING OF 1971.ACTA.SCI.NAT.ACAD. SCI.BOHEMOSLOV.6(7):3-44.

[492] LING,S.W. (1967)FEEDS AND FEEDING OF WARM-WATER FISHES IN PONDS IN ASIA AND THE FAR EAST.FAO FISH.REP.NO.44, (3):291-309.
VOL.3:291-309.

[493] LISSMANN,H.W.
MACHIN K.E. (1963)
ELECTRIC RECEPTIONS IN A NON-ELECTRIC FISH (CLARIAS). NATURE,LOND.199(4888):88-89.

[494] LOCKLEY,G.J. (1949)THE FAMILIES OF FRESHWATER FISHES OF TANGANYIKA TERRITORY.E.A.AGRIC.J.X1V:

[495] LOCKLEY,G.J. (1960)THE FISHING INDUSTRY OF TANGANYIKA.IN:REPORT ON THE TRAINING CENTRE ON FISHERY SURVEYS FOR THE COUNTRIES OF THE AFRICAN REGION AT TANGA, TANGANYIKA VOLII ESSAYS ON THE STATUS OF FISHERIES DEVELOPMENT IN SELECTED COUNTRIES AND TERRITORIES OF AFRICA. FAO/UN.PP 184-190.

[496] LOISELLE,P.V. (1971)THE FISHES OF THE HEADWATERS OF THE ZIO RIVER. AQUARIUM,N.J.4(8):32-51.

[497] LOISELLE,P.V. (1971)THE FISHES OF THE ZIO MARSHES.AQUARIUM,N.J.4(9):32 -55.

[498] LOMBARD
VAN DER WAAL,B.C.W. (1972)
FISH CULTURE EXPERIMENTS AT MARBLE HALL.STH.AFR. IBP/PF WORKING GROUP ANN REP.NO3:47-49.

[499] LONNBERG,E. (1895)NOTES ON FISHES COLLECTED IN THE CAMEROONS BY MR.Y.SJOSTEDT. OFV.AK.FORM. PP.179-195.

[500] LONNBERG,E. RENDAHL,H. (1920) ON SOME FRESHWATER FISHES FROM LOWER CONGO. ANN. MAG.NAT.HIST.,LONDON 9(6):167-176.

[501] LONNBERG,E. RENDAHL,H, (1922) SOME NEW SILURIDS FROM THE CONGO. ANN.MAG.NAT. HIST. 9(10):122-127. THE NIGER RIVER.) BUL.MUSEUM PARIS:211-213 (IN FRENCH).

[502] LORTEL,L. (1882)[FISHES AND REPTILES OF LAKE TIBERIAS AND SEVERAL OTHER PARTS OF SYRIA.]ARCH.MUS.LYON.III. FISHES:99-182.(IN FRENCH)

[503] LOVE,R.M. (1970)THE CHEMICAL BIOLOGY OF FISHES,ACADEMIC PRESS. LONDON.PP 600 .

[504] LOWE-MCCONNELL,R.H. (1970)ECOLOGICAL STUDIES ON TROPICAL FRESHWATER FOOD FISHES. PROC.IBP/UNESCO REG.MEET. INLAND WAT. BIOL.SOUTHEAST ASIA,KUALA LUMPUR + MALACCA, 5TH-11TH MAY,1969, PP91-103

[505] LUYEYI,A. (1971)[COLD BLOODED VERTEBRATES IN THE COLLECTION OF THE O.N.R.D.] HOMME NATURE, KINSHASA 1:129-140. (IN FRENCH).

[506] LYDEKKER,R. (1885)INDIAN TERTIARY AND POST TERTIARY VERTEBRATA. TERTIARY FISHES.PAL.IND.(10)III:241-258.

[507] MACKIEWICZ,J.S. BEVERLEY-BURTON,M. (1967) MONOBOTHRIODES WOODLANDI SP.NOV.(CESTOIDAE, CARYOPHYLLIDAE) FROM CLARIAS MELLANDI BOULENGER (CYPRINIFORMES(CLARIIDAE) IN ZAMBIA.AFRICA.PROC HELMINTH.SOC.WASH, 34(2):125-128.

[508] MACKIEWICZ,J.S. MURHAR,B.M. (1972) REDESCRIPTION OF BOVIENA SERIALIS(BOVIEN,1926) CESTOIDEA:CARYOPHYLLIDEA)FROM THE CATFISH CLARIAS BATRACHUS(L.) IN INDIA. J.HELMIN.XLVI(4) :399-405.

[509] MACLAREN,P.I.R. (1957)THE FISHERIES OF THE KAFUE FLOOD PLAIN.IN:SECOND SYMPOSIUM ON AFRICAN HYDROBIOLOGY AND INLAND FISHERIES,1956.CSA.PUB.NO.25:155.

[510] MAGLIO,V.J. (1971)VERTEBRATE FAUNAS FROM KUBI ALGI KOOBI FORA,AND ILERET AREAS, EAST RUDOLF, KENYA. NATURE,LONDON 231:248-249.

[511] MAJUMDAR,I. (1975)CYTOCHEMICAL STUDIES ON THE PERI NUCLEAR REGION IN THE GROWING OOCYTES OF A TELEOST FISH CLARIAS BATRACHUS.IND.SCI.CONG.ASSOC.PROC.62:194.

[512] MALAVIYA,R.B. (1973)THE YOLK-NUCLEUS IN THE OOCYTES OF CLARIAS BATRACHUS (LINN.).ZOOL.POL.22(3):189-196.

[513] MALETOUNGON,Z. (1975)[BEGINNING COMMERCIAL FISH CULTURE IN CENTRAL AFRICA.] IN:FAO/CIFA SYMPOSIUM ON AQUACULTURE IN AFRICA. CIFA/75/SE 18.PP10. (IN FRENCH).

[514] MANACOP,P.R. (1964)A BRAND NEW INDUSTRY: RAISING FISH IN RICE PADDIES. COFF. CACAO J. 7(7):149:160-161. REPORT).EAFFRO OCC.PAP.6.PP55.

[515] MANN,M.J. (1964)THE FISHERIES OF LAKE RUKWA,TANGANYIKA.(MIMEO

[516] MANN,M.J. (1964)REPORT ON THE FISHERIES SURVEY OF LAKE RUKWA TANGANYIKA.REP.EAFFRO.(1962/63):42-52.

[517] MANN,M.J. (1964)REPORT ON THE FISHERIES SURVEY OF LAKE RUDOLF. KENYA.REP.EAFFRO.(1962/63):53-62.

[518] MANN,M.J. (1965)REPORT ON A FISHERIES SURVEY OF LAKE KITANGIRI. TANZANIA.REP.EAFFRO(1964):43-49.

[519] MANN,M.J. (1967)THE FISHERIES OF LAKE RUKWA,TANGANYIKA.EAFFRO OCC.PAP.(6)(2ND ED.).

[520] MANN,M.J. (1969)A BRIEF REPORT ON A SURVEY OF THE FISH AND FISH-ERIES OF THE TANA RIVER, WITH SPECIAL REFERENCE TO THE PROBABLE EFFECTS OF THE PROPOSED BARRAGES EAFFRO OCC.PAP. 10.PP13.

[521] MANN,M.J. (1971)SOME TAXONOMICAL NOTES ON THE FISH FAUNA OF THE

BARINGO AREA.AFR.J.TROP.HYDROBIOL.FISH.1(1):25-34.

[522] MANTER,H.W. PRITCHARD, M.H. (1964) [ZOOLOGICAL MEDICAL MISSION OF THE MANIEMA WORMS: TREMATODA].ANNLS MUS.R.AFR.CENT.SER.8VO, (CONGO, LEOPOLDVILLE):P.L.G.BENOIT,1959.5. NO.132:75-101. (IN FRENCH)

[523] MAN,H.S.U., HODGKISS,I.J. (1977) STUDIES ON THE ICHTHYO-FAUNA IN PLOVER COVE RESERVOIR HONG KONG.PART I:SEQUENCE OF FISH POPULATION CHANGES.J.FISH.BIOL.10(5):493-503.

[524] MARLIER,G. (1938)[CONSIDERATIONS OF THE ACCESSORY ORGANS USED FOR AIR-BREATHING IN SOME TELEOSTS].ANN.SOC.ROY.ZOOL. BELG.69:163-185.(IN FRENCH)

[525] MATTHES,H. (1960)[A NOTE ON THE REPRODUCTION OF FISH OF LAKE TANGANYIKA].IN:THIRD SYMPOSIUM ON HYDROBIOLOGY AND INLAND FISHERIES-PROBLEMS OF MAJOR LAKES,1960 CSA,PUB.63:107-112.(IN FRENCH)

[526] MATTHES,H. (1964)[THE FISH OF LAKE TUMBA AND OF THE REGION OF THE IKELA. A SYSTEMATIC AND ECOLOGICAL STUDY.] MUS.R.AFR.CENT.SER.8VO,(126)PP.204.(IN FRENCH)

[527] MATTHES,H. (1967)THE FISHES AND FISHERIES OF THE RUAHA RIVER BASIN ,TANZANIA.(SYSTEMATICS,ECOLOGY,ZOOGEOGRAPHY, FISHERIES).EAFFRO OCC.PAP.(9).PP19.

[528] MAUNGBOON,M. (1961)THE COLLECTION OF CLARIAS BATRACHUS FRY FROM NATURAL WATERS. THAI FISH.GAZ.14(2):100-111.

[529] MCALLISTER,D.E. (1968)THE EVOLUTION OF BRANCHIOSTEGALS AND ASSOCIATED OPERCULAR,GULAR AND HYIOD BONES AND THE CLASSIFICATION OF TELEOSTOME FISHES LIVING AND FOSSIL. BULL.NAT.MUS.CAN.(221).

[530] MCLAREN,D.S. (1961)SOURCES OF CAROTENE AND VITAMIN A IN LAKE PROVINE ,TANGANYIKA. ACTA TROP. 18(1):78-80

[531] MEECHAM,K. (1975)AQUACULTURE IN MALAWI. IN:FAO/CIFA SYMPOSIUM ON AQUACULTURE IN AFRICA. CIFA/75/SC.1.PP6.

[532] MEGGITT,F.J. (1930)REPORT ON A COLLECTION OF CESTODES MAINLY FROM EGYPT.1V.CONCLUSION PARASITOLOGY,22(3), 338-354.

[533] MEINKEN,H. (1961)[CONTRIBUTIONS TO THE KNOWLEDGE OF THE FAUNA OF THE SENEGAL RIVER. FISH FROM BAOULE AND BAKOY.] BULL.MUS.HIST.NAT. PARIS (2) 32:506-512 (IN FRENCH).

[534] MENON,A.G.K. (1950)ON A REMARKABLE BLIND SILUROID FISH OF THE FAMILY CLARIIDAE FROM KERALA(INDIA).REC.IND.MUS. 48:59-66.

[535] MENON,A.G.K. (1951)DISTRIBUTION OF THE CLARIID FISHES AND ITS SIGNIFICANCE IN ZOOGEGRAPHICAL STUDIES. PROC. .NAT. INST.SCI.IND. 17(4):291-299.

[536] MENON,A.G.K. (1951)ON CERTAIN FEATURES IN THE ANATOMY OF HORAGLANIS MENON.J.ZOOL.SOC.IND.3:249-253.

[537] MENON,M.A.S. (1961)ON A COLLECTION OF FISH FROM LAKE CHILKA,ORISSA. REC.INDIAN MUS. 59:41-69.

[538] MICHA,J.C. (1971)POND CULTURE OF AFRICAN CATFISHES. FAO AQUACULT.BULL. 3(4):6.

[539] MICHA,J.C. (1971)EXTERNAL BLISTERS IN CLARIAS. FAO AQUACULT.BULL. 3(4):13.

[540] MICHA,J.C. (1973)[SYNTHESIS OF RESEARCH ON REPRODUCTION,STOCKING AND PRODUCTION OF THE CLARIAS LAZERA.] CENTRE PISCOICOLE NATIONAL FAO.BANGUI. (IN FRENCH).

[541] MICHA,J.C. (1973)[STUDIES ON THE FISH POPULATIONS OF THE UBANGUI AND ATTEMPTS AT SELECTION AND ADAPTATION OF SEVERAL SPECIES TO AQUACULTURE PONDS].CENTRE TECHNIQUE FORESTIER TROPICAL. NOGENT-SUR-MARNE. PP110. (IN FRENCH).

[542] MICHA,J.C. (1974)[STUDIES ON THE FISH POPULATIONS OF THE UBANGUI AND ATTEMPTS AT SELECTION AND ADAPTATION

OF SEVERAL SPECIES TO AQUACULTURE PONDS].THESIS FOR THE DEGREE OF DOCTOR OF ZOOLOGICAL SCIENCES UNIV.OF LIEGE,BELGIUM. PP . (IN FRENCH).

[543] MICHA,J.C. (1975)[SYNTHESIS OF RESEARCH ON REPRODUCTION, STOCKING AND PRODUCTION IN AFRICAN CATFISH: CLARIAS LAZERA VAL.]BULL.FR.PISCIC.256:77-87(IN FRENCH)

[544] MICHA,J.C. (1975)SUMMARY OF PRELIMINARY TESTS FOR REPRODUCTION, REARING AND PRODUCTION OF AN AFRICAN SILURID: CLARIAS LAZERA. IN:FAO/CIFA SYMPOSIUM ON AQUACULTURE IN AFRICA GHANA,30SEPT 1975, FAO-CIFA--75/SE-5:PP23.

[545] MICHA,J.C. FRANK,V. (1975) [BIOLOGY OF THE PRINCIPAL SPECIES USED IN FISH CULTURE IN AFRICA.] IN: FAO/CIFA SYMPOSIUM ON AQUACULTURE IN AFRICA. CIFA/75/SR 8. PP39. (IN FRENCH).

[546] MILLS,H.D. (1956)THE AFRICAN MUDFISH CLARIAS LAZERA. IBADAN UNIVERSITY PRESS,IBADAN.PP48.

[547] MINSHULL,J.L. (1969)INTRODUCTION TO THE FOOD WEB OF LAKE SIBAYI, NORTHERN ZULULAND.SUPPL.NEWS LETT.LIMNOL.SOC.STH. AFR(13):20-25.

[548] MINSHULL,J. (1975)1974 FISHERY STATISTICS-LAKE KYLE.REPORT OF DEPT. OF NAT.PARK WDL ,MAN.RHODESIA.PP14.

[549] MOHANTY,S.K. (1975)FURTHER ADDITIONS TO THE FISH FAUNA OF THE CHILKA LAKE.J.BOMBAY NAT.HIST.SOC.72(3):863-866.

[550] MONOD,T. (1951)[CONTRIBUTION TO THE STUDY OF THE FAUNA OF MAURITANIA.FRESH-WATER FISHES.] BULL.INST.FRANC. AFR.NOIRE.13:802-812. (IN FRENCH)

[551] MONOD,T. (1955)[CONTRIBUTION TO THE STUDY OF THE FAUNA OF MAURITANIA. FRESH-WATER FISH (3RD NOTE).] BULL. INST.FRANC.AFR.NOIRE 17A:590-591. (IN FRENCH)

[552] MONOD,T. (1968)[THE UROPHORE COMPLEX OF TELEOST FISHES.] MEM. INST.FOND.AFRIQUE NOIRE (81) (IN FRENCH)

[553] MONTEL,R. (1905)[TRYPANOSOME OF A FISH OF COCHINCHINE.] C.R.SOC. BIOL.LVIII:1016-1017 (IN FRENCH)

[554] MOOKERJEE,H.K. MAZUMDAR,S.R. DASGUPTA,B. (1941) ON THE NATURE OF THE URINOGENITAL PAPILLA IN CLARIAS BATRACHUS AND HETEROPNEUSTES FOSSILIS. IND.J.VET.SCI.ANI.HUS.(2):334-336.

[555] MOOKERJEE,H.K. (1945)IDENTIFICATION OF EGGS OF COMMON FRESHWATER FISHES OF BENGAL.SCI.CULT.,CALCUTTA 11(1):48.

[556] MOOKERJEE,H.K. MAZUMDAR,S.R. (1950) SOME ASPECTS OF THE LIFE HISTORY OF CLARIAS BATRACHUS.PROC.ZOOL.SOC.BENGAL.3:71-79.

[557] MORAIS,R.T. (1974)[ICHTHYOLOGY OF THE ZAMBEZI RIVER-PROGRESS REPORT (MIMEO.REPORT)ICHTHYOLOGICAL STUDIES IN CABORA BASSA]AGRICULTURE&FOREST SERVICES DEPT.L.MARQUE MOZAMBIQUE. PP18.(IN PORTUGUESE)

[558] MORAVEC,F. (1974)THE DEVELOPMENT OF PARACAMALLANUS CYATHOPHARYNX (BAYLISS,1923):NEMATODA:CAMALLANIDAE).FOLIA PARASITOL.21(4):333-343.

[559] MORGAN,P.R. (1971)THE LAKE CHILWA TILAPIA AND ITS FISHERY. AFR.J. TROP. HYDROBIOL.FISH.1(1):51-58.

[560] MORTIMER,M.A.E. (1959)FISHING IN THE LUNDAZI DAM.RHOD.AGRIC.J.56(2):50-60.

[561] MORTIMER,M.A.E. BELL-CROSS,G. (1960) OBSERVATIONS ON THE MOVEMENT OF FISH IN A FISH LADDER. JOINT FISH RES.ORG.ANN.REP.NO.9:67.

[562] MORTIMER,M.A.E. (1964)A NOTE ON THE GROWTH OF CLARIAS(SILUROIDEA: CLARIIDAE).THE PUKU,OCC.PAP.DEPT.GAME&FISH.N. RHODESIA,NO.2:81-83.

[563] MORTIMER,M.A.E. (1965)FISH PRODUCTION FROM A STREAM IN NORTHERN RHODESIA.PROC.CENT.AFR.SCI.MED.CONG.LUSAKA,1963:405-

414.

[564] MOUSSA,T.A. (1956)MORPHOLOGY OF THE ACCESSORY AIR BREATHING ORGANS OF THE TELEOST,CLARIAS LAZERA(C&V.). J.MORPH.98(1):125-160.

[565] MOUSSA,T.A. (1957)PHYSIOLOGY OF THE ACCESSORY RESPIRATORY ORGANS OF THE TELEOST, CLARIAS LAZERA (C& V.).J.EXP. ZOOL. 136(3):419-54.

[566] MUKHERJEE,S. BHATTACHARYA,S. (1975) CHANGES IN THE KIDNEY PEROXIDASE ACTIVITY IN FISH EXPOSED TO SOME INDUSTRIAL POLLUTANTS. ENVIRON. PHYSIOL.BIOCHEM. 5(5):300-307.

[567] MUKHERJEE,S., BHATTACHARYA,S. (1977) VARIATIONS IN THE HEPATO PANCREATIC ALPHA AMYLASE ACTIVITY IN FISHES EXPOSED TO SOME INDUSTRIAL POLLUTANTS.WATER RESEARCH.11(1):71-74

[568] MUKHOPADHYAY,P.K. (1977)STUDIES ON THE ENZYMATIC ACTIVITIES RELATED TO VARIED PATTERN OF DIETS IN THE AIR BREATHING CATFISH CLARIAS BATRACHUS.HYDROBIOLOGIA.52(2-3): 235-237.

[569] MUNRO,J.L. (1965)FEEDING RELATIONSHIPS AND PRODUCTION OF FISH IN A SOUTHERN RHODESIAN LAKE.PH.D. THESIS,UNIVERSITY OF LONDON,ENGLAND.PP277.

[570] MUNRO,J.L. (1967)THE FOOD OF A COMMUNITY OF EAST AFRICA FRESH-WATER FISHES.J.ZOOL.LOND. 151:389-415.

[571] MUNSHI,J.S. (1961)THE ACCESSORY RESPIRATORY ORGANS OF CLARIAS BATRACHUS(LINN.).J.MORPH.109:115-139.

[572] MUNSHI,J.S.D. SINGH,B.N. (1968) THE STRUCTURE OF THE RESPIRATORY EPITHELIUM OF CERTAIN FRESHWATER AND AIR BREATHING FISHES. J.ZOOT.(INDIA), 9:91-107.

[573] MUNSHI,J.S.D. SINGH,B.N. (1968) ON THE MICRO-CIRCULATORY SYSTEM OF THE GILLS OF CERTAIN FRESHWATER TELEOSTAN FISHES. J.ZOOL. 154(3):365-376

[574] MURRAY,J.L. (1975)SELECTION OF ZOOPLANKTON BY CLARIAS GARIEPINUS (BURCHELL) IN LAKE MCILWAINE A EUTROPHIC RHODESIAN RESERVOIR.MSC THESIS,UNIV.OF RHODESIA, RHODESIA.PP82.

[575] MYERS,G.S. (1938)NOTES ON ANSORGIA,CLARISILURIS,WALLAGO AND CERATOGLANIS, FOUR GENERA OF AFRICAN AND INDO-MALAYAN CATFISHES. COPEIA ANN. ARBOR :98

[576] MYERS,G.S. (1956)A NOTE ON THE ABYSSINIAN CATFISH,CLARIAS DEPRESSUS MYERS. STANF. ICTHYOL.BULL. 7:14.

[577] MYERS,B.J. KUNTZ,R.E. WELLS,W.H. (1962) HELMINTH PARASITES OF REPTILES,BIRDS AND MAMMALS IN EGYPT V11. CHECK LIST OF NEMATODES COLLECTED FROM 1948 TO 1955.CAN.J.ZOOL.40(4): 531-538.

[578] NAGVI,R.H. (1966)HYDROSTATIC SIGNIFICANCE OF ACCESSORY RESPIRATORY ORGANS IN SOME AIR BREATHING FISHES. COPEIA :136-139.

[579] NAIR,P.V. (1960)STUDIES ON THE ACCESSORY GLANDS AND TESTES OF SOME SILUROID FISHES.II.CLARIAS BATRACHUS. IND. SCI.CONGR.ASS.PROC.47.SESS.PART III:477.

[580] NAIR,P.V. (1965)STUDIES OF THE MALE REPRODUCTIVE SYSTEM OF SOME SILUROID FISHES. PARTS 1 AND2. INDIAN ZOOTOM.MEM. NO.9:1-35.

[581] NALAVADE,M.N., VARUTE,A.T. (1976) HISTOCHEMICAL STUDIES ON THE MUCINS OF THE VERTEBRATE TONGUES.PART 7:HISTOCHEMICAL ANALYSIS OF MUCO SUBSTANCES IN THE TONGUE OF SOME FISHES. ACTA HISTOCHEMICA.57(2):245-262.

[582] NARAIN,D. (1937)CYTOPLASMIC INCLUSIONS IN THE OOGENESIS OF SACCHOBRANCHUUS FOSSILIS, CLARIAS BATRACHUS,AND

ANABAS SCANDENS.Z.ZELLFORSCH.26:623-640.

[583] NAWAR,G. (1954)ON THE ANATOMY OF CLARIAS LAZERA.1. OSTEOLOGY. J.MORPH. 94:551-586.

[584] NAWAR,G. (1955)ON THE ANATOMY OF CLARIAS LAZERA.2. THE MUSCLES OF THE HEAD AND PECTORAL GIRDLE.J.MORPH.97:23-38

[585] NAWAR,G. (1955)ON THE ANATOMY OF CLARIAS LAZERA,3.THE VASCULAR SYSTEM. J.MORPH.97(2):179-214.

[586] NAWAR,G. (1959)OBSERVATIONS ON THE SEMINAL VESICLE OF THE NILE CATFISH CLARIAS LAZERA.ANN.MAG.NAT.HIST. 13 SER.2:444-448.

[587] NAWAR,G.
YOAKIM,E.G. (1962)
A STUDY ON THE FECUNDITY OF THE NILE CAT-FISH CLARIAS LAZERA VALENCIENNES & CUVIER 1840. ANN. MAG.NAT.HIST.(13),5(55):385-389.

[588] NELLEN,W .
RUCKES,E. (1975)
[EXPLOITING LAKE ASSAD, ARABIAN REPUBLIC SYRIA, FOR FISHING AND POSSIBILITIES OF SUPPORT FROM THE GERMAN FEDERAL REPUBLIC .] (TECHNICAL ASSISTANCE PROGRAM). KIEL,GFR. PP124. (IN GERMAN).

[589] NICHOLS,J.T.
GRISCOM,L. (1917)
FRESHWATER FISHES OF THE CONGO BASIN OBTAINED BY THE AMERICAN MUSEUM CONGO EXPEDITION 1909-1915. NEW YORK BULL.AMER.MUS.NAT.HIST. 37:653-756.

[590] NICHOLS,J.T.
LA-MONTE,F.R. (1933)
NEW FISHES FROM THE KASAI DISTRICT OF THE BELGIAN CONGO.AMER.MUS.NOV.N.Y. 656:1-6.

[591] NICHOLS,J.T.
LA-MONTE,F.R. (1934)
MORE NEW FISHES FROM THE KASAI DISTRICT OF THE BELGIAN CONGO. AMER.MUS.NOV.N.Y. 723:1-6.

[592] NICHOLS,J.T.
LA-MONTE,F.R. (1953)
TWO NEW AFRICAN CATFISHES FROM EASTERN FRENCH EQUATORIAL AFRICA.AMER.MUS.NOVIT.NO.1648:1-4.

WATER FISH IN LAKE VICTORIA,UGANDA(NEMATODA: HEDRURIDAE).REVUE ZOOL.BOT.AFR.75(1/2):77-81.

[593] NORMAN,J.R. (1928)TWO NEW FISHES FROM LAKE VICTORIA. ANN.MAG.NAT. HIST. 2:104-106.

[594] OGILVIE,V.E. (1967)THE GENERAL BIOLOGY OF CLARIAS BATRACHUS.(REVIEW OF WORK OF THAI FISHERIES DEPT).FLORIDA GAME AND FRESH WATER FISH COMMISSION.(MIMEO.REPORT).PP3

[595] OGILVIE,V.E.
GOODRICK,R.L. (1968)
A PRELIMINARY INVESTIGATION OF AN ALBINO CLARIID CATFISH,CLARIAS SP.INTRODUCED INTO PALM BEACH COUNTRY.FLORIDA GAME AND FRESHWATER FISH COMMISSION.(MIMEO.REPORT).PP19.

[596] OSUJI,F.N.C. (1974)COMPARATIVE STUDIES ON THE SUSCEPTIBILITIES OF DIFFERENT GENERA OF DRIED FISH TO INFESTATION BY DERMESTES MACULATUS AND NECROBIA RUFIPES. NIGER.J.ENTOMOL.1(1):63-68.

[597] OVAIS,M. (1974)ON THE PRESENCE OF A FORKED BARBEL IN THE CATFISH CLARIAS BATRACHUS(LINN.).CURR.SCI. 43(4):125-126.

[598] OVAIS,M.
GUPTA,S.S. (1975)
HISTAMINE CONTENT OF THE DIGESTIVE TRACT OF A CATFISH CLARIAS BATRACHUS IN RELATION TO SPAWNING, SEX, AND SEASONAL VARIATION.IND.J.PHYSIOL.& PHARMACOL.19(2):90-93.

[599] OVAIS,M.
GUPTA,S.S.
BHAGWAT,A.W. (1976)
ACETYL CHOLINE CONTENT OF THE BRAIN AND VISCERAL ORGANS OF THE TELEOSTAN FISH HETEROPNEUSTES FOSSILIS AND CLARIAS BATRACHUS.IND.J.PHYSIOL.& PHARMACOL.20(3):177-179.

[600] OVAIS,M. (1976)ADRENERGIC AND CHOLINERGIC RECEPTORS IN THE STOMACH OF THE TELEOSTEAN FISHES CLARIAS BATRACHUS AND HETEROPNEUSTES FOSSILIS.IND.J.ZOOL.4(1):7-13.

[601] PALMER,C.A. (1973)A PRELIMINARY INVESTIGATION OF SOME ASPECTS OF FEEDING AND DIGESTION IN CLARIAS GARIEPINUS. B.SC PROJECT,UNIVERSITY OF RHODESIA,RHODESIA.PP92.

[602] PANCERI,P. (1873)[ON ALBINISM IN CLARIAS ANGUILLARIS.]REND.ACC. NAP. SEPT.1873.(NOT SEEN IN ORIGINAL-IN ITALIAN)

[603] PANCERI,P. (1874)[NOTE ON MALE ORGANS OF CLARIAS ANGUILLARIS.] ANN.MUS.GENOV.VI.PP.361-365.(NOT SEEN IN ORIGINAL IN ITALIAN)

[604] PANDEY,K.C. (1970)ON A NEW PHYLLODISTOME METACERCARIA (PHYLLODISTOMUM LUCKNOWENSIS) AND ITS ADULT. INDIAN J.ZOOTOMY,11(1):22-35.

[605] PAPERNA,I. (1961)STUDIES ON MONOGENETIC TREMATODES IN ISRAEL.3. MONOGENETIC TREMATODES OF THE CYPRINIDAE AND CLARIIDAE OF THE LAKE GALILLEE.BAMIDGEH,13(1): 14-29.

[606] PAPERNA,I. (1965)MONOGENETIC TREMATODES COLLECTED FROM FRESHWATER FISHES SOUTHERN GHANA.BAMIDGEH, 17(4):107-111.

[607] PAPERNA,I. (1968)ECTOPARASITIC INFECTIONS ON FISH OF VOLTA LAKE, GHANA.BULL.WILDLIFE DIS. ASS. 4(4):135-137.

[608] PAPERNA,I. THURSTON,J.P. (1968) MONOGENETIC TREMATODES(DACTYLOGYRIDAE) FROM FISH IN UGANDA. REVUE ZOOL. BOT. AFR. 79(3/4): 284-294.

[609] PAPERNA,I. (1969)MONOGENETIC TREMATODES OF THE FISH OF THE VOLTA BASIN AND SOUTH GHANA. BULL.INST.FR. AFR.NOIRE,31, SER. A:840-880.

[610] PAPERNA,I. (1974)HOSTS,DISTRIBUTION AND PATHOLOGY OF INFECTIONS WITH LARVAE OF EUSTRONGYLIDES(DIOCTOPHYMIDAE, NEMATODA)IN FISHES FROM EAST AFRICAN LAKES. J.FISH BIOL.6(1):67-76

[611] PARDUE,G.B. (1970)TEMPERATURE TOLERANCE OF CLARIAS BATRACHUS. FAO FISH.CULT.BULL.2(3):6.

[612] PATHAK, S.P. REDDY, B.R. (1962) THE COMPONENT ACIDS OF THE FATS OF SOME INDIAN FRESHWATER FISHES. BIOCHEM.J.85:618-620.

[613] PAWAPOOTANON,O. (1965)[THE CULTIVATION OF PLADUK(CLARIAS BATRACHUS L.) IN THE VICINITY OF BANGKOK].THESIS,KASETSART UNIVERSITY,FISHERIES FACULTY(IN THAI).

[614] PELLEGRIN,J. (1913)[NEW FISH FROM THE OGOOUE CAUGHT BY ERNEST HAUG.] PARIS BULL.SOC.ZOOL. 38:272-275 (IN FRENCH)

[615] PELLEGRIN,J. (1920)[FISHES FROM LAGOONS OF THE IVORY COAST. DESCRIPTION OF TWO NEW SPECIES.] PARIS BULL.SOC ZOOL. 45:115-121 (IN FRENCH).

[616] PELLEGRIN,J. (1928)[FISH FROM CHILOANGO AND THE CONGO COLLECTED BY THE EXPEDITION OF DR SCHOUTENDEN (1920-1922).] ANN.MUS.CONGO BELGE ZOOL. 3(1):49 (IN FRENCH)

[617] PELLEGRIN,J. (1930)[FISHES OF THE OGOOUE, KOUILOU, ALIMA AND OF THE SANGHA COLLECTED BY MR.A.BANDON.DESCRIPTIONS OF FIVE SPECIES AND FIVE NEW GENUS.]BULL.SOC.ZOOL. FR.55:196-210. (IN FRENCH)

[618] PETEL,G . PLANQUETTE,P. (1975) [THE USE OF ELECTRIC FISHING IN TROPICAL FISH CULTURE PONDS.]NOTES DOC.PECHE PISCIC.(NOUV.SER.) 11:75-85.(IN FRENCH).

[619] PETERS,W.C.H. (1868)[NATURAL HISTORY RESEARCH CRUISE TO MOSSAMBIQUE.] ZOOLOGIE.IV.FLUSSFISCHE.BERLIN,PP.32.(IN GERMAN)

[620] PETERS,W.C.H. (1882) SB.NAT.FR.PP.374.

[621] PETRICK,F.O. (1975)ANATOMY OF THE SWIMBLADDERS OF 7 FAMILIES OF TRANSVAAL FRESH WATER FISHES.J.LIMNOL.SOC.STH.AFR. 1(1):17-22.

[622] PFEFFER,G. (1896)[THE ANIMAL WORLD OF EAST AFRICA AND NEIGHBOURING REGIONS.]LEIF.V. DIE FISCHE OST-AFRIKAS.BERLIN. PP72.

[623] PHAM,A.
RAUGEL,B. (1975)
[REARING OF SILURID FISH AT BOUAKE (IVORY COAST). RESULTS OBTAINED FROM DECEMBER 1974 TO JUNE 1975.] NOTES DOC.PECHE PISCIC.(NOUV.SER.) 11:36-42. (IN FRENCH)

[624] PHAM,A. (1975)[ON THE MASS PRODUCTION OF ALEVINS OF CLARIAS LAZERA VAL. IN THE BOUAKE STATION,IVORY COAST.] NOTES ET DOCUMENTS SUR LA PECHE ET LA PISCICULTURE. NOUVELLE CENTRE TECHNIQUE FORESTIER TROPICAL. SERIE NO 10:49-579(IN FRENCH

[625] PIENAAR,U.DE V. (1968)THE FRESHWATER FISHES OF THE KRUGER NATIONAL PARK. KOEDOE (11): 1-82.

[626] PIETSCHMANN,V. (1939)[ZOOLOGIC MATERIAL FROM ERITREA COLLECTED BY G.MULLER, DURING THE EXPEDITION OF THE MILAN INSTITUTE OF SCIENCE AND THE MUSEUM OF TRIESTE. PART VI FISH.] ATTI.MUS.STOR.NAT.TRIESTE 14:179 -186 (IN ITALIAN).

[627] PIKE,T. (1968)EFFECTS OF SEASONAL CHANGES ON THE ACTIVITY OF TILAPIA MOSSAMBICA AND CLARIAS GARIEPINUS IN LAKE SIBAYI.MIMEO.REPORT BY NATAL PARKS,GAME AND FISH PRESERVATION BOARD.PP6.

[628] PIKE,T (1969)COMPOSITION,RELATIVE ABUNDANCE AND SIZE RANGE OF FISH POPULATIONS IN LAKE SIBAYI. SUPP.NEWS LETT. LIMNOL.SOC.STH.AFR.13:38-43.

[629] PILLAY,T.V.R.(ED.) (1967)PROCEEDINGS OF THE WORLD SYMPOSIUM ON WARM-WATER POND FISH CULTURE.VOL1.FAO,ROME.PP.55.

[630] PITMAN,C.R.S. (1900)SURVEY OF THE FAUNA OF N. RHODESIA.N.RHOD.J.

[631] PIVNICKA,K. (1974)THE ZAMBEZI BARBEL CLARIAS GARIEPINUS(BURCHELL, 1822).IN:LAKE KARIBA A MAN MADE TROPICAL ECOSYSTEM IN CENTRAL AFRICA: FISH PRODUCTION OF A TROPICAL ECOSYSTEM.(ED.)BALON,E.K. :318-325.

[632] POLLARD,H.B. (1892)THE LATERAL LINE SYSTEM IN SILURIDS.MORPH.JB. ANAT.V:525-551.

[633] POLL,M. (1941)[A SYSTEMIC STUDY AND MORPHOLOGY OF A COLLECTION OF FISH FROM UELE(BELGIAN CONGO) INCLUDING THREE NEW SPECIES.]BULL.MUS.HIST.NAT.BELG.17(50):18- .(IN FRENCH).

[634] POLL,M. (1941)[NEW FISH FROM THE IVORY COAST].REV.ZOOL.BOT.AFR. 34:133-143. (IN FRENCH).

[635] POLL,M. (1942)[THE FISHES OF LAKE TUMBA, BELGIAN CONGO]. BULL. MUS.ROY.HIST.NAT.BELG.18(36):1-25.(IN FRENCH).

[636] POLL,M. (1942)[DESCRIPTION OF A NEW GENUS OF CLARIAS ORIGINALLY FROM THE BELGIAN CONGO.]REV.ZOOL.BOT.AFR.36:94-100.(IN FRENCH)

[637] POLL,M. (1942)[NOTES ON THE OSTEOLOGY OF THE DOLICHALLABES MICROPHTHALMUS(POLL) AND REMARKS ON THE EVOLUTION OF THE CLARIIDAE.]ANN.SOC.ZOOL.BELG. 73:222-235.(IN FRENCH).

[638] POLL,M. (1943)[DESCRIPTIONS OF NEW FISH FROM LAKE TANGANYIKA BELONGING TO THE FAMILYS, CLARIIDAE AND CICHLIDAE] REV.ZOOL.BOT.AFR.37:303-318.(IN FRENCH).

[639] POLL,M. (1943)[DESCRIPTION OF TANGANIKALLABES MORTIAUX GEN.NOV. SP.NOV. OF THE FAMILY CLARIIDAE.]REV. ZOOL. BOT. AFR.37:129-133.(IN FRENCH)

[640] POLL,M. (1946)[A REVISION OF THE ICTHYOLOGIC FAUNA OF LAKE TANGANYIKA.] ANN.MUS.CONGO ZOOL.SER.1(4):141-364. (IN FRENCH).

[641] POLL,M. (1948)[FISHES COLLECTED IN KATANGA BY H.J.BREDO]. BULL. MUS.HIST.NAT.BELG.24(21):1-24.(IN FRENCH).

[642] POLL,M.
LAMBERT,J. (1958)
A NEW CYPRINOID AND A NEW CLARIIDAE FROM THE CONGOLESE FOREST. REV.ZOOL.BOT.AFR. 56:328-339.

[643] POLL,M.
GOSSE,J.P. (1963)
[CONTRIBUTION TO THE SYSTEMATIC STUDY OF THE

ICTHYOLOGICAL FAUNA OF THE CENTRAL CONGO.] ANN. MUS.ROY.AFR.CENTRALE ZOOL. 116:43-111 (IN FRENCH)

[644] POLL,M. (1967)[CONTRIBUTION TO THE ICTHYOLOGICAL FAUNA OF ANGOLA.] PUBL. CULT.CO.DIAM. ANGOLA 75:15-381 (IN FRENCH)

[645] POOLEY,A.C. (1972)FEEDING BEHAVIOUR OF BARBEL(CATFISH). LAMMERGEYER NO 17:65.

[646] POPTA C.M.L. (1905)[PRELIMINARY DESCRIPTIONS OF NEW SPECIES OF FISH COLLECTED IN CENTRAL BORNEO BY DR A.W. NIEUWENHUIS IN 1898 AND 1900.] NOTES LEYDEN MUS. XXIV :179-202 (IN FRENCH)

[647] POPTA,C.L. (1919)DESCRIPTION OF CLARIAS NIGERIAE N.SP. FROM THE WARI,MOUTH OF THE NIGER. WEST AFRICA ZOOL.MED. LEIDEN 5:4

[648] POTT,R.MCC. (1969)THE FISH - LIFE OF THE PONGOLO RIVER BEFORE THE CLOSURE OF THE J.G. STRIJDOM DAM.M.SC. THESIS, UNIVERSITY OF WITWATERSRAND,RSA.

[649] POTT,R.MCC. (1969)THE FISH LIFE OF THE PONGOLA RIVER BEFORE THE CLOSURE OF THE J.G.STRYDOM DAM. M.SC. THESIS,UNIV. OF WITWATERSRAND, SOUTH AFRICA.

[650] PRAKASH,R. (1953)THE HEART OF THE COMMON INDIAN CATFISH, HETEROPNEUSTES FOSSILIS (BLOCH),WITH SPECIAL REFERENCE TO THE CONDUCTING SYSTEM.PROC.ZOOL. SOC.BENG.6:113-118.

[651] PRUDHOE,S. (1957)TREMATODA. EXPLOR. PARC NATN. UPEMBA MISS.G.F. DE WITTE,(1946-49), 48:1-28.

[652] QUASIM,S.Z. ET AL. (1960) THE MEASUREMENT OF CO2 PRODUCED BY AIR BREATHING FISHES AND EVIDENCE OF THE RESPIRATORY FUNCTION OF THE ACCESSORY RESPIRATORY ORGANS.PROC. INDIAN ACAD.SCI.52 B(1):19-26.

[653] QUASIM,S.Z. HASAN,R. (1961) A HYDROSTATIC FUNCTION OF THE ACCESSORY RESPIRATORY ORGANS IN AIR-BREATHING FISHES. NATURE, LOND.191 (4786):396-397.

[654] RABIE,F. ABDEL MAGID,A.M. GUMA,A,K.A. KARRAR,O. (1972) EVOLUTION OF CATALASE IN FISH. COMP.BIOCHEM. PHYSIOL.(PHYSIOL.) 43A(4):1053-1055.

[655] RADEK,G. (1972)AQUARIUM FISH ENDANGER THE FISH FAUNA OF FLORIDA. AQUARIEN MAG.6(11):482-483.

[656] RAHIMULLAH,M. DAS,B.K. (1936) OBSERVATIONS ON THE CONFLUENCE OF THE MEDIAN FINS IN AN AIR BREATHING CATFISH 'MAROOF'CLARIAS BATRACHUS LINN).BULL.SOC.PORTUG.SCI.NAT.12(20):155-165.

[657] RAI,P. PANDE,B.P. (1967) ON THE TREMATODE GENUS EUMASENIA (MASENIIDAE)IN INDIAN FRESHWATER FISHES.AGRA.UNIV.J.RES.16(3):1-5

[658] RAJBANSHI,V.K. (1966)ON THE CUTANEOUS TASTEBUD SYSTEM OF A FEW FRESHWATER FISHES. SCI. CULT. 32:381-382.

[659] RAJBANSHI,V.K. (1966)A NOTE ON THE STUDY OF GUSTATORY SENSES OF THE BARBELS OF FRESHWATER TELEOSTS.NATURWISSENSCHA-FTEN.53(8):208-209.

[660] RAJBANSHI,V.K. (1966)THE CUTANEOUS SENSE-ORGANS OF CLARIAS BATRACHUS (GRONOVIOUS).ZOOL.ANZ.176(5):379-384.

[661] RAMASWAMI,L.S. SUNDARARAJ,B.I. (1956) INDUCED SPAWNING OF THE INDIAN CATFISH. SCIENCE, 123:1080.

[662] RAMASWAMI,L.S. SUNDARARAJ,B.I. (1957) SOME ASPECTS OF INDUCED SPAWNING IN CATFISH HETEROPNEUSTES. NATURWISS,44:384.

[663] RAMASWAMI,L.S. SUNDARARAJ,B.I. (1957) INDUCED SPAWNING OF THE INDIAN CATFISH

HETEROPNEUSTES FOSSILIS WITH PITUITARY INJECTIONS. ACTA ANAT.31:551-562.

[664] RAMASWAMI,L.S.
LAKSHAMAN,A.B. (1958)
SPAWNING CATFISH WITH MAMMALIAN HORMONES.NATURE, LOND.182:122-123.

[665] RAMASWAMI,L.S.
SUNDARARAJ,B.I. (1958)
INDUCED SPAWNING IN THE CATFISH HETEROPNEUSTES FOSSILIS WITH MAMMALIAN CHORIONIC GONADOTROPHINS. ACTA ANAT.32:230-234.

[666] RAMASWAMI,L.S.
LAKSHAMAN,A.B. (1959)
ACTION OF MAMMALIAN HORMONES ON THE SPAWNING OF CATFISH.J.SCIENT.IND.RES.18C:185-191.

[667] RAMA DEVI,P. (1973)LYTOCESTUS LONGICOLLIS SP. NOV.(CESTOIDEA: CARYOPHYLLIDEA) FROM THE CATFISH CLARIAS BATRACHUS (LINN.) IN INDIA. J.HELMINTHOL. 47(14) :415-420.

[668] RAO,P.D.P. (1966)ON THE SACCUS VASILOSUS IN SOME INDIAN FRESHWATER TELEOSTS.NATURWISSENSCHAFTEN, 53(9): 233.

[669] RAO,P.D.P.
BETOLE,U.K.
KONDAWAR,V.V. (1972)
CHANGES IN THE PITUITARY-INTER-RENAL AXIS AFTER A GONADECTOMY IN THE CATFISH,CLARIAS BATRACHUS (LINN.).ACTA ZOOL.STOCKH.53(2):135-145.

[670] RAO,P.D.P.
BETOLE,U.K. (1973)
CHANGES IN THE HYPOTHALAMO-NEUROHYPOPHYSIAL COMPLEX AFTER GONADECTOMY IN THE CATFISH CLARIAS BATRACHUS(LINN.),.ZOOL.BEITR.19(3):319-333.

[671] RAO,P.D.P.
DABHADE,K.B. (1973)
IN VITRO DEMONSTRATION OF CHANGES IN THE PREOPTICONEUROHYPOPHYSIAL COMPLEX AFTER ELECTRIC STIMULATION OF THE OLFACTORY TRACT IN THE CATFISH,CLARIAS BATRACHUS(LINN.). Z.ZELLFORSCH. MIKROSK.ANAT.139(1):95-99.

[672] RASHEED, S. (1965)A PREMLIMINARY REVIEW OF THE GENES RHABDOCHONA RAILLIET,1916, WITH DESCRIPTION OF A NEW AND RELATED GENUS, ACTA PARASIT, POL) 13(39/46): 407-424.

[673] RATCLIFF,C. (1972)TRAWLING-CENTRAL AFRICA STYLE.FISH.NEWS INTER. SEPT 1972 :22-29.

[674] RATCLIFFE,C. (1974)COMMERCIAL SMALL CRAFT PAIR TRAWLING TRIALS LAKE CHILWA MALAWI 1971.AFR.J.TROP HYDROBIOL.FISH.3(1): 61-78.

[675] REED,W. (1967)FISH AND FISHERIES OF NORTHERN NIGERIA. NIGERIA,MIN.OF AGRIC. PP 226.

[676] REGAN,C.T. (1908)DESCRIPTIONS OF NEW FRESHWATER FISHES FROM CHINA AND JAPAN . ANN.MAG.NAT.HIST.,LONDON. 1:149-153.

[677] REJ,S.
MITRA,B. (1975)
VARIATION OF MUSCLE LIPIDS IN DIFFERENT SEASONS SIZE AND WEIGHT GROUPS OF CLARIAS BATRACHUS.IND. SCI.CONG.ASSOC.PROC.62:195.

[678] RICARDO-BERTRAM,C.K. (1943)THE FISHES OF THE BANGWEULU REGION.J.LINN.SOC. LONDON.41:183-217.

[679] RISHI,K.K. (1976)KARYOTYPIC STUDIES ON 4 SPECIES OF FISHES. NUCLEUS(CALCUTTA).19(2):95-98.

[680] RIZKALLA,W. (1963)THE PITUITARY GLAND OF THE TELEOST CLARIAS LAZERA C&V. PROC.ZOOL.SOC.U.A.R. 1:127-146.

[681] RIZKALLA,W. (1969)STUDIES ON THE THYMUS OF THE NILE TELEOST FISH, CLARIAS LAZERA C&V. ACTA VET.HUNG. 19:331-342

[682] RIZKALLA,W. (1969)STUDIES ON THE ADRENAL GLANDS AND CORPUSCLES OF STANNIUS OF THE TELEOST NILE FISH CLARIAS LAZERA C&V. ACTA VET. HUNG. 19:343-350.

[683] RIZKALLA,W. (1970)STUDIES ON THE GONADS OF THE TELEOST NILE FISH CLARIAS LAZERA C&V.,WITH SPECIAL REFERENCE TO THEIR ENDOCRINE TISSUES. ACTA VET.HUNG. 20:129-138

[684] RIZKALLA,W. (1970)THE MORPHOLOGY OF THE PINEAL ORGAN IN THE TELEOST CLARIAS LAZERA C&V. ACTA BIOL. HUNG. 21:25-33

[685] ROUSHDY,H.M.
IMAN,A.E. (1971)
GROWTH RATE MEASUREMENTS IN GAMMA IRRADIATED CATFISH Z,NATURFORSCH.(B),26(9):962-967.

[686] ROUSHDY,H.M.
EL-KASHEF,H.S.
IMAN,A.E. (1975)
BIOCHEMICAL AND PHYSIOLOGICAL CHANGES IN EGYPTIAN NILE FISH SUBJECTED TO VARYING LEVELS OF GAMMA IRRADIATION.IN:IMPACTS OF NUCLEAR RELEASES INTO THE AQUATIC ENVIRONMENT. INT .ATOMIC ENERGY ASSOC. STI/PUB-406(VIENNA):415-430.

[687] ROY,A.
SEN,P.B. (1941)
THE ESTIMATION OF CYSTINE,TYROSINE.AND TRYPTOPHANE IN SOME COMMON EDIBLE FISHES OF BENGAL.ANN.BIOCHEM.EXP.MED.1:321-324.

[688] SACHS,W.B. (1961)[AN ALBINO CLARIAS.] AQUAR. TERRAR. Z. 14:268-269 (IN GERMAN)

[689] SAHA,K.C. (1939)DISTRIBUTION OF FREE AND TOTAL ASCORBIC ACID IN THE LIVER AND MUSCLE OF BENGAL FRESHWATER FISH. J.INDIAN CHEM SOC.16:511-514.

[690] SAHA,K.C. (1941)NICOTINIC ACID CONTENT OF FISH.ANN.BIOCHEM.EXP. MED.1:75-78.

[691] SAHA,M.P.
PANDEY,P.K.
OJHA,J.
MUNSHI,J.S.D. (1976)
HISTOCHEMICAL LOCALIZATION OF LIPID IN THE RESPIRATORY MUSCLES OF A FRESH WATER AIR BREATHING SILUROID FISH CLARIAS BATRACHUS.ZEITSCHRIFT FUER MIKROSKOPISCH-ANATOMISCHE FORSCHUNG.90(2): 290-301.

[692] SAIGAL,B.N.
GHOSH,A.
DATTA,A.K. (1974)
OBSERVATIONS ON THE CARBOHYDRATE DIGESTING ENZYMES IN CARNIVOROUS CATFISH HETEROPNEUSTES FOSSILIS. J.INL.FISH.SOC.INDIA.6:83-84.

[693] SAKSENA,J.N. (1960)STUDIES ON A NEW SPECIES OF THE GENUS ORIENTOCREADIUM (TREMATODA(ALLOCREADIIDAE) FROM THE INTESTINE OF CLARIAS MAGUR. PROC. NATN ACAD. SCI.INDIA,(B),30:83-86.

[694] SANDON,H. (1950)AN ILLUSTRATED GUIDE TO FRESHWATER FISHES OF THE SUDAN.SUDAN NOTES AND RECORDS,KHARTOUM.

[695] SANGA,H.C.S.
BHATT,S.D.
BISHT,J.S. (1975)
THE CAUDAL NEUROSECRETORY SYSTEM OF A FRESHWATER TELEOST CLARIAS BATRACHUS (LINN.) ZOOL .BEITR .,21(3):473-481.

[696] SAOUD,M.F.A. (1976)A GENERAL SURVEY OF THE PROTOZOAN BLOOD PARASITES OF SOME NILE FISHES FROM THE SUDAN.REV.ZOOL.AFR. 90(2):313-322.

[697] SARIG,S. (1964)FISHERIES AND FISH CULTURE IN ISRAEL IN 1963. BAMIDGEH 16(4):111-128.

[698] SARIG,S. (1966)FISHERIES AND FISH CULTURE IN ISRAEL IN 1965. BAMIDGEH 18(2):37-50.

[699] SARIG,S. (1967)FISHERIES AND FISH CULTURE IN ISRAEL IN 1966. BAMIDGEH 19(2/3):29-45.

[700] SARIG,S. (1977)FISHERIES AND FISH CULTURE IN ISRAEL IN 1976. BAMIDGEH 29(3):75-86.

[701] SASTRY,K.V. (1972)AMINO TRIPEPTIDASE ACTIVITY IN THREE TELEOST FISHES.PROC.INDIAN ACAD.SCI.(B),76(6):251-257.

[702] SASTRY,K.V. (1974)DISTRIBUTION OF LIPASE IN THE DIGESTIVE TRACT OF TWO TELEOST FISHES. ACTA.HISTOCHEM.48(2): 320-325.

[703] SASTRY,K.V. (1975)OPTIMUM PH FOR THE ACTIVITY OF LIPASE IN THE DIGESTIVE SYSTEM OF 2 TELEOST FISHES.ZOOLOGICA

POLONIAE 25(2/3):23-30.

[704] SASTRY,K.V. (1977)PEPTIDASE ACTIVITY IN A FEW TELEOST FISHES.ZOOL. BEITR.23(1):29-36.

[705] SATHYANESAN,A.G. (1961)A COMPARATIVE STUDY OF THE PITUITARY OF SOME FRESHWATER TELEOSTS.REC.INDIAN MUS.59:305-319.

[706] SATHYANESAN,A.G. (1968)PATTERN OF THE HYPO-THALAMO-NEUROHYPOPHYSIAL COMPLEX IN SOME INDIAN TELEOSTS,AS REVEALED BY BULK STAINING PROCEDURES. ADVAN.ABSTR.CONTRIB. FISH,AQUATIC SCI.INDIA 11(4):41.

[707] SATHYANESAN,A.G. (1969)AN IN SITU STUDY OF THE PREOPTICO-NEUROHYPO-PHYSIAL COMPLEX OF THE FRESHWATER TELEOST CLARIAS BATRACHUS(L.).Z.ZELLFORSCH.98:202-216.

[708] SATHYANESAN,A.G. (1970)THE NUCLEUS LATERALIS TUBERIS OF CLARIAS BATRACHUS L. ZOOL.BEITR. 16:257-262

[709] SATHYANESAN,A.G. (1970)THE NUCLEUS PREOPTICUS OF THE FRESHWATER CATFISH CLARIAS BATRACHUS L. ZOOL.BEITR. 16:247-255

[710] SATHYANESAN,A.G. (1971)STRUCTURE OF THE NEUROHYPOPHYSIS OF CLARIAS BATRACHUS(L.) WITH A NOTE ON THE TETRAPOD-LIKE CHARACTERISTIC OF ITS HYPOTHALAMOHYPOPHYSICAL VASCULARIZATION.Z.MIKROSK.-ANAT.FORSCH.84(1): 37-51.

[711] SATHYANESAN,A.G. (1971)STRUCTURE OF THE HYPOTHALAMUS OF THE FRESHWATER TELEOST CLARIAS BATRACHUS(L.).Z.MIKROSK.-ANAT. FORSCH.84(4):429-448.

[712] SATHYANESAN,A.G. (1971)HYPOTHALMO-HYPOPHYSIAL VASCULARIZATION AND ITS RELATIONSHIP WITH THE NEUROSECRETORY SYSTEM IN THE CATFISH CLARIAS BATRACHUS(L.).ZOOL.BEITR. 17(1):1-11.

[713] SATHYANESAN,A.G. (1972)HYPOTHALMO-HYPOPHYSIAL VASCULARIZATION IN A TELEOST FISH WITH SPECIAL REFERENCE TO ITS TETRAPODAN FEATURES.ACTA ANAT.81(3):349-366.

[714] SATHYANESAN,A.G. HAIDER,S. (1975) A COMPARISON BETWEEN THE CORTICOTROPHS AND THE LEAD HEMATOXYLIN POSITIVE CELLS OF THE PARS INTERMEDIA IN THE FRESH WATER TELEOST CLARIAS BATRACHUS.IND.SCI.CONG.ASSOC.PROD.62:159.

[715] SATHYANESAN ,A.G . JOY,K.P. (1976) MONOAMINE OXIDASE LOCALIZATION IN THE EPENDYMA AND INFUNDIBULAR RECESS IN THE CATFISH CLARIAS BATRACHUS AND ITS PROBABLE SIGNIFICANCE. EXPERIENTIA.32(7):943-944.

[716] SATPUTE,L.R. AGARWAL,S.M. (1974) SEASONAL INFECTION OF CLARIAS BATRACHUS (BLOCK) LYTOCESTUS INDICAIS MOGHE AND PARASITIC EFFECTS ITS HAEMATOLOGY AND HISTOPATHOLOGY. INDIAN J. EX BIOL.12(6):584-586.

[717] SATPUTE,L.R. AGARWAL,S.M. (1974) 'DIVERTICULOSIS' OF THE FISH DUODENUM INFESTED WITH CESTODES. INDIAN J.EXP.BIOL. 12(4):373-375.

[718] SAUVAGE,H.E. (1878)[ON THE ICHTHYOLOGY OF THE OGOOUE.]BULL.SOC. PHILOM.(7)III:90-103(IN FRENCH)

[719] SAUVAGE,H.E. (1880)[STUDIES ON THE FISH FAUNA OF THE OGOOUE.] N.ARC MUS.(2)III:1-56(IN FRENCH.)

[720] SAUVAGE,H.E. (1882)A COLLECTION OF FISHES MADE BY M CHAPER IN THE ASSINIE TERRITORY IVORY COAST.BULL.SOC.Z.FR. 188 :313-325.

[721] SAUVAGE,H.E. (1882) BULL.SOC.Z.FR:.313-325.

[722] SAUVAGE,H.E. (1883)[DESCRIPTIONS OF THE FISH FAUNA OF THE WEST OF ASIA AND MORE PARTICULARLY ON THE FISHES COLLECTED BY MR.CHANTRE DURING HIS TRAVELS IN THAT REGION.]N.MUS.(2)VII.PP1-41.(IN FRENCH)

[723] SAXENA,D.B. (1954)

SYSTEM OF HETEROPNEUSTES FOSSILIS BLOCH. (ACTINOPTERYGII:OSTARIOPHYSI:SILUROIDEA).CURR SCI.23:363.

[724] SAXENA,D.B. (1958)EXTENT OF THE GILL SURFACE IN THE TELEOSTS HETEROPNEUSTES FOSSILIS BLOCH AND CLARIAS BATRACHUS LINN. PROC.NAT.ACAD.SCI.INDIA(B)28: 258-263.

[725] SAXENA,D.B. (1958)CIRCULATION OF BLOOD IN THE RESPIRATORY REGION OF SOME FRESHWATER CATFISHES OF INDIA. PROC. INDIAN ACAD.SCI.28 B III:264-283.

[726] SAXENA,D.B. (1960)ON THE ASPHYXIATION AND INFLUENCE OF CARBON DIOXIDE ON RESPIRATION OF AIR BREATHING FISH, HETEROPNEUSTES FOSSILIS AND CLARIAS BATRACHUS. J.ZOOL.SOC.INDIA,12(1):114-124.

[727] SAXENA,D.B. (1963)COMPARATIVE STUDY OF THE GILL AREA IN SOME FRESHWATER CATFISHES OF INDIA. J.ZOOL.SOC. INDIA.

[728] SAXENA,D.B. (1963)A REVIEW ON ECOLOGICAL STUDIES AND THEIR IMPORTANCE IN THE PHYSIOLOGY OF AIR BREATHING FISHES. ICTHYOLOGICA,II:116-128.

[729] SAXENA,D.B.
CHITRAY,B.B. (1966)
ON THE CORRELATION BETWEEN THE BODY LENGTH WITH GUT AND INTESTINAL LENGTHS OF SOME FRESHWATER FISHES OF INDIA ICHTHYOLOGICAL, KANPUR 5:15-52

[730] SAXENA,P.K. (1966)SOME SALIENT FEATURES IN THE CRANIAL NERVES OF CERTAIN TELEOSTEAN FISHES JAP.J. ICHTHYOL. 14:99-100.

[731] SAXENA,P.K. (1966)CRANIAL NERVES OF THE COMMON CATFISH CLARIAS BATRACHUS LINN. JAP.J.ICHTHYOL. 14:91-98

[732] SCHOONBEE,H.J. (1969)NOTES ON THE FOOD HABITS OF FISH IN LAKE BARBERSPAN,WEST. TRANSVAAL,SOUTH AFRICA. VERH. INTERNAT.VEREIN.LIMNOL.BD. 17:689-701.

[733] SCHURMANNS-STEKHOVEN,J.H(1937). PARASITIC NEMATODA. IN: EXPLOR. PARC NATN. ALBERT MISS. G.F. DE WITTE,(1933-1935),FASC4: 1-40.

[734] SELAPUTIN,G. (1907)[A STUDY OF THE SKELETONS OF SILURIDS AND THE SKULL OF CLARIAS.] MOSKVA,TRD.SRAVNIT-ANATOM. INST.UNIV.,1-5:;-38. VIDE ALSO:MOSKVA,ZAP.UNIV., 22 (IN RUSSIAN)

[735] SEN,T.K. (1975)FURTHER LIGHT ON FRESHWATER FISH FAUNA OF ANDAMAN ISLANDS. SEAFOOD EXPORT J. 7(2):31-33 .

[736] SHAFFI,S.A.
JAFRI,A.K.
KHAWAJA,D.K. (1974)
DISTRIBUTION OF ACID AND ALKALINE PHOSPHATASE ACTIVITY IN THE GASTROINTESTINAL TRACT OF THE FRESHWATER CATFISH,CLARIAS BATRACHUS (LINN.) BROTERIA,43(1-2):39-44.

[737] SHAFFI,S.A.
JAFRI,A.K.
KHAWAJA,D.K. (1974)
ALKALINE PHOSPHATASE ACTIVITY IN THE OVARY OF THE CATFISH,CLARIAS BATRACHUS(LINN.) DURING MATURATION. CURR.SCI.43(2):51.

[738] SHAFFI,S.A.
JAFRI,A.K. (1974)
THE ACTIVITY OF ACID AND ALKALINE PHOSPHATASES IN THE EGGS OF SOME FRESH WATER TELEOSTS.IND.J.FISH. 21(1):296-298.

[739] SHANBHAG,A.B.
NADKARNIS,V.B. (1977)
HISTOLOGICAL AND HISTOCHEMICAL STUDIES ON THE INTERRENAL TISSUE AND CHROMAFFIN CELLS IN 2 FRESH WATER TELEOSTS CHANNA STRIATUS AND CLARIAS BATRACHUS.ZOOLOGISCHER ANZEIGER 198(1-2):109-114.

[740] SHARMA,S. (1968)NEW GLANDULAR STRUCTURE IN THE KIDNEY OF CLARIAS BATRACHUS (LINN.).NATURE, LOND. 217:85.

[741] SHARMA,S. (1969)THE MODES OF VASCULAR SUPPLY IN THE KIDNEYS OF TELEOSTS.ANN.ZOOL.AGRA.5: 111-114

[742] SHARMA,S. (1971)HOMOLOGY OF THE SO-CALLED 'HEAD-KIDNEY' IN CERTAIN INDIAN TELEOSTS. ANN.ZOOL.,AGRA 7(2):19-40.

[743] SHERNI,K.D. JAFRI,A.K. (1975) TOTAL CHOLESTEROL CONTENT OF THE BRAIN AND SPINAL CORD OF SOME FRESH WATER FISHES.IND.J.MED.RES. 63(5):733-735.

[744] SHETTY,H.P.C. (1972)INDUCED BREEDING OF CLARIAS BATRACHUS. FAO AQUACULT.BULL. 5(1):4.

[745] SIDDIQI,M.A. (1966)A COMPARATIVE STUDY OF SOME BIOCHEMICAL CONSTITUENTS IN DIFFERNT TISSUES OF SOME FRESH WATER TELEOSTS.II.TOTAL CHOLESTEROL.SCI.CULT. 32(136-138.

[746] SIDDIQUI,A.H. SIDDIQUI,A.Q. (1969) A COMPARATIVE STUDY OF THE FREE AMINO ACID MAKE-UP OF SIX SPECIES OF FRESHWATER FISHES FROM NORTH INDIA. LABDEU 7(B):141-143.

[747] SIDDIQUI,N. (1975)EFFECT OF FEEDING,SPAWNING AND SIZE ON CHEMICAL CONSTITUENTS OF THE BLOOD PLASMA OF CLARIAS BATRACHUS (LINN) INDIAN J. EXP.BIOL.13(2):203-205.

[748] SIDTHIMUNKA,A. EKURU,P. (1959) [BREEDING AND CULTIVATION OF PLADUK IN PONDJ. THAI FISH.GAZ.12(3):253-259 (IN THAI).

[749] SIDTHIMUNKA, A. PAWAPOOTANON,U. SANGLERT,J. (1968) THE CULTURE OF CATFISH (CLARIAS SPP.) IN THAILAND FAO FISH REP. 44(5):196-204.

[750] SIDTHIMUNKA,A. (1972)THE CULTURE OF PLA DUK(CLARIAS SP).REPORT NO. 12, INLAND FISH. DIV.DEPT.OF.FISH.BANGKOK,THAILAND. PP 17.

[751] SIDTHIMUNKA,A. (1972)PRODUCTION OF CATFISH (CLARIAS SP) AND ITS PROBLEMS IN THAILAND.SEMINAR ON THE PROBLEMS OF INNOVATING AGRICULTURAL TECHNOLOGY IN THAILAND. MAY 1972:17-19.

[752] SIDTHIMUNKA,A. (1973)CULTURE OF CATFISH ON THAILAND FARMS. FISH FARM. INT.1:81-83.

[753] SINGH,H.R. (1969)FURTHER CONTRIBUTIONS TO THE STUDY OF THE TELENCEPHALIC NUCLEAR AREAS OF SOME INDIAN FISHES. PROC.NATN.ACAD.SCI.INDIA 40(SECT.B):99-112.

[754] SINGH,H.R. (1969)ON THE STRUCTURE AND RELATIONS OF THE HABENULA IN SOME TELEOSTS.ZOOL. BEITR. 15: 457-464

[755] SINGH,H.R. (1969)FURTHER CONTRIBUTIONS TO THE STUDY OF THE VALVULA CEREBELLI OF SOME TELEOSTS.PROC. NATN.ACAD.SCI. INDIA 40(SECT.B):16-20

[756] SINGH,H.R. (1970)THE STUCTURE AND RELATIONS OF THE TORUS LONGITUDINALIS IN SOME TELEOSTS ZOOL. BEITR.16:83-91

[757] SINGH,H.R. KHANNA,S.S. (1970) THE CYTOARCHITECTURE AND THE FIBRE CONNECTIONS OF THE OPTIC TECTUM OF SOME TELEOSTS.ZOOL BEITR. 16:129-140

[758] SINGH, B.N. HUGHES,G.M. (1971) RESPIRATION OF AN AIR-BREATHING CATFISH CLARIAS BATRACHUS (LINN.).J.EXP.BIOL.55(2):421-434.

[759] SINGH,S.P. (1971)STUDIES ON THE PHYSIOLOGY OF ALIMENTARY CANAL OF CLARIAS BATRACHUS (LINN.).INDIAN J.ZOOTOMY. 12(2):151-153.

[760] SINGH,H.R. KHANNA,S.S. (1972) ON THE STRUCTURE AND RELATIONS OF THE MESENCEPH-ALIC TEGMENTUM OF THE BRAIN IN SOME TELEOSTS. ZOOL.BEITR.28(1):159-166.

[761] SINGH,B.N. (1976)METABOLIC ENERGY REQUIREMENTS AND ENERGY BUDGETS

OF CERTAIN INDIAN AIR BREATHING FISHES.MARATHWADA UNIV.J.SCI.15(8):3-9.

[762] SINHA,A.L. (1973)LENGTH WEIGHT RELATIONSHIP OF A FRESH WATER CATFISH CLARIAS BATRACHUS.IND.J.ZOOT.14(2): 97-102.
AQUATIC SCI.ABSTRACTS 1969-1976(NOV).

[763] SINHA,A.L. (1973)LENGTH WEIGHT RELATIONSHIP OF A FRESH WATER CATFISH CLARIAS BATRACHUS.IND.J.ZOOT.14(2):97-102.

[764] SINHA,G.M. (1975)A HISTOCHEMICAL STUDY OF THE MUCOUS CELLS IN THE BUCCO-PHARYNGEAL REGION OF FOUR INDIAN FRESHWATER FISHES IN RELATION TO THEIR ORIGIN, DEVELOPMENT, OCCURRENCE AND PROBABLE FUNCTIONS. ACTA. HISTOCHEM 53(2):217-223.

[765] SIRCAR,A.K. (1970)MORPHOLOGY OF THE URINOGENITAL SYSYEM OF SOME SILUROID FISHES. PROC.ZOOL.SOC.CALCUTTA 23: 93-117.

[766] SITASITH,P. (1970)[CULTIVATION OF PLA DUK AT SMUTPRAKARN PROVINCE] THAI FISH.GAZ.23(1):69-71 (IN THAI).

[767] SIVALINGAM,S. (1975)THE BIOLOGY OF CULTIVABLE BRACKISH WATER AND MARINE FIN FISH IN AFRICA. IN:FAO/CIFA SYMPOSIUM ON AQUACULTURE IN AFRICA. CIFA/75/SR 1 PP 8.

[768] SMITH,H.M. (1906)
SEARLE,ALVIN NOTES ON A COLLECTION OF FISHES FROM THE ISLAND OF MINDANAO PHILIPPINE ARCHIPELAGO, WITH DESCRIPTIONS OF NEW GENERA AND SPECIES WASHINGTON D.C.,PROC.BIOL.SOC. 19:73-82

[769] SMITH,H.M. (1934)CONTRIBUTIONS TO THE ICHTHYOLOGY OF SIAM. IX. SIAMESE CLARIID CATFISHES.J.SIAM SOC.9(3):287-325

[770] SMITH,H.M. (1943)THE FRESHWATER FISHES OF SIAM OR THAILAND.U.S.

[771] SOONG,MIN KONG (1950)FISHES OF THE MALAYAN PADI-FIELDS.III.KELI: CAT- NAT.MUS.BULL.188:346-353.
FISHES.MALAYAN NATURE J. (5):86-91.

[772] SORENSEN,W. (1882)[RESEARCH ON TETRODONS(?) AND CLARIAS.] NAT.TIDS.XIV.PP.374-407.

[773] SOULSBY,J.J. (1960)THE MWERU-WANTIPA FISHING RECORD.RHOD.AGRIC.J. 57(4):331-337.

[774] SOULSBY,J.J. (1960)THE MWERU-WENTIPA FISHING RECORD.IN:REPORT ON THE TRAINING CENTRE ON FISHERY SURVEYS FOR THE COUNTRIES OF THE AFRICAN REGION AT TANGA TANGANYIKA. VOLII ESSAYS ON THE STATUS OF FISHERIES DEVELOPMENT IN SELECTED COUNTRIES AND TERRITORIES OF AFRICA. FAO/UN. :141-157.

[775] SOUTHWELL,T. (1939)
LAKE,F. ON A COLLECTION OF CESTODA FROM THE BELGIAN CONGO.ANN.TROP.MED.PARASIT.33:107-123:

[776] SPINAGE,C.A. (1970)CATFISH ON THE MOVE. ANIMALS 13:214-215.

[777] SPINAGE,C.A. (1971)SPECTACULAR MIGRATION OF CATFISH.AFR.WILD.LIFE. 25:70-71.

[778] SRINIVASACHAR,H.R. (1961)THE DEVELOPMENT OF THE SKULL IN CATFISHES. PART III THE DEVELOPMENT OF THE CHONDROCRANIUM IN HETEROPNEUSTES FOSSILIS (BLOCH)(HETEROPNEUSTIDAE) AND CLARIAS BATRACHUS (LINN)(CLARIIDAE). GEGENBAURS JB. 101:373-405.

[779] SRIVASTAVA,C.M. (1961)
SINHA,A.N. CONTRIBUTION TO THE STUDY OF BARBELS OF FISHES. VESTN. CESKOSL.ZOOL. SPOLEC. 25:12-15.

[780] SRIVASTAVA,A.K. (1965)STUDIES ON THE MORPHOLOGY AND PHYSIOLOGY OF THE HEART OF CLARIAS BATRACHUS (LINN.) PART I: THE GROSS ANATOMY OF THE HEART. ZOOL.ANZ. 175:310-316

[781] SRIVASTAVA,G.J. (1967)A NEW SPECIES OF FRESHWATER FISH OF THE GENUS HEMIRAMPHUS CUV. FROM GORAKHPUR, UTTAR PRADESH, INDIA.J.BOMBAY NAT.HIST.SOC.64:93-94.

[782] SRIVASTAVA,A.K. (1968)STUDIES ON THE HEMATOLOGY OF CERTAIN FRESH WATER TELEOSTS.I.ERYTHROCYTES.ANAT.ANZ.123(3):

233-249.

[783] SRIVASTAVA,G.I. (1968)FISHES OF EASTERN UTTAR PRADESH,VISHWAVIDYALAYA PRAKASHAN.BHAIRAVNATH,VARANASI,INDIA.PP163.

[784] SRIVASTAVA,S. SATHYANESAN,A.G. (1971) STRUCTURE OF THE THYROID OF SOME TELEOSTS HAVING ACCESSORY RESPIRATORY ORGANS Z.MIKROSKOP.ANAT.FORSCH.83(2):237-245

[785] SSENTONGO,G.W. MANN,M.J. (1971) ON THE FISH SPECIES OF LAKE BARINGO. FRESHWATER FISH.RES.ORGAN.ANN.REP. 1970:20-27.

[786] SSENTONGO,G.W. (1974)ON THE FISHES AND FISHERIES OF LAKE BARINGO. AFR.J.TROP.HYDROBIOL.FISH.3(2):95-105.

[787] STEINDACHNER,F. (1866)ICICTHYOLOGICAL BULLETIN(IX).JVERH.ZOOL-BOT. GES.WIEN:761-766

[788] STEINDACHNER,F. (1869) SB.AK.WIEN,IX.PP.945-995.

[789] STEINDACHNER,F. (1894)ITHE FISHES OF LIBERIA.JNOTES LEYDEN MUS. XVI:1-96.(IN GERMAN)

[790] STEINITZ,H. (1954)THE DISTRIBUTION AND EVOLUTION OF THE FISHES OF PALESTINE. PUBL.HIDROBIOL.RES.INST.UNIV. ISTANBUL 1B:225-275.

[791] STERBA,G. (1962)FRESHWATER FISHES OF THE WORLD.LONGACRE PRESSLIT LONDON:395-397.

[792] SUNDARARAJ,B.I. GOSWAMI,S.V. (1966) EFFECTS OF MAMMALIAN HYPOPHYSIAL HORMONES , PLACENTAL GONADOTROPHINS,GONADAL HORMONES , AND ADRENAL CORTICOSTEROIDS ON OVULATION AND SPAWNING IN HYPOPHYSECTOMIZED CATFISH HETEROPNEUSTES FOSSILIS (BLOCH). J.EXPTL.ZOOL. 161:287-296.

[793] SYDENHAM,REV.D.H.J. (1975)OBSERVATIONS ON THE FISH POPULATIONS OF A NIGERIAN FOREST STREAM. ZOOL.AFR.89(2):257-272.

[794] SYMES,O.E. (1954)DISCUSSION ON SPECIES OF FISH FOR FISH FARMING. IN:SYMPOSIUM ON AFRICAN HYDROBIOLOGY AND INLAND FISHERIES,1952.CSA.PUB.NO.6:96.

[795] TADROS,G. (1966)ON A NEW CESTODE BOTHRIOCEPHALUS PRUDHDEI SP.NOV. FROM THE NILE CATFISH CLARIAS ANGUILLARIS WITH SOME REMARKS ON THE GENUS CLE STOBOTHRIUM LUHE, 1899,BULL ZOOL SOC. EGYPT.21: 74-88.

[796] TADROS, G. (1968)A REDESCRIPTION OF POLYONCHOBOTHRIUM CLARIAS (WOODLAND,1925) MEGGITT, 1930(BOTHRIOCEPHALIDAE(CESTODA) WITH A BRIEF REVIEW OF THE GENUS POLYONCHOBOTHRIUM DIESING, 1854 AND THE IDENTITY OF THE GENERA TETRACAMPOS WEDL, 1861, SENGA DOLLFUS, 1935, AND ONCHOBOTHRIO YAMAGUTI,1959.J.VET.SCI.U.A.R. 5(1):53-84.

[797] TAIT,C.C. (1967)MASS FISH MORTALITIES.FISH.RES.BULL.ZAMBIA(3): 28-30.

[798] TAIT, C.C. (1967)OBSERVATIONS ON PREDATION BY THE WHITE PELICAN, PELECANUS ONOCROTALUS, AND BY SILURID FISH OF THE GENUS CLARIAS,AND A NOTE ON CORMORANT PREDATION. FISH.RES.BULL. ZAMBIA.(3): 31-32.

[799] TAJON,M.M. KINGSCOTE,A.A. (1970) A STUDY OF THE EFFICACY OF N-TRITYLMORHOLINE AS A MOLLUSCICIDE AGAINST LYMNEA AURICULARIARUBIGINOSA. PHILIPP. J. ANIM. IND. 27:9-16.

[800] TANDON,K.K. SHARMA,C.B. (1966) THE FISH FAUNA OF THE RIVER GHAGGAR IN THE VICINITY OF CHANDIMANDIR. RES.BULL. PUNJAB UNIV. N.S. SCI. 16:333-337.

[801] TANDON,K.S. JOSHI,B.D. (1973)

WISS,Z. STUDIES ON THE PHYSIOPATHOLOGY OF BLOOD OF FRESH WATER FISHES INFECTED WITH TWO NEW FORMS OF TRYPANOSOMES. ZOOL.185(304):207-221.

[802] TANDON,R.S. (1974)
JOSHI,B.D. EFFECT OF TRYPANOSOME INFECTION ON BLOOD GLUCOSE LEVELS OF SOME FRESH WATER TELEOSTS.J.INL.FISH. SOC.INDIA.6:81-82.

[803] TANDON,R.S. (1975)
JOSHI,B.D. STUDIES ON THE PHYSIOLOGY OF BLOOD OF SOME FRESH WATER FISHES OF INDIA PART1 NORMAL BLOOD GLUCOSE AND LACTIC ACID LEVELS.J.INL.FISH.SOC. INDIA, 7:1-6.

[804] TANDON,R.S. (1976)
CHANDRA,S. CYCLIC CHANGES IN SERUM CHOLESTEROL LEVELS OF FRESH WATER CATFISH CLARIAS BATRACHUS.ZEITSCHRIFT FUER TIERPHYSIOLOGIE,TIERERNAEHRUNG UND FUTTERMIT TELKUNDE .36(4):179-183.

[805] TANDON,R.S., (1977)
CHANDRA,S. CYCLIC CHANGES IN BLOOD UREA LEVELS OF FRESH WATER CATFISH CLARIAS BATRACHUS.ZEITSCHRIFT FUER TIER PHYSIOLOGIE + FUTTERMITTELKUNDE.38(4):211-214.

[806] TARTARINOV,L.P. (1960)[EVOLUTION OF THE APPARATUS DIVIDING BLOOD STREAMS IN VERTEBRATES HEART.] ZOOL.ZH. 39:1218 -1231 (IN RUSSIAN WITH ENGLISH SUMMARY).

[807] TAYLOR,J. (1830)ON THE RESPIRATION ORGANS AND AIR BLADDER OF CERTAIN FISHES OF THE GANGES. GLEANINGS IN SCIENCE.2:169-176.

[808] THAKUR,N.K. (1974)
PAL,R.N.
KHAN,H.A. EMBRYONIC AND LARVAL DEVELOPMENT OF HETEROPNEUSTES FOSSILIS. J.INL.FISH.SOC.INDIA. 6:33-44.

[809] THAKUR,N.K. (1974)
PAL,R.N.
KHAN,H.A. EMBRYONIC AND LARVAL DEVELOPMENT OF HETEROPNEUSTES FOSSILIS.J.INL.FISH.SOC.INDIA.6:33-44.

[810] THOMAS,J.D. (1958)THREE NEW DIGENETIC TREMATODES, EMOLEPTALEA PROTEROPORA N.SP.(CEPHALOGINIMIDAE: CEPHALOGONIMINAE) PHYLLODISTOMUM SYMMETRORCHIS N.SP. AND PHYLLODISTOMUM GHANENSE N.S.P. (GORGODERIDAE: GORGODERINAE) FROM WEST AFRICAN FRESHWATER FISHES. PROC. HELMINTH.SOC. WASH. 25(1):1-8.

[811] THOMAS,E.I. (1963)STUDIES ON THE HEART OF CLARIAS MAGUR (C&V.). AGRA.UNIV.J.RES.12(3):59-67.

[812] THOMAS, J.D. (1966)ON THE BIOLOGY OF THE CATFISH CLARIAS SENEGALENSIS,IN A MAN-MADE LAKE IN THE GHANAIAN SAVANNA WITH PARTICULAR REFERENCE TO ITS FEEDING HABITS. PROC. ZOOL. SOC.LOND. 148:476-514.

[813] THYS VAN ANDERNAUDE,D. (1968)ADDENDUM TO THE 'FRESHWATER FISHES FERNANDOPO' ON SOME FISHES COLLECTED BY COL.J.J.SCHEEL. REV. ZOOL BOT. AFR. 78(1/2): 123-128.

[814] TILAK, R. (1963)RELATIONSHIPS BETWEEN THE OSTEOCRANIUM AND WEBERIAN APPARATUS IN TWO INDIAN CATFISHES OF THE GENUS CLARIAS (SILURIDAE).COPEIA,(4):623-9.

[815] TILAK,R. (1967)STUDIES ON THE OSTEOLOGY OF THE NEMATOGNATHINE GIRDLE IN RELATION TO TAXONOMY. J.ZOOL.SOC. INDIA 19:101-110.

[816] TOBOR,J.G. (1973)THE FISH AND FISHERIES OF THE LOWER YOBE RIVER IN THE NORTH-WESTERN SECTOR OF LAKE TCHAD. ANN.REPORT DEPT.FISH.NIGERIA:70-75.

[817] TUNGSANGA,S. (1963)
MENASVETA,P.
SIDTHIMUNKA,A. INDUCED SPAWNING IN CATFISH (CLARIAS MACROCEPHALUS

GUNTHER) BY PITUITARY HORMONE INJECTION. PROC. INDO-PACIF. FISH.COUN. 10(11):205-213.

[818] TOOTS,H (1968)REPORT ON TEST FISHING AND AND ANGLING IN CLEVELAND DAM FROM 1962-1968. INTERNAL REPORT FISHERIES RESEARCH CENTER(HENDERSON),MIN.OF AGRIC R&SS, RHODESIA.PP19.

[819] TOOTS,H. (1969)REPORT ON TEST FISHING AT MARANDELLAS TOWN COUNCIL NYAMBUIA AND NYAKAMBIRA DAMS(9/4/69 - 11/4/69.INTERNAL REPORT,FISHERIES RESEARCH CENTER (HENDERSON),MIN.OF AGRIC,R&SS,RHODESIA.PP7.

[820] TOOTS,H. (1969)REPORT ON NGONDOMA DAM: SURVEY AND TESTFISHING, 15-18 APRIL,1969,INTERNAL REPORT FISHERIES RESEARCH CENTER(HENDERSON), MIN.OF AGRIC.R&SS, RHODESIA.PP4

[821] TOOTS,H. (1970)FIRST REPORT ON SILALABUHWA DAM.INTERNAL REPORT FISHERIES RESEARCH CENTER(HENDERSON),MIN.OF AGRIC R&SS,RHODESIA.PP3.

[822] TOOTS,H. (1971)SILALABUHWA DAM(SECOND REPORT).INTERNAL REPORT FISHERIES RESEARCH CENTER(HENDERSON),MIN.OF AGRIC R&SS,RHODESIA.PP9.

[823] TOOTS,H. (1971)NGONDOMA DAM(SECOND REPORT).INTERNAL REPORT FISHERIES RESEARCH CENTER(HENDERSON),MIN.OF AGRIC R&SS,RHODESIA.PP10.

[824] TOOTS,H. (1972)REPORT ON CHIMWEMBWE DAM.INTERNAL REPORT FISHERIES RESEARCH CENTER(HENDERSON), MIN.OF AGRIC.R&SS,RHODESIA.PP8.

[825] TOOTS,H. (1973)FARM DAMS FOR FISH PRODUCTION.RHOD.AGRIC.J.71(6) 133-138.

[826] TOOTS,H. (1974)REPORT ON TESTFISHING IN GWENORO & IMPALI DAMS. INTERNAL REPORT DEPT OF NAT.PARK &WLF.MAN,MIN.OF LANDS,RHODESIA.PP8.

[827] TOOTS,H. (1974)REPORT ON OBSERVATIONS AND TESTFISHING IN PINDI PARK DAM IN THE BANKET AREA.INTERNAL REPORT , DEPT.OF NAT.PARK &WLF.MAN.MIN.OF LANDS,RHODESIA. PP10.

[828] TOOTS,H. (1974)REPORT ON TESTFISHING AND OBSERVATIONS OF NGAMO AND WHITE WATERS DAMS.INTERNAL REPORT , DEPT.OF NAT.PARK &WLF.MAN.MIN.OF LANDS,RHODESIA.PP7.

[829] TOOTS,H. (1974)REPORT ON TESTFISHING IN NGONDOMA DAM(22/4/74 TO 27/4/74.INTERNAL REPORT , DEPT.OF NAT.PARK &WLF. MAN. MIN.OF LANDS,RHODESIA.PP8.

[830] TORTONESE,E. (1934)[ON ANOTHER AFRICAN SPECIES OF THE GENUS CLARIAS] BULL.MUS.ZOOL.ANAT.COMP. TORINO 44:231-239. (IN ITALIAN).

[831] TREWAVAS,E. (1936)DR KARL JORDAN'S EXPEDITION TO SOUTH WEST AFRICA AND ANGOLA: THE FRESHWATER FISHES. NOVIT.ZOOL. TRING 40:63-74

[832] TREWAVAS,E. (1962)FISHES OF THE CRATER LAKES OF THE NORTH-WESTERN CAMEROONS. BONN. ZOOL. BEITR. 13:146-192.

[833] TRIPATHI,S.D. SHARAF,R.R. (1974) AN ARTIFICIAL FEED FOR AIR BREATHING FISHES. LABDEV,PART B :LIFE SCIENCES.12(4):151-152.

[834] TRIPATHI,S.D. SHARAF,R.R. (1974) AN ARTIFICIAL FEED FOR AIR BREATHING FISHES.LABDEV PART B:LIFE SCIENCES.12(4):151-152.

[835] TRISTRAM,H.B. (1884)THE SURVEY OF WESTERN PALESTINE.FRESHWATER FISHES. LONDON. :162-177.

[836] TUBANGUI,M.A. (1931)TREMATODE PARASITES OF PHILIPPINE VERTEBRATES. III.FLUKES FROM FISH AND REPTILES.PHILIPP.J.SCI. 44:417-423.

[837] TWEEDIE,M.W.F. (1952)NOTES ON MALAYAN FRESH-WATER FISHES.NO.3 THE ANABANTOID FISHES.NO.4.SOME NEW AND INTERESTING RECORDS.NO.5.MALAYAN NAMES.BULL.RAFFLES MUS. SINGAPORE NO.24:63-95.

[838] VAILLANT,L. (1895)[ON THE NATURE OF THE DORSAL SPINES OF SILURIDS

AND CYPRINOIDS C.R.JAC. CXXI :909-911.(IN FRENCH)

[839] VAILLANT,L. (1895)[ON THE TERRESTRIAL HABITS OF THE AFRICAN SILURID CLARIAS LAZERA, CUVIER AND VALENCIENNES.]BULL. MUS.NH. PP.271-272. (IN FRENCH)

[840] VAILLANT,L. (1897)[THE DISTRIBUTION OF THE SILUROIDES IN SPACE AND TIME.]REV.SCI.(4)VII(685-689.(IN FRENCH)

[841] VAILLANT L. (1904)[A FEW REPTILES,BATRACHIANS AND FISH OF UPPER TONKIN.] BULL.MUS. PARIS :297-301 (IN FRENCH)

[842] VANDEL,A. (1964)[CAVE BIOLOGY. THE BIOLOGY OF CAVE DWELLING ANIMALS.] PARIS :1-619. (IN FRENCH)

[843] VAN DER LINGEN,M.I. (1962)REPORT ON LAKE MCILWAINE. F.R.C.INVEST. REPORT. (MIMEO,REPORT)DEPT.R&SS.MIN.AGRIC.RHODESIA.

[844] VAN DER LINGEN,M.I. (1962)REPORT ON NGESI LAKE,NO. I,II&III.F.R.C.INVEST. REPORT.(MIMEO.REPORT)DEPT.R&SS.MIN.AGRIC. RHODESIA.

[845] VAN DER LINGEN,M.I. (1962)REPORT ON LAKE SEBAKWE,NO.I&II.F.R.C.INVEST. REPORT.(MIMEO,REPORTS)DEPT.R&SS.MIN.AGRIC. RHODESIA.

[846] VAN DER LINGEN,M.I. (1965)SOME PROBLEMS OF BREEDING BIOLOGY OF FISH IN IMPOUNDED WATERS.KARIBA RESEARCH SYMPOSIUM.LAKE KARIBA FISHERIES RESEARCH INSTITUTE,KARIBA:97-103

[847] VAN DER WAAL,B.C.W. (1970)[BREEDING AND PRODUCTION OF CLARIAS GARIEPINUS.] DEPT. OF NATURE CONSERVATION, LOWVELD FISHERIES STATION, 1969-1970, MARBLE HALL,TVL.RSA. PROJECT NO. 5.PP30-43.(IN AFRIKAANS)

[848] VAN DER LINGEN,M.I. (1971)A CASE STUDY OF LAKE KARIBA.(MIMEO.REPORT)DEPT. NAT.PARKS.&WLDF.MAN.SALISBURY,RHODESIA.PP 35.

[849] VAN DER WAAL,B.C.W. (1972)[INVESTIGATIONS INTO ASPECTS OF THE ECOLOGY, BREEDING AND PRODUCTION OF CLARIAS GARIEPINUS (BURCHELL)1822.]M.SC. THESIS,AFRIKAANS UNIVERSITY OF THE RAND,RSA.(IN AFRIKAANS)

[850] VAN DER WAAL,B.C.W. (1974)OBSERVATIONS ON THE BREEDING HABITS OF CLARIAS GARIEPINUS(BURCHELL).J.FISH.BIOL.6(23-27.

[851] VAN DER WAAL,B.C.W. SCHOONBEE,H.J. (1975) AGE AND GROWTH STUDIES OF CLARIAS GARIEPINUS (BURCHELL):CLARIIDAE) IN THE TRANSVAAL,SOUTH AFRICA. J.FISH BIOL.7:227-233.

[852] VAN SCHOOR,D.J. (1968)SURVEY OF THE FISHES IN THE ORANGE AND VAAL RIVERS FROM ALIWAL NORTH TO DOUGLAS.NEWS. BULL.ZOOL. SOC. S.AFR. 9(2):8-9.

[853] VARGHESE,T.J. DEVARAJ,K.V. SATYANARAYANA-RAO,G.P. (1973) FOOD OF THE JUVENILES OF THE CATFISH CLARIAS BATRACHUS(LINN.).J.INLAND FISH.SOC.INDIA, BARRACKPORE,5:78-81.

[854] VASSILIADES,G. (1979)[A NEW PARACAMALLANUS (NEMATODA:CAMALLANIDSE), FROM A FISH CLARIIDAE OF SANGALKAN (SENEGAL).] ANNLS. PARASIT. LUM. COMP. 45:441-447. (IN FRENCH

[855] VELASQUEZ,C.C. (1959)STUDIES ON PARASITES OF PHILIPPINE EDIBLE FISHES. 1.FAMILY CLINOSTOMIDAE IN OPHICEPHALUS STRIATUS BLOCH,CLINOSTOMUM PHILIPPINENSIS SP.NOV. WITH NOTES ON THE DEFINITIVE HOSTS. PHILIPP.J.SCI. 88(3): 263-278.

[856] VERMA,S.R. TYAGI,M.P. (1974) A COMPARATIVE HISTOLOGICAL STUDY OF THE STOMACH OF A FEW TELEOST FISHES(PART1).MORPHOL. JAHRB. 120(2):244-253.

[857] VERRIER,M.L. (1928)[STUDIES ON THE EYES AND VISION OF FISH.] BULL. BIOL.FRANCE BELG. SUPPL. 11:1-222 (IN FRENCH)

[858] VERRIER,M.L. (1928)[THE STRUCTURE OF THE EAR OF CLARIAS BATRACHUS AND AMEIRUS NEBULOSUS LESUEUR, WITH RESPECT TO HABITAT AND BIOLOGY OF THESE TWO SILURIDS.] BULL. SOC.ZOOL.FR. 52:581-594 (IN FRENCH).

[859] VINCIGUERRA,D. (1880)[FIRST CONTRIBUTION TO THE FISH FAUNA OF THE

ISLAND OF BORNEO.JANN.MUS.GENOV.XVI.PP161-182. (IN ITALIAN)

[860] VINCIGUERRA,D. (1893)DI ALCUNI PESCI RACCOLTI NEL PAESE DEI SOMALI DALL INGRE L. BRICHETTI-RIBECCHI.ANN.MUS.GENOVA, (2)XIII:448-455.

[861] VINCIGUERRA,D. (1895)ESPLORAZIONE DEL GIUBA E DEI SUOI AFFLUENTI COMPIUTA DAL CAP.V.BOTTEGO DURANTE GLI ANNI 1892-1893 SOTTO GLI AUSPICII DELLA SOCIETA GEOGRAPHICA ITALIANA.PESCI.ANN.MUS.GENOVA(2)XV: 21-60.

[862] VINCIGUERRA,D. (1897)PESCI RACCOLTI DA DON EUGENIO DEI PRINCIPI RUSPOLI DURANTE L ULTIMO SUO VIAGO NELLE REGIONI DEI SOMALI E DEI GALLA. ANN.MUS.GENOVA (2)XVIII:24-31.

[863] VOLKER,A. (1968)ECOLOGICAL INVADERS TILTING THE SCALES. SCI.NEWS. 94(17): 423-425.

[864] WANICHAKORN,P. (1957)THE CULTIVATION OF CATFISH(CLARIAS SPP.)THAI FISH.GAZ.10(1):39-49.

[865] WANICHAKORN,P. (1962)RAISING OF YOUNG CATFISH IN FLOATING BASKETS. THAI.FISH.GAZ.15(3):271-276.

[866] WANICHAKORN,P. (1964)CULTURE OF CATFISH (CLARIAS SPP.).FISH CULTURE EXTENSION PAMPHLET,FISHERIES DEPT.BANGKOK, THAILAND

[867] WARD,J.W. DAVIS,H.C. (1974) SOME ASPECTS OF THE HEMATOLOGY OF CLARIAS BATRACHUS(LINN.).Q.J.FLA.ACAD.SCI.

[868] WASAWO,D.P.S. (1964)SOME PROBLEMS OF UGANDA SWAMPS.PROC.PAP.I.U.C.N. PUBLS.N.S. 4:196-204.

[869] WATANABE,K.O. (1971)PHYSICAL CHARACTERISTICS AND CHEMICAL COMPOSITION OF FRESH BREAM,MUDSUCKER,TIGER-FISH,AND BARBEL FROM LAKE KARIBA.FISH.RES.BULL.ZAMBIA(5):163-173.

[870] WEBER,M. (1897)[BULLETINS ON THE FAUNA OF SOUTH AFRICA.I. BULLETINS OF FRESHWATER FAUNA OF SOUTH AFRICA]. ZOOL.JAHRB.SYST. X:135-199.

[871] WEDL,C. (1861)[ON THE HELMINTH FAUNA OF EGYPT].(2ABT.)SBR,DT. AKAD. LAND WISS. WIEN, 44:463-482. (IN GERMAN)

[872] WEIR,J.S. (1969)STUDIES ON CENTRAL AFRICAN PANS. 3. FAUNA AND PHYSIO-CHEMICAL ENVIRONMENT OF SOME EPHEMERAL POOL HYDROBIOLOGIA 33:93-116.

[873] WEIR,J.S. (1972)DIVERSITY AND ABUNDANCE OF AQUATIC INSECTS REDUCED BY INTRODUCTION OF THE CLARIAS GARIEPINUS TO POOLS IN CENTRAL AFRICA.BIOL. CONSERV.4(3):169-175.

[874] WELCOMME,R.L. (1964)THE ECOLOGY OF THE FISHES OF A SMALL RIVER OF THE LAKE VICTORIA BASIN.ANN.REP.EAFRO.(1962/ 1963):32-35.

[875] WELCOMME,R.L. (1969)THE BIOLOGY AND ECOLOGY OF THE FISHES OF A SMALL TROPICAL STREAM.J.ZOOL.LOND.158:485-529.

[876] WELCOMME,R.L. (1970)A DESCRIPTION OF CERTAIN INDIGENOUS FISHING METHODS FROM SOUTHERN DAHOMEY. AFR.J.TROP. HYDROBIOL. FISH. 1:129-140.

[877] WELMAN,J.B. WORTHINGTON,E.B. (1943) THE FOOD OF THE CROCODILE (CROCODILUS NILOTICUS) PROC.ZOOL.SOC.LOND.113:108-112.

[878] WELMAN,T.B. (1948)PRELIMINARY SURVEY OF THE FRESHWATER FISHERIES OF NIGERIA.LAGOS,GOVERNMENT PRINTER,NIGERIA.

BOT.AFR.LIX,3-4(329-363.

[879] WELSCH,U. STORCH,V. (1969) [ON THE FINE STRUCTURE OF TASTE BUDS OF THE CATFISH (CLARIAS BATRACHUS (L.) AND KRYPTOPTERUS BICIRRHIS(C.&V.)).JZ.ZELLFORSCH.MIKROSK.ANAT.100 :552-559.(IN GERMAN).

[880] WHITEHEAD,P.J.P. (1959)THE ANADROMOUS FISHES OF LAKE VICTORIA.REV.ZOOL.

[881] WHYTE,S .A . (1975)DISTRIBUTION, TROPHIC RELATIONSHIPS AND BREEDING HABITS OF THE FISH POPULATIONS IN A TROPHIC LAKE BASIN.(LAKE BOSUMTWI, GHANA) J.ZOOL.117(1):25-56.

[882] WILLEM,V. (1951)[CONTRIBUTIONS TO THE STUDY OF THE RESPIRATORY ORGANS OF THE TELEOSTS: CLARIAS AND HETEROBRANCHUS].BULL.MUS.HIST.NAT.BELG.27(36)1-8 (IN FRENCH)

[883] WILLIAMS,N.V. (1960)A REVIEW OF THE KAFUE RIVER FISHERY.RHOD.AGRIC.J. 57(1):86-92.

[884] WILLIAMS,N.V. (1960)A REVIEW OF THE KAFUE RIVER FISHERY.IN:REPORT ON THE TRAINING CENTRE ON FISHERY SURVEYS FOR THE COUNTRIES OF THE AFRICAN REGION AT TANGA, TANGANYIKA. VOLII ESSAYS ON THE STATUS OF FISHERIES DEVELOPMENT IN SELECTED COUNTRIES AND TERRITORIES OF AFRICA. FAO/UN.:128-141.

[885] WILLMER,E.N. (1934)SOME OBSERVATIONS ON THE RESPIRATION OF CERTAIN TROPICAL FRESHWATER FISHES. J.EXP.BIOL.11: 283-306.

[886] WOODLAND,W.N.F. (1924)ON A NEW SPECIES OF THE CESTODARIAN GENUS CARYOPHYLLAEUS FROM AN EGYPTIAN SILUROID. PROC. ZOOL. LOND. 1924(2):529-532.

[887] WOODLAND,W.N.F. (1925)ON SOME REMARKABLE NEW MONOTICELLIA-LIKE AND OTHER CESTODES FROM SUDANESE SILUROIDS. Q.JL.MICROSC.SCI. 69(4):703-729.

[888] WOODLAND,W.N.F. (1937)SOME CESTODES FROM SIERRA LEONE. I.ON WENYONIA LONGICUADA SP.N. AND PROTEO CEPHALUS BIVITELLATU SP.N. PROC. ZOOL. SOC. LOND. 1936(4): 931-937

[889] WOODLAND,W.N.F. (1937)SOME CESTODES FROM SIERRA LEONE. II A NEW CARYOPHYLLAEID, MARSYPOCEPHALUS AND POLYONCHOBOTHRIUM. PROC.ZOOL.SOC. LOND. 1937(2):189-197.

[890] WORTHINGTON,E.B. (1933)III. FISHES (OTHER THAN CICHLIDAE) OF LAKE BANGWEULU AND ADJOINING REGIONS WITH DESCRIPTIONS OF THREE SPECIES.ANN.MAG.NAT.HIST.SER.10,V. XII,1933:34-52.

[891] WORTHINGTON,E.B. (1933)THE FISHES OF LAKE NYASA (OTHER THAN CICHLIDAE). PROC.ZOOL.SOC.LONDON:285-316.

[892] YANNI,M. (1961)LIPIDES AND WATER CONTENT OF THE TISSUES OF CLARIAS LAZERA.Z.VERGL.PHYSIOL.45,(1):50-55.

[893] YANNI,M.H. (1961)EFFECT OF EXCESS GLUCOSE ON THE CARBOHYDRATES, WATER AND FAT CONTENTS OF THE TISSUES OF CLARIAS LAZERA.Z.VERGL.PHYSIOL.45:227-232.

[894] YANNI,M. (1961)STUDIES ON CARBOHYDRATE CONTENT OF THE TISSUES OF CLARIAS LAZERA.Z.VERGL.PHYSIOL.45(1):56-60.

[895] YANNI,M. (1962)EFFECT OF STARVATION ON THE CONTENTS OF THE WATER AND LIPIDES OF TISSUES OF CLARIAS LAZERA.Z.VERGL. PHYSIOL.45(4):390-395.

[896] YANNI,M. (1962)EFFECTS OF STARVATION ON CARBOHYDRATE CONTENT OF TISSUES AND ON RELATIVE WEIGHTS OF ORGANS OF CLARIUS LAZERA.Z.VERFL.PHYSIOL. 45(3):315-21.

[897] YANNI,M. (1963)THE EFFECT OF FASTING ON THE TOLERANCE OF CLARIUS LAZERA TO EXCESS GLUCOSE Z.VERGL.PHYSIOL. 46(3):314-20

[898] YANNI,M. (1964)THE ASSIMILATION OF FRUCTOSE AND GALACTOSE IN THE DIFFERENT TISSUES OF CLARIAS LAZERA.Z.VERGL. PHYSIOL.49(2):130-137.

[899] YANNI,M. (1964)EFFECT OF INSULIN AND EXCESS GLUCOSE ON THE CARBOHYDRATES,WATER AND FAT CONTENTS OF THE TISSUES OF CLARIAS LAZERA.Z.VERGL.PHYSIOL.48:624-631.

[900] YANNI,M. (1972)DEPOSITION AND UTILIZATION OF OLIVE OIL BY DIFFERENT TISSUES OF CLARIAS LAZERA WITH REFERENC E TO THE DEGREE OF SATURATION OF DEPOT FAT. ZOOL SOC.EGYPT BULL.24:31-41.

[901] YEH,L.S. (1960)ON A COLLECTION OF CAMALLANID NEMATODES FROM

FRESHWATER FISHES IN CEYLON.J.HELMINTH.34(1-2):
107-116.

Plate 1. Dentition of upper jaw of *Clarias gariepinus* that would previously have been named *Clarias mossambicus* (from the Hartbeespoort Dam, Republic of South Africa).



Plate 2. Dentition of upper jaw of typical *Clarias gariepinus* (from Hartbeespoort Dam, Republic of South Africa).

Plate 3. Liver of dissected *Clarias gariepinus* showing (arrow) hepatic blood vessel leaving main body of liver to link to the lateral liver lobe. Note orange coloured flesh - the flesh of the African catfish has a venison type texture (from Sand-River Lake, Swaziland).

Plate 4. Lateral liver lobe exposed externally on *Clarias gariepinus*. Entry made laterally immediately posterior to skull: note tubercles on skull plates (from Sand-River Lake, Swaziland).